

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA REG. NO. 279-3426

2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN Alabama, Arkansas, Delaware, Georgia, Kentucky, Maine, Maryland, Massachusetts, Michigan, New Jersey, North Carolina, Ohio, Rhode Island, South Carolina, Tennessee, Virginia, and West Virginia.

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

CROP	PEST	RATE OF APPLICATION
Soybeans	Kudzu bug or Bean Plataspid	4.0 fl oz / acre

DIRECTIONS FOR USE: Through coverage is essential to achieve control. Apply by ground or air using sufficient spray volume to obtain full coverage of foliage. Multiple applications may be required

Soybeans - (21 Day PHI): Use a minimum spray volume of 10 gallons by ground and 2 gallons by air.

RESTRICTIONS:

Do not apply more than 24 fluid ounces product (0.15 lbs active ingredient) per acre per season.

Do not make applications less than 7 days apart.

Do not graze or harvest treated soybean forage, straw, or hay for livestock feed.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED MUSTANG MAXX INSECTICIDE LABEL (EPA REG. NO. 279-3426) MUST BE FOLLOWED.

THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

2(ee) 2013-05-23

Mustang and FMC -FMC Trademarks
©2013 FMC Corporation