

**RESTRICTED USE PESTICIDE**

**Toxic to fish and aquatic organisms.**

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**



EPA Reg. No. 279-3426

**2(ee) Recommendation**

**FOR DISTRIBUTION AND USE  
IN ALL STATES \*  
with the exception of Idaho, Oregon, Texas, and  
Washington**

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING**

CROP	PEST	RATE OF APPLICATION
<p><b>Berries Crop Group including:</b> Blackberry; Loganberry; Red and Black Raspberry; Blueberry, highbush and lowbush; Currant; Elderberry; Gooseberry; Huckleberry; and cultivars and/or hybrids of these</p> <p><b>Stone Fruit Crop Group including:</b> Apricot; Cherry (sweet and tart); Nectarine; Peach; Plum; Plumcot; Prune (fresh)</p> <p><b>Grapes</b></p>	<p><b>Vinegar Flies (adults) (<i>Drosophila spp.</i>)</b></p> <p><b>Spotted Wing Drosophila</b></p>	<p><b>4.0 fl oz/A (0.025 lb ai/A)</b></p>

**DIRECTIONS FOR USE:** Through coverage is essential to achieve control. Apply by ground or air using sufficient spray volume to obtain full coverage of foliage. Multiple applications may be required

**BERRIES CROP GROUP - (1 Day PHI):** Use a minimum spray volume of 20 gallons by ground and 2 gallons by air.

**STONE FRUIT CROP GROUP - (14 Day PHI):** For ground application, use a minimum spray volume of 20 gallons per acre for concentrate spray or 100 gallons per acre for dilute sprays. For aerial application, use a minimum spray volume of 10 gallons per acre. Do not apply as a ULV spray. Do not feed or allow livestock to graze on cover crops from treated orchards.

**GRAPES - (1 Day PHI):** Use a minimum spray volume of 10 gallons by ground and 2 gallons by air.



FMC Corporation  
Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

2(ee) 2014-02-18

Mustang and FMC -FMC Trademarks  
©2014 FMC Corporation

**RESTRICTIONS:**

Do not apply more than 24 fluid ounces product (0.15 lbs active ingredient) per acre per season.

Do not make applications less than 7 days apart.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED MUSTANG MAXX INSECTICIDE LABEL (EPA REG. NO. 279-3426) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

\*Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.



FMC Corporation  
Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

2(ee) 2014-02-18

Mustang and FMC -FMC Trademarks  
©2014 FMC Corporation