

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee)
AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**



EPA Reg. No. 84876-1-279

2(ee) Recommendation
FOR DISTRIBUTION AND USE ONLY IN California
THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)
IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

CROP	PEST	RATE OF APPLICATION
Grape	Powdery Mildew (<i>Erysiphe necator</i>)	18.3 – 20.4 fl oz / acre.

DIRECTIONS FOR USE: Apply in a preventive spray schedule. Make the first application of Fracture Fungicide before bloom and continue applications using spray intervals of up to 14 days in low to moderate disease pressure. Use full labeled rates (20.5 – 24.4 fl oz/A) when disease pressure is severe.

Alternate applications with another labeled effective fungicide with a different mode of action.

Do not make more than five foliar applications of Fracture Fungicide per crop season.

Do not make more than two sequential applications of Fracture Fungicide before alternating to a labeled fungicide with a different mode of action.

Do not harvest until 1 day after last application.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED Fracture Fungicide LABEL (EPA REG. NO. 84876-1-279) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

