

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA REG. NO. 84876-1-279

<p>2(ee) Recommendation</p> <p>FOR DISTRIBUTION AND USE ONLY IN Florida, Georgia, North Carolina, and South Carolina</p>
<p>IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING</p>
<p>This Product Bulletin Contains New or Additional Directions For Use Which Are Recommended By FMC Corporation Under FIFRA Section 2(ee)</p>

CROP	PEST	RATE OF APPLICATION
Tomato	Southern Blight <i>Sclerotium rolfsii</i>	24.4 – 36.6 fl oz/A (0.4-0.6 lb ai/A)
<p>DIRECTIONS FOR USE: Apply sprays of Fracture® Fungicide following soil applications of Fontelis® Fungicide or other fungicide labeled for Southern Blight control in tomato. Apply Fracture® Fungicide as a directed spray to the soil. Ensure thorough coverage of the lower stems and soil surface surrounding the plants.</p>		
<p>RESTRICTIONS: Do not make more than two sequential applications of Fracture® Fungicide before alternating to a labeled fungicide with a different mode of action. Do not make more than 5 applications of Fracture® Fungicide per harvest cycle.</p>		
<p>PHI: 1 Day</p>		
<p>ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED FRACTURE® FUNGICIDE (EPA REG. NO. 84876-1-279) LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.</p>		



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

FMC and Fracture are trademarks of FMC Corporation.
Fontelis® Fungicide is a trademark of DuPont Crop Protection
©2017 FMC Corporation
Fracture_2ee_Tomato Southern Blight_03-24-2017