



**24(c) Special Local Need Label
For Use in Fallow or Soybeans with
Reduced Rotation to Sugarcane**

**FOR DISTRIBUTION AND USE ONLY
IN LOUISIANA**

**EPA Reg. No. 279-9560
EPA SLN No. LA-140004**

This label for Authority Maxx Herbicide is valid until September 8, 2019, or until withdrawn, canceled or suspended.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CROPS	WEED	USE RATE Oz by wt./Acre (lb a.i.)*			
		% Organic Matter	Coarse	Medium	Fine
Fallow and Soybean	Morningglory, entireleaf	1.0 – 2.0	5 - 6 (0.2) – (0.25)	6 - 7 (0.25) – (0.29)	7 - 8 (0.29) – (0.33)
	Morningglory, ivyleaf		2.0 – 4.0	6 - 7 (0.25) – (0.29)	7 - 8 (0.29) – (0.33)
	Morningglory, red				
	Nutsedge, yellow				

Refer to the AUTHORITY MAXX section 3 label for information on soil types under the COARSE, MEDIUM, and FINE categories.

*Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range. Do not apply to soils with pH >7.5.

Fallow and Soybean Application

Authority Maxx may be applied to fallow or soybean fields to be planted in the fall to sugarcane at 5.0 to 9.6 ounces per acre. Make application of Authority Maxx at least 60 days prior to the planting of sugarcane. Apply either by air in a minimum of 5 gallons of spray per acres or by ground equipment in a minimum of 15 gallons of spray per acre.

Apply to newly prepared seedbeds as a preemergence prior to weed emergence. If weeds are present at time of application it is recommended to add a registered burndown herbicide to improve postemergence control. Observe all directions for use, precautions and restrictions on the label of the burndown herbicide.

Restrictions

- Do not apply less than 60 days prior to planting sugarcane.
- Do not apply more than 9.6 oz (0.396 pound active) total per twelve-month period. The twelve-month period is considered to begin upon the initial AUTHORITY MAXX herbicide application.
- Do not apply more than one application of AUTHORITY MAXX herbicide per growing season.
- Do not apply more than 0.375 lb a.i. of the sulfentrazone active total per acre per twelve-month period.
- Do not apply to soils with pH >7.5.
- Do not harvest sugarcane for 14 months after the AUTHORITY MAXX herbicide application.
- Do not apply to coarse soils classified as sand, which have less than 1% organic matter.

CROP ROTATIONAL INTERVALS

Shown below are the minimum intervals in days from the time of AUTHORITY MAXX herbicide application until AUTHORITY MAXX herbicide treated soil may be replanted. Refer to the complete label for crop rotation intervals for all other crops.

CROP	INTERVAL (Days)
Sugarcane	60

**MANUFACTURER’S CONDITIONS AND RISK OF USE FOR FEDERAL
INSECTICIDE FUNGICIDE AND RODENTICIDE ACT (FIFRA) SECTION 24(c)
REGISTRATION**

NOTICE: This Section 24(c) label will only be utilized by growers located in the State of Louisiana. **SUCH GROWER USERS AGREE FMC CORPORATION WILL NOT BE HELD RESPONSIBLE FOR FAILURE TO PERFORM AND/OR CROP RESPONSE AND OR YIELD LOSSES RESULTING FROM THE USE OF AUTHORITY MAXX HERBICIDE AS A PRE-EMERGE TREATMENT TO SUGARCANE.** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability on the commercial Authority Maxx Herbicide section 3 label and this 24(c) label before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

Authority and **FMC** - Trademarks of FMC Corporation
©2014 FMC Corporation. All Rights Reserved.

090814