

# SUPPLEMENTAL LABELING



EPA Reg. No. 71711-31

|             |       |     |             |
|-------------|-------|-----|-------------|
| TOLFENPYRAD | GROUP | 21A | INSECTICIDE |
|             | GROUP | 39  | FUNGICIDE   |

**For Use On: Leafy Greens (Crop Subgroup 4-16A)**

amaranth, Chinese; amaranth, leafy; aster, Indian; blackjack; cat's whiskers; cham-chwi; cham-na-mul; chervil, fresh leaves; chipilin; chrysanthemum, garland; cilantro, fresh leaves; corn salad; cosmos; dandelion, leaves; dang-gwi, leaves; dillweed; dock; dol-nam-mul; ebolo; endive; escarole; fameflower; feather cockscomb; Good King Henry; huauzontle; jute, leaves; lettuce, bitter; lettuce, head; lettuce, leaf; orach; parsley, fresh leaves; plantain, buckhorn; primrose, English; purslane, garden; purslane, winter; radicchio; spinach; spinach, Malabar; spinach, New Zealand; spinach, tanier; Swiss chard; violet, Chinese, leaves; cultivars, varieties, and hybrids of these commodities

**Use Pattern: Aerial Application**

This supplemental labeling expires 07/15/2025 and must not be used or distributed after this date.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for TORAC® Insecticide before applying. Use of TORAC Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for TORAC Insecticide.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

## APPLICATION RATE CHART FOR TORAC INSECTICIDE

| Leafy Greens (Crop Subgroup 4-16A) |  |   |
|------------------------------------|--|---|
| Pest                               | Rate/Acre  | Use Directions  |
| Leafhoppers                        | 14.0 to 21.0 fl oz/acre<br>(0.14 to 0.21 lb ai/acre) | <ul style="list-style-type: none"><li>• Aerial Application: Use a minimum of 5 gallons of water per acre.</li><li>• Ground Application: Use a minimum of 20 gallons of water per acre.</li></ul> <b>USE RESTRICTIONS</b> <ul style="list-style-type: none"><li>• Do not apply until at least fourteen (14) days after emergence or after transplanting to allow time for root establishment. This period of time should be extended if conditions at time of emergence or transplanting are not favorable to crop growth.</li><li>• Do not apply more than 42.0 fluid ounces (0.42 lb ai) per acre per crop cycle.</li><li>• Do not make more than 2 applications per crop cycle.</li><li>• Allow at least 14 days between applications.</li><li>• Do not make more than 4 applications per year.</li><li>• Preharvest Interval (PHI): 1 day</li></ul> <b>USE RECOMMENDATIONS - DISEASES</b> <ul style="list-style-type: none"><li>• Begin applications prior to onset of disease.</li><li>• Use of an agriculturally approved nonionic surfactant at 0.25% v/v may improve disease control.</li><li>• If weather conditions remain conducive to disease development, apply another registered fungicide product with a different mode of action 7 to 10 days later.</li><li>• Consult local extension recommendations or your agricultural consultant for information specific to your area.</li></ul> |
| Aphids (excluding lettuce aphid)   | 17.0 to 21.0 fl oz/acre<br>(0.17 to 0.21 lb ai/acre) |   |
| Powdery mildew                     | 21.0 fl oz/acre<br>(0.21 lb ai/acre)                 |   |
| Thrips                             |  |   |
| Armyworms*                         |  |   |
| Corn earworm*                      |  |   |
| Cutworm species*                   |  |   |
| European corn borer*               |  |   |
| Flea beetle*                       |  |   |
| Imported cabbageworm*              |  |   |
| Tobacco budworm*                   |  |   |
| Tomato hornworm*                   |  |   |
| Whitefly*                          |  |   |
| Downy mildew*                      |  |   |

\*suppression only

\*suppression only

©2022 Nichino America, Inc. Torac is a registered trademark of Nichino America, Inc. The Fujiichi-mark is a trademark of Nihon Nohyaku Co., Ltd.



Nichino America, Inc.  
4550 Linden Hill Road, Suite 501  
Wilmington, DE 19808  
888-740-7700