



2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA

DUPONT™ EXIREL® INSECT CONTROL

EPA Reg. No. 352-859

DIRECTIONS FOR USE FOR OVERHEAD CHEMIGATION APPLICATION IN POTATOES IN THE STATE OF TEXAS

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

Refer to main label for general use precautions and restrictions and to supplemental label for overhead chemigation application instructions.

DIRECTIONS FOR USE

Crop	Application Method	Target Pest	EXIREL® RATE		Last Application (days to Harvest)	REI (Hours)
			Lb. ai per acre	Fluid ounces product per acre		
Potatoes	Overhead Chemigation	Colorado potato beetle†	0.033 - 0.088	5 - 13.5	7	12
		Beet armyworm	0.044 - 0.088	7 - 13.5		
		European corn borer				
		Potato tuberworm*††				
		Yellow striped armyworm	0.065 - 0.011	10 - 17		
		Potato flea beetle* §	0.088 - 0.134	13.5 - 20.5		
		Green peach aphid*				
		Potato aphid* §				
		Potato psyllid				

Minimum application interval between treatments is 5 days.

Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR™ or cyantraniliprole containing products per crop; this is the total of seed piece treatment (potato), soil treatment, and foliar treatment.

* - For best performance use with an effective adjuvant. See "Use of Adjuvants" section.

§ - Suppression only. Use as part of an effective control program. Rotate with products with different modes of action.

† - Colorado potato beetle resistance management - Do not apply EXIREL® more than twice to a generation of Colorado potato beetle or within any 30 day period. Application(s) to the next generation of Colorado potato beetle must be with an effective product with a different mode of action. Do not apply EXIREL® for Colorado potato beetle control if any cyantraniliprole containing product was used at-plant either as a soil applied or seed treatment.

†† - Potato Tuberworm: EXIREL® may be applied at rates of 7 to 13.5 fl oz/A to control potato tuberworm. Begin application when field scouting indicates the presence of tuberworm adults and/or larvae. Potato tuberworm often have overlapping generations so repeat applications of EXIREL® may be needed based on scouting. Avoid treating successive generations with the same mode of action. It is important to protect the crop just prior to harvest when foliage starts to senesce. Use the higher rate of EXIREL® when tuberworm pressure is high. Failure to adequately control potato tuberworm larvae prior to crop senescence or vine kill increases the risk of tuber damage. Foliar sprays alone, by air or ground, may not provide adequate control of larvae in the mid to lower crop canopy. For best results, apply via overhead chemigation or integrate chemigation applications into the foliar spray program. For chemigation applications, apply in 0.1 to 0.2 acre inches of water and add MSO at 12 to 16 fl oz/acre. See supplemental label on CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS for instructions on overhead sprinkler chemigation.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the DuPont product(s) covered by this Bulletin. DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by DuPont on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2015 E.I. du Pont de Nemours and Company, 1007 Market Street, Wilmington, DE 19898