

## 2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



### DUPONT™ EXIREL® INSECT CONTROL

EPA Reg. No. 352-859

### FOR CONTROL OF CARROT WEEVIL IN ROOT VEGETABLES (CROP GROUP 1B EXCEPT SUGAR BEET), AND FLORENCE FENNEL IN THE STATE OF TEXAS

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ Exirel® insect control must be used only in accordance with directions on this bulletin or in separately published DuPont labeling.

DuPont Exirel® Insect control is recommended for control of carrot weevil in root vegetables (crop group 1B, except sugar beet) and Florence fennel.

Apply Exirel® insect control at 13.5-20.5 fl oz/A as a foliar spray using properly calibrated ground spray equipment. Initial application should be made at first sign of adult weevil presence.

For best results use the highest labeled rate of Exirel® (20.5 fl oz/A). Under moderate to high pest pressure, an additional application of Exirel® may be needed to maintain weevil populations below damaging levels. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for this pest.

Thorough spray coverage of is essential to achieve best results.

Read and follow the protection of pollinators section of the federal label when using during bloom.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

#### RESTRICTIONS

Do not apply a total of more than 0.4 lbs ai/A of CYAZYPYR® or cyantraniliprole containing products per calendar year.

The minimum retreatment interval is 5 days.

The pre-harvest interval is 1 day.

The field re-entry interval is 12 hours.

#### IMPORTANT NOTICE

**BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

**Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT  
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A