

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ EXIREL® INSECT CONTROL

EPA Reg. No. 352-859

CONCENTRATION BASED RATES FOR CONTROL OF ASIAN CITRUS PSYLLID, CITRUS LEAFMINER, AND DIAPREPES ROOT WEEVIL ADULTS IN YOUNG RESET OR SOLID-SET CITRUS TREES IN THE STATE OF FLORIDA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application. All applicable directions, restrictions and precautions on the EPA registered label must be followed. This recommendation must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for DuPont™ Exirel® insect control before applying.

DuPont™ Exirel® insect control must be used only in accordance with directions in this bulletin or in separately published DuPont labeling.

APPLICATION RATES, TIMING, METHOD AND PEST CONTROLLED

DuPont™ Exirel® insect control is recommended for control of Asian citrus psyllid, citrus leafminer and *Diaprepes* root weevil adults in young reset or solid-set citrus trees (crop group 10-10) up to 24 months old.

Application Rate: Apply DuPont™ Exirel® insect control at 16-20.5 fluid ounces per 100 gallons of water. For best performance, use of an adjuvant such as horticultural spray oil at 1 gallon per 100 gallons of spray volume (1% v/v) is recommended. Do not apply more than 0.133 lb. of CYAZYPYR® per acre per application (20.5 fl. oz./A Exirel®) or a total of more than 0.4 lb of CYAZYPYR® or other cyantraniliprole containing products per acre per calendar year.

Application Timing: DuPont™ Exirel® insect control should be timed to protect new flush. Applications should be made early during the feather flush stage when the new flush is no more than 40% expanded and before heavy insect damage is present.

Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees and density of foliage to ensure adequate coverage to wet the foliage. The amount of treatment solution per acre will vary based on the size of trees being treated and planting density. When using a rate of 20.5 fl. oz. DuPont™ Exirel insect control per 100 gallons of water, do not apply more than 100 gallons of spray solution per acre.

Application Equipment: Applications can be made with a handgun or hoop boom sprayer to young reset or solid-set citrus trees in sufficient spray volume to wet all plant foliage.

Read and follow the Protection of Pollinators section of the federal label when using during bloom.

SPRAY PREPARATION

Acidification of the spray tank: If the pH of the spray tank after all products have been added and mixed is above pH 8, adjust to pH 8 or less using a registered acidifying agent. If the spray tank pH is 8 or less no adjustment is necessary. Spray tanks of pH 8 or less can be held up to 8 hours before spraying. Do not store the spray mixture overnight in the spray tank.

RESTRICTIONS

Do not apply DuPont™ Exirel® insect control to the soil or through drip irrigation systems.

Do not apply a total of more than 0.4 lbs. ai/A of CYAZYPYR® or cyantraniliprole containing products per calendar year whether applications are foliar or soil-applied.

For resistance management purposes, applications of DuPont™ Exirel® insect control or other Group 28 insecticides should be used in rotation with other non-group 28 soil-applied or foliar insecticides.

The minimum retreatment interval is 7 days.

Expiration Date: 10/31/2023

The PHI (pre-harvest interval) is 1 day.

The REI (re-entry interval) is 12 hours.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

As of November 1, 2017, the USEPA registration for DuPont™ EXIREL® insect control was sold to FMC by DuPont.

Copies of the Exirel insect control product label are available at www.fmccrop.com/grower/

**EXIREL, CYAZYPYR and FMC are trademarks of FMC Corporation or an affiliate. ©2018 FMC Corporation.
2929 Walnut Street, Philadelphia, PA 19104**

R-4115 110118 10-31-18

Expiration Date: 10/31/2023