



2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA

DUPONT™ VERIMARK® INSECT CONTROL

EPA Reg. No. 352-860

FOR CONTROL OF STRIPED CUCUMBER BEETLE IN CUCURBIT VEGETABLES (CROP GROUP 9) IN THE STATE OF NEW YORK

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe; consult the State or Tribal agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ Verimark® insect control must be used only in accordance with directions on this bulletin or in separately published DuPont labeling.

DuPont™ Verimark® insect control is recommended for the control of striped cucumber beetle in cucurbit vegetables (EPA crop group 9).

APPLICATION METHOD, RATES AND TIMINGS

Tray Drench

Apply DuPont™ Verimark® insect control at 13.5 fluid ounces per acre (0.176 lb. ai/A) as a transplant tray drench no earlier than 72 hours prior to planting in the field.

Drip Chemigation

Apply DuPont™ Verimark® insect control at 10 fluid ounces per acre (0.130 lb ai/A) as a drip chemigation application.

DuPont™ Verimark® insect control must be applied uniformly in the root zone to ensure effective control. Drip tape must be placed directly underneath a single row of plants to ensure DuPont™ Verimark® insect control is applied in the root zone.

Allow 1 – 3 days for DuPont™ Verimark® insect control to be translocated into the aerial portions of the plant and to fully protect transplants following an at-planting application. Allow 2 – 5 days for DuPont™ Verimark® insect control to fully protect the plants following a drip application.

SPRAY PREPARATION

Adjusting the pH of the application solution: The pH of the application solution should be between pH 4 and 6. (See “Acidification of Application Solution” section of the container label.)

SOIL APPLICATION DIRECTIONS

DuPont™ Verimark® insect control may be applied as a transplant tray drench by growers or commercial transplant producers no earlier than 72 hours prior to planting in the field. For drip chemigation applications after planting, apply at the specified rates before insect populations reach locally determined action thresholds. Application via drip chemigation is more effective if made during the first half of the crop, approximately 2-3 weeks after transplanting to allow the root system time to develop in order to ensure adequate uptake of Cyazapyr®. Two applications using a 10-14 day interval may be required for effective control.

Refer to the SOIL APPLICATIONS section of the container label for additional use and application guidance.

Read and follow all label directions for the specified application methods as described on the Federal Label.

RESTRICTIONS

Minimum application interval between drip chemigation treatments is 10 days.

Do not apply more than 13.5 fluid ounces per acre (0.176 lb. ai/A) of DuPont™ Verimark® insect control at planting.

Do not apply more than 2 drip chemigation applications of DuPont™ Verimark® insect control per crop.

Do not make more than one drip application of DuPont™ Verimark® insect control per crop if an at plant application of Verimark was made.

Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® or cyantraniliprole containing products per calendar year whether the application is made to the soil or foliarly.

Re-entry interval (REI) is 4 hours.

PHI (pre-harvest interval) is 1 day.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2018 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A