

Cyflumetofen	Group	25	Acaricide
--------------	-------	----	-----------

Supplemental Label

Nealta®

Miticide

For use in Citrus Fruit and Tree Nuts

This supplemental label expires December 31, 2024, and must not be used or distributed after this date.

Active Ingredient:

cyflumetofen: (2-methoxyethyl(*RS*)-2-(4-*tert*-butylphenyl)-2-cyano-3-oxo-3-(α,α,α -trifluoro-*o*-tolyl)propionate)..... 18.7%

Other Ingredients:..... 81.3%

Total: 100.0%

Nealta® miticide contains 1.67 pounds of cyflumetofen per gallon.

EPA Reg. No. 7969-336
CAUTION/PRECAUCION

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Nealta® miticide** container label, EPA Reg. No. 7969-336, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Nealta** before applying.
- Use of **Nealta** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Nealta**.

Product Information

Nealta is a suspension concentrate formulation containing the active ingredient cyflumetofen, a member of the beta-ketonitrile class of chemistry.

When used as directed, **Nealta** provides knockdown and residual control of tetranychid mites on citrus fruit and tree nuts when used according to the label. **Nealta** is a highly active contact miticide on egg, nymph, and adult stage tetranychid mites. **Nealta** is not systemic and has no translaminar activity, thorough coverage of plant surfaces is necessary for effective control.

Nealta is compatible with most biological control products.

Refer to the **Nealta** product container label for **Pests Controlled**, IPM, and **Resistance Management** information.

Application Instructions

Refer to the **Nealta** container label for additional application instructions and restrictions.

Crop-specific Application Instructions

Crop	Pest	Product Use Rate per Application (fl ozs/A)	Minimum Time from Application to Harvest (PHI) days
Citrus Fruit (crop subgroup 10-10A) Calamondin Citron Citrus hybrids Mediterranean mandarin Orange, sour Orange, sweet Satsuma mandarin Tachibana orange Tangelo Tangerine (mandarin) Tangor Trifoliate orange and cultivars, varieties, and/or hybrids of these	See Pests Controlled under Product Information section	13.7 to 17.1	
Citrus Fruit (crop subgroup 10-10B) Australian desert lime Australian finger-lime Australian round lime Brown river finger lime Kumquat Lemon Lime Mount white lime New Guinea wild lime Russell River lime Sweet lime Tahiti lime and cultivars, varieties, and/or hybrids of these	See Pests Controlled under Product Information section	13.7	7
Citrus Fruit (crop subgroup 10-10C) Grapefruit Japanese summer grapefruit Pummelo Tangelo Uniq fruit and cultivars, varieties, and/or hybrids of these			

(continued)

Crop-specific Application Instructions *(continued)*

Citrus Fruit (crop subgroup 10-10A) (crop subgroup 10-10B) (crop subgroup 10-10C) *(continued)*

For maximum knockdown and residual control, apply **Nealta[®] miticide** at the first sign of mite infestation, and before the population increases, according to locally recommended treatment threshold levels in developing mite populations.

Apply **Nealta** in at least 100 gallons of spray solution per acre. For **small nonbearing trees**, apply in at least 50 gallons of spray solution per acre. Applying **Nealta** at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

Restrictions

DO NOT make **Nealta** applications at intervals shorter than 14 days.

DO NOT make more than 2 applications per season per crop when applying at 13.7 fl ozs. **DO NOT** make more than 1 application per season per crop when applying at 17.1 fl ozs.

Resistance Management. DO NOT make more than one **Nealta** application before using an effective miticide with a different mode of action.

DO NOT apply more than 27.4 fl ozs/A (0.36 lb cyflumetofen/A) per season per crop.

Maximum crop seasons per year: 1.

Crop-specific Application Instructions *(continued)*

Crop	Pest	Product Use Rate per Application (fl ozs/A)	Minimum Time from Application to Harvest (PHI) days
<p>Tree Nuts (crop group 14-12) African nut-tree Almond Beechnut Brazil nut Brazilian pine Bunya Bur oak Butternut Cajou nut Candlenut Cashew Chestnut Chinquapin Coconut Coquito nut Dika nut Ginkgo Guiana chestnut Hazelnut (filbert) Heartnut Hickory nut Japanese horse-chestnut Macadamia nut Mongongo nut Monkey-pot Monkey puzzle nut Okari nut Pachira nut Peach palm nut Pecan Pequi Pili nut Pine nut Pistachio Sapucaia nut Tropical almond Walnut, black Walnut, English Yellowhorn and cultivars, varieties, and/or hybrids of these</p>	<p style="text-align: center;">See Pests Controlled under Product Information section</p>	<p style="text-align: center;">13.7 to 17.1</p>	<p style="text-align: center;">7</p>

(continued)

Crop-specific Application Instructions *(continued)*

Tree Nuts (crop group 14-12) *(continued)*

For maximum knockdown and residual control, apply **Nealta[®] miticide** at the first sign of mite infestation, and before the population increases, according to locally recommended treatment threshold levels in developing mite populations.

Apply **Nealta** in at least 100 gallons of spray solution per acre. For **small nonbearing trees**, apply in at least 50 gallons of spray solution per acre. Applying **Nealta** at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

Restrictions

DO NOT make **Nealta** applications at intervals shorter than 14 days.

DO NOT make more than 2 applications per season per crop when applying at 13.7 fl ozs. **DO NOT** make more than 1 application per season per crop when applying at 17.1 fl ozs.

Resistance Management. DO NOT make more than one **Nealta** application before using an effective miticide with a different mode of action.

DO NOT apply more than 27.4 fl ozs/A (0.36 lb cyflumetofen/A) per season per crop.

Maximum crop seasons per year: 1.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF. 1108

Nealta is a registered trademark of BASF.

© 2023 BASF Corporation
All rights reserved.

007969-00336.20230419.NVA 2022-04-0390-0192
Supersedes: NVA 2021-04-390-0184

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

 **BASF**
We create chemistry