



Bayer Environmental Science
A Division of Bayer CropScience, LP
5000 CentreGreen Way, Suite 400
Cary, NC 27513
1-866-99BAYER (1-866-992-2937)

Cimarron Plus® Herbicide

Suppression of Yucca and Juniper with individual plant treatments

EPA Reg. No. 432-1572

Expiration: December 31, 2024

Product Bulletin

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label. Always read this 2(ee) bulletin and the entire label on the product container before using the product.

Bayer will not be responsible for losses or damages resulting from the use of these products in any manner not specifically instructed by Bayer. User assumes all risks associated with non-label use.

Suppression of Yucca and Juniper in Pasture and Rangeland

Apply Cimarron Plus® Herbicide in combination with triclopyr 4 ester in the spring through the fall prior to frost. Make application as an Individual Plant Treatment (IPT). Thorough coverage of all foliage and stems is necessary to optimize results. Spray canopy from all sides to wet but no to the point of dripping. Treatments may be made with equipment such as back pack sprayers, ATV, or hand sprayers. Use an adjustable conejet nozzle with an orifice size of X6 or X12 or equivalent. The application volume required will vary with the height and density of the brush and the application equipment used. For best results with juniper species, treat before plants reach 4 feet in height. Variation in weather (moisture and temperature extremes), plant physiological condition, soil type, and extend of root system will determine treatment effectiveness.

Application Use Rates

Apply Cimarron Plus® Herbicide at a rate of 1 ounce plus one gallon of triclopyr 4 ester plus a spray adjuvant and water to make 100 gallons of spray solution.

Spray Adjuvants

Crop oil concentrate (COC), Modified Seed Oil (MSO) or Modified Seed Oil/Organosilicone (MSO/OS) are the preferred adjuvants. Include 1 to 2 gallons of spray adjuvant per 100 gallons of spray solution. A dye may be added to the tank to help mark plants that have been sprayed.

IMPORTANT

Refer to the Cimarron Plus® Herbicide and any tank-mix partner labels for information on adjuvants; weed species controlled; additional use directions; rotations crop intervals and precautions. The most restrictive provision on any label should apply.

Do not apply more than 2.0 ounces of Cimarron Plus® Herbicide per acre per year as a result of broadcast or repeat applications.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any herbicide product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*