





**Table 1. Application Rates and Intervals for Dollar Spot on Turf**

Disease	Rate (fl.oz./1,000 sq.ft.)	Rate (fl.oz./Acre)	Application Interval (days)	Instructions
Dollar spot ( <i>Sclerotinia homoeocarpa</i> )	0.4 to 0.5	17.4 to 21.8 (0.452 to 0.567 lb. a.i./A)	14	Apply as a preventative treatment prior to or in the very early stage of disease development. Use the highest rate during periods of prolonged favorable disease conditions. Do not apply more than two consecutive applications of KABUTO® FUNGICIDE SC or other Group 7 containing fungicides before rotating to a registered fungicide with a different mode of action. Do not apply the next application of KABUTO® FUNGICIDE SC within 28 days of the previous two consecutive KABUTO® FUNGICIDE SC applications. Do not apply more than 3.2 fl.oz./1,000 sq.ft. per year or 140 fl.oz. per acre per year on turf. This is equal to 8 applications at the low rate (0.452 lb. a.i./A/application) or 6 applications at the high rate (0.567 lb. a.i./A/application). In the State of New York, do not apply more than 3 applications per year at the high rate of 0.567 lb. a.i./A/application (1.70 lb. a.i./acre/year) on turf.
Leaf Spot/Melting Out ( <i>Drechslera Poae</i> )	0.5	21.8 (0.567 lb. a.i./A)	14	

**Table 2. Dilution Table for Applications on Turf**

fl.oz. of KABUTO® FUNGICIDE SC per 100 gallons of spray solutions			
Use Rate (fl.oz. per 1,000 sq.ft.)	Spray Volume 2 gallons per 1,000 sq.ft.	Spray Volume 3 gallons per 1,000 sq.ft.	Spray Volume 4 gallons per 1,000 sq.ft.
0.4	20 fl.oz./100 gal	13.3 fl.oz./100 gal	10 fl.oz./100 gal
0.5	25 fl.oz./100 gal	16.7 fl.oz./100 gal	12.5 fl.oz./100 gal

**Table 3. Application Rates and Intervals for Spring Dead Spot on Turf**

Disease	Rate (fl.oz./1,000 sq.ft.)	Rate (fl.oz./Acre)	Application Interval (days)	Instructions
Spring Dead Spot ( <i>Ophiophaerella korrae</i> , <i>Ophiophaerella herpotricha</i> , <i>Ophiophaerella narmari</i> )	Application Program A 3.2	Application Program A 140 (3.65 lb. a.i./A)	Application Program A None	Apply as a preventative treatment either as a single treatment (Program A) or two treatments (Program B) beginning one month prior to dormancy to southern-type turf grasses, or when soil temperatures have declined to 70°F (21°C). Apply in a minimum spray volume of 2 gallons/1,000 sq.ft. (88 gallons/Acre) and immediately water in with 1/8" irrigation. Program B Use lower rates where damage to turf from spring dead spot is less noticeable and/or more tolerable (i.e. on higher cut turf such as tees and fairways). Use higher rates where damage to turf from spring dead spot is not tolerable (i.e. greens). Do not apply more than 3.2 fl.oz./1,000 sq.ft. (3.65 lb. a.i./A) of this product to turf grass per year. This use is prohibited in the state of New York.
	Application Program B  1st application: 0.5 to 1.6  2nd application: 0.5 to 1.6	Application Program B  1st application: 22 to 70 (0.56 to 1.825 lb. a.i./A)  2nd application: 22 to 70 (0.56 to 1.825 lb. a.i./A)	Application Program B  14-28 days	

**STORAGE AND DISPOSAL**

DO NOT contaminate water, food or feed by storage or disposal.  
**PESTICIDE STORAGE:** Store in original container, in a secured, dry, cool place separate from fertilizer, food, and feed. Avoid cross-contamination with other pesticides.  
 (cont. on next column)

**STORAGE AND DISPOSAL (cont.)**

**PESTICIDE DISPOSAL:** Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Nonrefillable container (equal to or less than 5 gallons). DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**LIMITED WARRANTY AND DISCLAIMER**


**IMPORTANT:** Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this LIMITED WARRANTY AND DISCLAIMER in their entirety and without exception. If the terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under abnormal conditions such as weather, presence of other materials, or the manner or use of application, etc. Such factors and conditions are beyond the control of the manufacturer, and **BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use, subject to the inherent risks described below. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OF HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THE PRODUCT.** The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this LIMITED WARRANTY AND DISCLAIMER in any manner.

Kabuto® is a registered trademark of Ishihara Sangyo Kaisha, Ltd.   
 Active ingredient made in India.



Employee-Owned  
**DISTRIBUTED BY**  
**PBI/GORDON CORPORATION**  
 P.O. BOX 860350  
 SHAWNEE, KANSAS 66286  
 PBIGordonTurf.com

642/3-2021 AP021820  
 EPA REG. NO. 71512-23-2217

**ATTENTION:** This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.