

**SPECIAL LOCAL NEED
24(C)LABELING**

**DuPont Crop
Protection**

**DUPONT™ SENTRALLAS®
HERBICIDE
90 DAY PLANTBACK RESTRICTION TO SOYBEANS AFTER A
SPRING APPLICATION TO
WINTER WHEAT TO CONTROL
BROADLEAF WEEDS INCLUDING CHICKWEED IN THE STATE
OF MARYLAND**

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MARYLAND

DUPONT™ SENTRALLAS® HERBICIDE

**EPA REG. NO. 352-897
EPA SLN NO. MD-18-0001
EXPIRATION DATE: 12/31/2023**

**FOR 90 DAY PLANTBACK RESTRICTION TO SOYBEANS
AFTER A SPRING APPLICATION TO WINTER WHEAT TO
CONTROL BROADLEAF WEEDS INCLUDING CHICKWEED IN
THE STATE OF MARYLAND**

PRODUCT INFORMATION

SENTRALLAS® Herbicide is used for selective postemergence control or suppression of broadleaf weeds in wheat (winter, spring and durum), oats, and barley not underseeded with legumes or grasses. SENTRALLAS® contains two active ingredients formulated as oil dispersion. SENTRALLAS® is to be mixed with water and applied as a uniform broadcast spray early postemergence to the crop, and to the main flush of actively growing broadleaf weeds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this label and the entire label on the product container before using this product.

SENTRALLAS® Herbicide may be used on winter wheat for control of select broadleaf weeds including chickweed in the state of Maryland.

Apply DuPont™ SENTRALLAS® after the crop is in the 2-leaf stage, but before the flag leaf is visible. SENTRALLAS® controls only those weeds that have germinated. Annual broadleaf weeds must be past the cotyledon stage, actively growing.

Do not apply to crops underseeded to legumes or grasses as injury to forage may result.

Applications of SENTRALLAS® to winter wheat that is stressed by severe weather conditions, drought (including low levels of subsoil moisture), near freezing temperatures prior to, at, and following time of application, low fertility, water-saturated soil, disease, or insect damage, May result in crop injury and reduced weed control.

For product information call 1-888-6-DUPONT

© 2017 E. I. du Pont de Nemours and Company, DuPont Agricultural Products, Wilmington, Delaware 19898

USE RATES

Use SENTRALLAS® at 7 to 14 fl oz/A for infestations of those weeds listed under the "WEEDS CONTROLLED/SUPPRESSED" section of the label affixed to the container. Consult table for use product rates required to control/suppress listed weeds.

Specific Use Restrictions for Wheat: Do not make more than two applications of SENTRALLAS® per crop season and do not exceed 14 fl oz/A when making any single application or 17.8 fl oz/A for the entire season to wheat.

GRAZING/HARVESTING RESTRICTIONS

Do not graze or harvest the treated forage within 7 days of application.

Do not harvest treated hay within 30 days of application.

Do not harvest treated grain within 45 days of application.

CROP ROTATION/PLANT-BACK RESTRICTIONS

Soybeans may be planted 90 days after the application of SENTRALLAS® herbicide in the state of Maryland. For all other plant-back restrictions see the label affixed to the container.

SPRAY ADJUVANTS

Include a spray adjuvant with applications of SENTRALLAS®. In addition, an ammonium nitrogen fertilizer may be used.

See TANK MIXTURES for additional information on adjuvant recommendations for certain tank mixtures. Consult your Ag dealer or applicator, local DuPont fact sheets and technical bulletins prior to using an adjuvant system. If another herbicide is tank mixed with SENTRALLAS® select adjuvants authorized for use with both products. Products must contain only EPA exempt ingredients.

NONIONIC SURFACTANT (NIS)

- Apply 0.06 to 0.25% v/v (0.5 to 2 pt per 100 gal of spray solution).
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

CROP OIL CONCENTRATE (COC) - PETROLEUM OR MODIFIED SEED OIL (MSO)

- Apply at 1% v/v (1 gal per 100 gal spray solution). MSO adjuvants may be used at 0.5% v/v if specified on local DuPont product literature or service policies.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

AMMONIUM NITROGEN FERTILIZER

- Use 2 qt/A of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 lb/A of a spray-grade ammonium sulfate (AMS). Use 4 qt/A UAN or 4 lb/A AMS under arid conditions.
- See TANK MIXTURES With Liquid Nitrogen Fertilizer for instructions on using fertilizer as a carrier in place of water.

IMPORTANT

BEFORE USING SENTRALLAS® , READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by DuPont. User assumes all risks associated with such non-directed use.

This product may not be available in all states. See your local DuPont retailer or representative for details and availability in your state.