



# EsplAnade<sup>®</sup>

## F

## HERBICIDE

For Preemergent Weed Control in Conifer and Hardwood Production Areas

**ACTIVE INGREDIENT:**

Indaziflam (CAS No: 950782-86-2) ..... 19.05%

**OTHER INGREDIENTS:** ..... 80.95%

**TOTAL:** ..... 100.00%

EPA Reg. No. 432-1517

Contains 1.67 pounds of indaziflam per gallon

**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

See Back Panel for First Aid Instructions and Booklet for Complete Precautionary Statements and Directions for Use.

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY**

Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-800-331-2867

Nonrefillable Container

Net Contents

**2.5 Gallons**

84488305

84401528B 160419AV5

## FIRST AID

<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything to an unconscious person.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eyes open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>

**For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577**

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**Note to physician:** No specific antidote is available. Treat symptomatically.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, applicators and other handlers must wear:

- long-sleeved shirt and long pants
- shoes plus socks
- waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### ENGINEERING CONTROLS STATEMENTS:

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY RECOMMENDATIONS:

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean water mark. Do not contaminate water when disposing of rinseate or washwater. This product may impact water through spray drift or runoff. Follow directions for use to avoid spray drift and runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential of this product entering water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

**Groundwater Advisory:** This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory:** This pesticide may impact water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

#### DIRECTIONS FOR USE

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**Read the entire label before using this product**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**SHAKE CONTAINER WELL BEFORE USING.**

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Waterproof gloves

## PRODUCT INFORMATION

Esplanade F Herbicide is a selective, preemergent, alkylazine herbicide for control of grasses and broadleaf weeds in conifer and hardwood production areas.

Esplanade F Herbicide controls weeds by reducing the emergence of seedlings through inhibition of cellulose biosynthesis (CB Inhibitor). Necrosis or yellowing may also be observed if the herbicide is applied to herbaceous tissue such as leaves and green stems of susceptible plants. The herbicide needs to be activated prior to weed germination for most effective control. For maximum activity against germinating weeds, Esplanade F Herbicide requires rainfall (minimum 0.25 inches) within several weeks after application to activate the herbicide. Esplanade F Herbicide has minimal post-emergent activity and generally does not control weeds that have emerged. A labeled post-emergent herbicide may be mixed with Esplanade F Herbicide to control existing weeds. Esplanade F Herbicide does not control tubers, rhizomes, and woody vegetation.

Esplanade F Herbicide can be applied to terrestrial conifer and hardwood production areas that contain areas of casual water of a temporary nature as a result of surface water collecting in equipment wheel ruts or in other depressions created by management activities. Esplanade F Herbicide may be applied by ground or aerial equipment (Helicopter only).

The level of weed control is dependent on many variables including soil texture, moisture, temperature, weed species present, the amount of weed seed present in the soil, and the crop canopy.

Weed control activity may be reduced if the application is made to soil covered in heavy crop or weed debris that prevents a uniform distribution of the product reaching the soil.

## USE RESTRICTIONS

- Do not apply Esplanade F Herbicide to soil where standing water is present except as specified on this label.
- Do not apply this product to frozen or snow covered soil.
- Do not contaminate water intended for irrigation and domestic use.
- Do not exceed 7 fl oz/A of Esplanade F Herbicide in a single application.
- Do not exceed 10 fl oz/A of Esplanade F Herbicide within a calendar year or in a 12-month period from the previous application.
- Do not apply Esplanade F Herbicide through an irrigation or chemigation system.
- For aerial applications, do not apply Esplanade F within 25 feet of lakes, reservoirs, rivers, streams, marshes, ponds, estuaries and commercial fish ponds. Refer to the Spray Drift Management section of this label for more information.
- Not for sale, sale into, distribution and/or use in Nassau and Suffolk counties of New York State.
- Do not use in hardwood production areas in California.

## APPLICATION INFORMATION

Apply Esplanade F Herbicide with a properly calibrated sprayer according to the manufacturer's directions and check periodically to be certain that the equipment is working properly prior to each use. Uniform application is essential for satisfactory weed control. Avoid overlap. Shut off spray booms while starting, turning, slowing, or stopping to avoid off-target application.

For ground application, use 10-100 gallons of spray volume per acre to ensure adequate coverage. For aerial application, use 5-30 gallons of spray volume per acre. Use a sufficient volume of water and an effective delivery system to ensure thorough coverage of the soil. The use of a hand-held or backpack sprayer is allowed, especially when treating smaller areas. The water volume and use rates are the same on a given area as if treating with a much larger boom sprayer. When using a hand-held or backpack sprayer, do not exceed the use rate restrictions stated on this label.

Esplanade F Herbicide may be used for preemergent residual control of certain weeds in establishment of conifer and hardwood species. Make a broadcast application of 3.5-7 fl oz /A of Esplanade F Herbicide prior to planting (site prep) or after transplanting (release) when the soil has settled. Applications made to trees under stress may injure these trees. Use 7 fl oz for longer residual control. The total of Esplanade F Herbicide applied in a 12 month period, however, must not exceed 10 fl oz/A. Esplanade F Herbicide may be tank-mixed with other pre and post emergent herbicides for broader spectrum of control and for control of existing weeds provided that all restrictions for both products are observed.

## MIXING INSTRUCTIONS

Ensure that the application equipment has been thoroughly cleaned from previous use before using to apply Esplanade F Herbicide.

Fill the spray tank with 1/2 of the required volume of water prior to the addition of Esplanade F Herbicide. Add the proper amount of Esplanade F Herbicide, and then add the rest of the water. Maintain sufficient agitation to ensure an adequate spray mixture during application.

If Esplanade F Herbicide is to be applied in a tank mixture with other pesticides, add the appropriate amounts of the tank mix partners in the following order: (a) products in water-soluble packaging (WSP), (b) Wettable powder (WP), (c) Wettable granule (WG) or other dry flowables, (d) fertilizers, (e) Esplanade F Herbicide, (f) other aqueous suspension products (SC), (g) soluble liquids (SL), (h) emulsifiable concentrates and other organic-solvent based formulations. Continue to fill the tank with water to the desired volume while agitating. **Maintain sufficient agitation during application to ensure a uniform spray mixture.**

**Re-suspending Esplanade F Herbicide in Spray Solution:** Like other suspension concentrates (SC's), this product will settle if left standing without agitation. Reagitate the spray solution for a minimum of 10 minutes before application.

## TIMINGS, USE RATES, AND MAXIMUM SEASONAL RATE FOR ESPLANADE F HERBICIDE

Apply Esplanade F Herbicide prior to germination of targeted weeds. Esplanade F Herbicide does not generally control weeds that have emerged. Esplanade F Herbicide needs to reach the soil surface for best efficacy. Removing debris prior to application will facilitate contact with soil. The desired rate of this product depends on the residual weed activity required and restrictions on the maximum amount of Esplanade F Herbicide per season. Esplanade F Herbicide may be applied at 3.5-7 fl oz/A either as a broadcast treatment during site preparation or to established stands, or as a banded spray to the rows of trees, prior to or after planting.

Do not exceed 7 fl oz of this product for a single application. Applications of Esplanade F Herbicide must not exceed 10 fl oz/A within the calendar year, or in a 12-month period from the time of the first application. As such, full dosages may be applied to all soil types and in

all regions where trees are grown. When tank mixing Esplanade F Herbicide with other products, follow the regionally-specific, crop-tree-specific, and soil-specific guidelines found on other product labels.

For late fall applications, apply Esplanade F Herbicide prior to when the ground freezes. Length of control may vary outside of the stated limits. Factors including soil type, rainfall, and the amount of vegetation at the time of treatment may affect weed control. Lower rates of Esplanade F Herbicide may be effective for sandy soils, whereas fine textured soils may require higher rates. If the herbicide is not activated by rainfall, control may not be satisfactory.

**Site Preparation** – Apply this product after plowing, bedding, burning, or any other mechanical or cultural practice. Excessive soil disturbance after application will substantially reduce the preemergent activity of the product.

Application during site preparation can be made by ground or air (helicopter only). Accurate calibration of equipment is necessary to achieve uniform droplet distribution and coverage on the soil surface. If excessive vegetation is present, insufficient amounts of Esplanade F Herbicide will reach the soil surface to ensure uniform preemergent weed control.

Conifers that can be planted include loblolly pine, longleaf pine, ponderosa pine, shortleaf pine, slash pine, sugar pine, Virginia pine, coast redwood, Douglas fir, and western larch.

Hardwoods that can be planted include oaks.

For tree species not listed above, site preparation treatments may be applied if the user has prior experience with Esplanade F Herbicide. Without prior experience, it is recommended that small areas be treated with Esplanade F Herbicide to determine selectivity on specific tree species before large scale treatments are made. The user accepts all responsibility for injury on any tree species not listed above.

**Herbaceous Weed Control** – Apply this product after transplanting as a broadcast spray, or by a banded application to beds or rows. Esplanade F Herbicide may be applied over-the-top of conifers and hardwoods. For hardwood release, apply Esplanade F Herbicide before crop trees break dormancy. Applications made over-the-top may injure trees at bud swell stage or beyond. Applications may be done by air or ground equipment.

Conifers that can be treated with Esplanade F Herbicide include loblolly pine, longleaf pine, ponderosa pine, shortleaf pine, slash pine, sugar pine, Virginia pine, coast redwood, and Douglas fir.

Applications made over-the-top may injure trees at bud swell stage or beyond. Hardwoods that can be treated include oaks.

In areas where other conifer species may be mixed in with the conifers listed above, Esplanade F Herbicide may be applied if the user has prior experience with, Esplanade F Herbicide on the other conifer species. With no prior experience, it is advised that either a small area of plantings be tested for conifer safety prior to treating larger areas, or make no application of Esplanade F Herbicide in these areas. The addition of a surfactant is not recommended with application made over the top of seedlings because tree damage may occur. Applications made over the top of white or red fir may result in severe injury or tree death.

Application made to trees under stress may cause tree injury.

## Spray Drift Management

Spray equipment and weather affect spray drift. Consider all factors when making application decisions. Where states have more stringent regulations, they must be observed. Avoiding spray drift is the responsibility of the applicator or grower. To reduce the potential for drift, the ground and aerial application equipment must be set to apply medium to large droplets (i.e., ASAE Standard 572) with corresponding spray pressure. Use high flow rate nozzles to apply the highest practical spray volume. With most nozzle types, narrower spray angles produce larger droplets. Follow the nozzle manufacturer's directions on pressure, orientation, spray volume, etc., in order to minimize drift and optimize coverage and control.

### Sensitive Areas

Sensitive areas to this product are defined as bodies of water (ponds, lakes, rivers, and streams), wetland, habitats of endangered species and non-labeled agricultural crop areas. Applicators must take all precautions necessary to keep spray drift from reaching sensitive areas. Only apply this product when the potential for drift to adjacent sensitive areas is minimal (e.g. when wind is blowing away from the sensitive areas). The applicator is responsible for considering all these factors when making decisions. Do not apply under circumstances where possible drift to unprotected persons, food, forage, or other plantings that might be damaged, as crops thereof may be rendered unfit for sale, use, or consumption.

### Wind

Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing. Many factors influence spray drift potential including droplet size, equipment type and local terrain. Drift potential increases if wind is in excess of 10 mph, gusty, or below 2 mph (due to inversion potential). Always make applications when there is some air movement to determine the direction and distance of possible spray drift. The applicator should be familiar with local conditions and how it may influence spray drift.

### Temperature Inversion

A surface temperature inversion (i.e., increasing temperature with increasing altitude) greatly increases the potential for drift. Avoid application when conditions are favorable to inversion. Presence of ground fog is a good indicator of a surface temperature inversion.

### Controlling Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that still provide sufficient coverage and control. Uniform spray coverage is important to maximize weed control. Applying larger droplets will reduce drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions such as wind speed, temperature and humidity, and temperature inversion situations.

Spray volume, pressure, and nozzle selection are all important for reducing drift. Select a high flow rate nozzle to apply the highest practical spray volume. High flow rate nozzles produce larger droplets. Use lower spray pressures within the recommended range for the nozzle. If a higher flow rate is needed, increase the nozzle size instead of increasing pressure. Lower spray pressures produce larger droplets. Also, consider using low-drift nozzles.

Drift control additive may also be used with most spray equipment to reduce the potential for drift. When using a drift control additive, read and follow all directions on the additive label.

Shielding the boom or individual nozzles may also reduce the potential for drift. However, it is the responsibility of the applicator to verify that the shield does not interfere with uniform spray coverage.

When applying by air (helicopter only), orient nozzles so that the spray is applied backwards, parallel to the airstream. This practice will produce larger droplets than other nozzle orientations. Use a boom length and position that prevents the spray from entering the rotor vortices. Set the boom and make applications at the lowest height that safely permits uniform coverage of the soil and minimizes droplet evaporation.

### Tank Mix Combinations

Esplanade F Herbicide may be tank mixed with any herbicide product registered for use in conifer and hardwood areas. The list of herbicides that can be tank mixed with Esplanade F Herbicide includes but is not limited to: 2,4-D, aminopyralid, fosamine, Finale® VU glyphosate, hexazinone, imazapyr, metsulfuron, sulfometuron, and triclopyr. In areas where other conifer species may be mixed in with the conifers listed above, Esplanade F Herbicide may be applied if the user has prior experience with, Esplanade F Herbicide on the other conifer species. With no prior experience, it is advised that either a small area of plantings be tested for conifer safety prior to treating larger areas, or make no application of, Esplanade F Herbicide in these areas.

Follow all use restrictions on this label and for all tank mix partners and use the most restrictive use pattern for the labels of all products in a tank mixture. It is the pesticide user's responsibility to ensure that all products in the mixtures are registered for the intended use.

Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another). Apply mixtures so that the spray solution covers the soil surface in a uniform manner. If uniform coverage is not achieved, preemergent activity will be inconsistent.

## COMPATIBILITY TESTING WITH OTHER PESTICIDES

A compatibility test must be conducted with any potential tank mix partner with Esplanade F Herbicide. Using a clear container, conduct the test as described below:

Fill the container three-quarters full with water.

1. Add the appropriate amounts of tank mix partners in the following order: (a) products in water-soluble packaging (WSP), (b) Wettable powder (WP) (c) Wettable granule (WG) or other dry flowables, (d) fertilizers, (e) Esplanade F Herbicide (f) other aqueous suspension products (SC), (g) soluble liquids (SL), (h) emulsifiable concentrates and other organic-solvent based formulations. Shake or gently stir after each addition to mix thoroughly.
2. After adding all ingredients, let the mixture stand for 15 minutes and look for separation, large flakes, precipitates, gels, and heavy oily film or other signs of incompatibility.
3. If the compatibility test shows signs of incompatibility, do not tank mix the product tested with Esplanade F Herbicide.

## Resistance Management Guidelines

Continual use of herbicides with a single mode of action encourages the development of resistant weeds. Esplanade F Herbicide is a Group 29 Herbicide that contains the active ingredient indaziflam. Esplanade F Herbicide may be used in programs with other preemergence herbicides with different modes of action. No known resistance to Esplanade F Herbicide exists, and there are no known instances of cross resistance between this product and other classes of herbicides, or modes of action. Performance of this product is not affected by the presence of biotypes resistant to glyphosate, triazines, ALS-inhibiting, growth regulator, or other herbicide modes of action. When resistance of a specific weed is confirmed, rotation of Esplanade F Herbicide in one season followed by a preemergent herbicide with another mode of action in the subsequent season, for example, will reduce existing populations and minimize further development of resistant weeds. Contact a Bayer CropScience LP representative for the latest information on resistance management guidelines for this product.

## Weeds Controlled or Suppressed by Esplanade F Herbicide

### Broadleaf Weeds Controlled

American black nightshade	<i>Solanum americanum</i>	Kochia	<i>Kochia scoparia</i>
Bittercress	<i>Cardamine sp.</i>	Lambsquarters, common	<i>Chenopodium album</i>
Buckweed, wild (seedlings only)	<i>Polygonum convolvulus</i>	Lawn burweed	<i>Soliva pterosperma</i>
California burclover	<i>Medicago polymorpha</i>	Little mallow	<i>Malva parviflora</i>
Canada thistle, common (seedlings)	<i>Cirsium arvense</i>	Long-stalk phyllanthus	<i>Phyllanthus tenellus</i>
Carpentweed	<i>Mollugo verticillata</i>	Panicle willowweed	<i>Epilobium paniculatum</i>
Chickweed, common	<i>Stellaria media</i>	Plantain, Buckhorn	<i>Plantago lanceolata</i>
Chickweed, Mouse-ear	<i>Cerastium vulgatum</i>	Plantain, Paleseed	<i>Plantago virginica</i>
Clover, White	<i>Trifolium repens</i>	Prostrate knotweed	<i>Polygonum aviculare</i>
Corn speedwell	<i>Veronica arvensis</i>	Prostrate pigweed	<i>Amaranthus blitoides</i>
Cudweed, Linear-leaf/purple	<i>Gnaphalium purpureum</i>	Prostrate spurge	<i>Euphorbia humifusa</i>
Curly dock (seedlings)	<i>Rumex crispus</i>	Purslane, common	<i>Portulaca oleracea</i>
Cutleaf evening primrose	<i>Oenothera laciniata</i>	Ragweed, common	<i>Ambrosia artimisiifolia</i>
Dandelion, cat's ear	<i>Hypochoeris radicata</i>	Red tasselflower	<i>Emilia sonchifolia</i>
Dandelion, common (seedlings)	<i>Taraxacum officinale</i>	Redmaids	<i>Calandrinia caulescens</i>
Doveweed	<i>Murdannia nudiflora</i>	Redroot pigweed	<i>Amaranthus retroflexus</i>
Eclipta	<i>Eclipta alba</i>	Redstem fleabane/Storks-bill	<i>Erodium cicutarium</i>
Evening primrose, common	<i>Oenothera biennis</i>	Russian Thistle	<i>Salsola tragus</i>
Evening primrose, cutleaf	<i>Oenothera laciniata</i>	Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Filaree, redstem	<i>Erodium cicutarium</i>	Sowthistle, Annual	<i>Sonchus olerachus</i>
Fleabane, blackleaved	<i>Conza bonariensis</i>	Spotted catsear	<i>Hypochoeris radica</i>
Florida pusley	<i>Richardia scabra</i>	Swinecress	<i>Coronopus didymus</i>
Gromwell, Yellow	<i>Amsinckia calycina</i>	Tropic ageratum	<i>Ageratum conyzoides</i>
Groundsel, common	<i>Senecio vulgaris</i>	Velvetleaf	<i>Abutilon theophrasti</i>
Hairy fleabane	<i>Erigeron bonariensis</i>	Wild buckwheat (seedlings)	<i>Polygonum convolvulus</i>
Hairy nightshade	<i>Solanum sarrachoides</i>	Wild mustard	<i>Sinapis arvensis</i>
Henbit	<i>Lamium amplexicaule</i>	Yellow starthistle	<i>Centaurea solstitialis</i>
Horseweed/Marestail	<i>Erigeron canadensis</i>		

### Grasses and Sedges Controlled

Annual bluegrass	<i>Poa annua</i>	Crabgrass, Henry	<i>Digitaria adscendens</i>
Annual bromegrass	<i>Bromus spp.</i>	Crabgrass, Large/Hairy	<i>Digitaria sanguinalis</i>
Barnyardgrass, common	<i>Echinochloa crus-galli</i>	Crabgrass, Smooth	<i>Digitaria ischaemum</i>
Crabgrass	<i>Digitaria species</i>	Cheatgrass	<i>Bromus secalinus</i>

### Grasses and Sedges Controlled (continued)

Downy brome	<i>Bromus tectorum</i>	Medusahead	<i>Taeniatherum caput-medusae</i>
False chamomile	<i>Matricaria maritima</i>	Mouse barley	<i>Hordeum murinum</i>
Foxtail brome	<i>Bromus rubens</i>	Sedge, annual	<i>Cyperus</i> spp.
Foxtail, Giant	<i>Setaria faberi</i>	Rice flatsedge	<i>Cyperus iria</i>
Foxtail, Green	<i>Setaria viridis</i>	Ryegrass, Italian	<i>Lolium multiflorum</i>
Foxtail, Yellow	<i>Pennisetum glaucum</i>	Ryegrass, Perennial	<i>Lolium perenne</i>
Goosegrass	<i>Eleusine indica</i>	Sprangletop	<i>Leptochloa</i> spp.
Guineagrass	<i>Panicum maximum</i>	Tufted lovegrass	<i>Eragrostis pectinacea</i>
<b>Weeds Suppressed</b>			
Black medic	<i>Medicago lupulina</i>	Southern brassbuttons	<i>Cotula australis</i>
Black mustard	<i>Brassica nigra</i>	Sunflower, common	<i>Helianthus</i> spp.
False chamomile	<i>Matricaria maritima</i>	Vetch, purple	<i>Vicia benghalensis</i>
London rocket	<i>Sisymbrium irio</i>	Wild carrot	<i>Daucus carota</i>
Prickly lettuce	<i>Lactuca serriola</i>	Woodsorrell, yellow	<i>Oxalis stricta</i>
Sesbania, hemp	<i>Sesbania exaltata</i>	Woodsorrel/Oxalis	<i>Oxalis</i> species
Sida, prickly/teaweed	<i>Sida spinosa</i>		

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store product in original container only. Store in cool, dry place.

**Pesticide Disposal:** Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" designation.

**Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons)**

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

**Rigid Non-refillable containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 pounds)**

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

Do not transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire, or other emergency, contact BAYER CROPSCIENCE LP at 1-800-334-7577, day or night.

## CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, plant injury and other property damage, as well as other unintended consequences may result because of factors beyond the control of Bayer CropScience LP. Those factors include, but are not limited to, weather conditions, presence of other materials or the manner of use or application. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd) and Esplanade® are registered trademarks of Bayer.

**Produced for:**

**Bayer Environmental Science  
A Division of Bayer CropScience LP  
5000 CentreGreen Way, Suite 400  
Cary, NC 27513**

**Bayer**



# EsplAnade<sup>®</sup>

## F

## HERBICIDE

For Preemergent Weed Control in Conifer and Hardwood Production Areas

<b>ACTIVE INGREDIENT:</b>	
Indaziflam (CAS No: 950782-86-2)	19.05%
<b>OTHER INGREDIENTS:</b>	
	80.95%
<b>TOTAL:</b>	100.00%

EPA Reg. No. 432-1517

Contains 1.67 pounds of indaziflam per gallon

## KEEP OUT OF REACH OF CHILDREN CAUTION

See Panel for First Aid Instructions and Booklet for Complete Precautionary Statements and Directions for Use.

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY**  
Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-800-331-2867

### FIRST AID

<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eyes open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> <li>Move person to fresh air.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

For **MEDICAL** Emergencies Call 24 Hours A Day  
1-800-334-7577

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**Note to physician:** No specific antidote is available. Treat symptomatically.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or

clothing. Avoid breathing spray mist. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean water mark. Do not contaminate water when disposing of rinseate or washwater. This product may impact water through spray drift or runoff. Follow directions for use to avoid spray drift and runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential of this product entering water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

**Groundwater Advisory:** This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory:** This pesticide may impact water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store product in original container only. Store in cool, dry place.

**Pesticide Disposal:** Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" designation.

**Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons)**

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinseate into application equipment or a mix tank or store rinseate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

Do not transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire, or other emergency, contact BAYER CROPSCIENCE LP at 1-800-334-7577, day or night.

Nonrefillable Container  
Net Contents

**2.5 Gallons**  
84488305

84401528B 160419AV5

Bayer

Produced for:  
Bayer Environmental Science  
A Division of Bayer CropScience LP  
5000 CentreGreen Way, Suite 400  
Cary, NC 27513

(01)007857404997 19