

Supplemental Labeling

E.I. du Pont de Nemours and Company 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Zest™ WDG

EPA Reg. No. 352-560

Reduced CRI in Sorghum

This supplemental labeling expires January 31, 2024 and must not be used or distributed after this date.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Zest™ WDG before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Zest WDG according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Zest WDG.

DIRECTIONS FOR USE

CROP ROTATION

Observe the following rotational intervals when using Zest WDG at a maximum of 1.33 ounces (0.062 lb ai):

ZEST WDG ROTATIONAL CROP GUIDELINE - 1

No soil pH restrictions

Rotational Crop	Interval in Months
Corn (field, pop, seed, sweet*)	Anytime
Soybeans	0.5 (15 days)
Cereals, spring (barley, oats, rye, wheat)	8
Cereals, winter (barley, oats, rye, wheat)	4
Cotton	10
Dry beans, Peas, Snap beans	10
Alfalfa**	12
Red Clover**	12
Sorghum	10
Sorghum (hybrids containing ALS-resistant traits)	18
Other Crops	See Rotational Crop guidelines 2 and 3

* Except the sweet corn varieties "Merit", "Carnival", and "Sweet Success", for which the minimum time interval is 15 months.

**Except for the state of Kansas east of Highway 75, for Minnesota east and south of the Red River Valley and for the states east of the line formed by the western borders of Iowa, Missouri, Arkansas, and Louisiana, where the minimum time interval is 10 months.

ZEST WDG ROTATIONAL CROP GUIDELINE - 2

With soil pH <7.5 restrictions

Crop	Rotational Interval in Months	
	pH 7.5	pH > 7.5
Sunflowers*	11	18
All other crops not listed in Rotational guidelines 1 or 2	See Rotational guideline 3	

*Precipitation following application must exceed 14" prior to planting sunflowers.

ZEST WDG ROTATIONAL CROP GUIDELINE - 3

With soil pH <6.5 restrictions

Crop	Rotational Interval in Months	
	pH 6.5	pH > 6.5
Sugarbeets*, potatoes**	10	18***
All other crops not listed in Rotational guidelines 1 or 2	10	18

* Except on irrigated sites in Colorado, Wyoming, Nebraska, Texas, Michigan, and Ohio, where precipitation following application must exceed 25" prior to planting beets, where the interval is 10 months on soils with pH < 7.5. Sites in Minnesota east and south of the Red River Valley may follow these guidelines provided maximum rates of Zest WDG **DO NOT** exceed 0.67 ounces (0.031 lb ai).
 **Irrigated potatoes following irrigated corn treated in the States of Washington, Oregon, Idaho, or Utah can be planted 10 months after using Zest WDG on sprinkler irrigated corn with no soil pH restrictions, providing the maximum use rate on corn does not exceed 1.0 ounce product (0.046 lb ai) per year. Corn treated with Zest WDG must be grown to maturity and receive a minimum of 18 inches of irrigation water before potatoes can be planted at this rotation interval. Injury to potatoes may occur if less than 18 inches of irrigation is used on the previous corn crop. Zest WDG may not be used in a tankmix or sequential application program with other ALS-inhibiting herbicides including "Beacon" (AI: primisulfuron-methyl; Reg. No. 100-705).
 ***In North Dakota and northwest Minnesota, the cumulative precipitation in the 18 months following application must exceed 28" in order to rotate to sugarbeets or potatoes.

ROTATIONAL CROP GUIDELINES - 4 may be observed when using a single application of Zest WDG per year with a maximum use rate of 0.67 ounces product (0.031 lb ai). Extend rotational intervals to 12 months if drought conditions prevail after application and before the rotational crop is planted, unless sprinkler irrigation has been applied and totals greater than 15" during the growing season.

ZEST WDG ROTATIONAL CROP GUIDELINES - 4

With 0.67 ounces (0.031 lb ai) maximum use rate

Crop	Rotational Interval in Months
Alfalfa*	10
Canola	10
Flax**	10
Sorghum	10
Sorghum (hybrids containing ALS-resistant traits)	18
Potato	10
Red clover	10
Sunflower	10

*On sprinkler irrigated fields in Idaho, Utah, and Northern Nevada it is best to use deep fall tillage including plowing prior to planting alfalfa. Product degradation may be less on furrow irrigated soils and may result in some crop injury.

**Extend rotational intervals to 18 months if drought conditions prevail after application and before the rotational crop is planted, unless sprinkler irrigation has been applied and totals greater than 15" during the growing season.

® TM Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners.

R635-001
 EPA accepted 01/28/2021
 Replaces: Initial