

FIFRA Section 24(c) Special Local Need

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF GEORGIA

RESTRICTED USE PESTICIDE

For sale to and use only by Certified Applicators.

READ THE ENTIRE PRODUCT LABELING FOR XTENDIMAX® WITH VAPORGRIP® TECHNOLOGY BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS FIFRA SECTION 24(c) SPECIAL LOCAL NEED LABEL.

When using XtendiMax® With VaporGrip® Technology as permitted according to this 24(c) Special Local Need label, read and follow all applicable directions, restrictions, and precautions on the container label and booklet provided with the product container, on applicable supplemental labeling for the product, and on this 24(c) Special Local Need labeling. This labeling, and all other product labeling authorizing the desired application, must be in the possession of the user at the time of pesticide application.

This FIFRA 24(c) Special Local Need label expires on 12/20/2020 and must not be used or distributed after this date.



EPA Reg. No. 524-617

EPA SLN No. GA-190001

DICAMBA	GROUP	4	HERBICIDE
---------	-------	---	-----------

ADDITIONAL APPLICATION RESTRICTIONS FOR USE ON DICAMBA-TOLERANT COTTON OR DICAMBA-TOLERANT SOYBEANS IN GEORGIA.

Keep out of reach of children

CAUTION!

In case of an emergency involving this product, call collect, day or night, 314-694-4000.

Bollgard II®, Bollgard® 3, Roundup Ready®, Roundup Ready 2 Xtend®, XtendiMax®, XtendFlex® and VaporGrip® are registered trademarks of Monsanto Technology LLC. All other trademarks are the property of their respective owners.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

See the "PRODUCT INFORMATION" and "APPLICATION EQUIPMENT AND TECHNIQUES" sections of the XtendiMax® With VaporGrip® Technology product label for important use information. In the event that there are any inconsistencies with the directions for use between this 24(c) label and any other labeling for this product, follow the directions for use on this 24(c) label.

TYPES OF APPLICATIONS: Preplant; At-Planting; Preemergence; Postemergence (In-crop)

XtendiMax® With VaporGrip® Technology is approved by U.S. EPA to be used in the following states, subject to county restriction as noted: Alabama, Arkansas, Arizona, Colorado, Delaware, Florida (excluding Palm Beach County), Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Jersey, New Mexico, New York, North Carolina, North Dakota, Oklahoma, Ohio, Pennsylvania, South Carolina, South Dakota, Tennessee (excluding Wilson County), Texas, Virginia, West Virginia, Wisconsin.

RESTRICTIONS/LIMITATIONS

USING PESTICIDES WISELY TRAINING:

The certified applicator applying this product on cotton or soybeans, in-crop, must attend Using Pesticides Wisely training, on an annual basis, that is approved by the Georgia Department of Agriculture. For more information on this training, contact the Georgia Department of Agriculture at 404-656-4958 or your local County Cooperative Extension Service office.

Tank Mix Restrictions:

XtendiMax® With VaporGrip® Technology may be mixed with any other labeled cotton or soybean herbicides listed at www.xtendimaxapplicationrequirements.com. Glufosinate containing products (Liberty, etc) contain ammonium and are not approved as a tank-mix partner. Also, do not include any type of ammonium salts such as ammonium sulfate (AMS) and urea ammonium nitrate.

ADDITIONAL APPLICATION METHODS

HOODED ROW-MIDDLESPRAY APPLICATIONS:

Applications may be made with any standard spray tip for hoods as long as the system generates droplets that are coarse or larger in size (>341microns vmd). Contact UGA Extension or your local Bayer representative for questions.

Application Requirements:

1. Hoods must remain in contact with the soil while the herbicide application is being made.
2. A maximum of 6 mph is required to ensure hoods remain in contact with the soil.
3. Cotton or soybeans must be at least 15 inches tall at time of application.

Application Timing:

Cotton with XtendFlex® Technology: In-crop application of this product may be made from emergence up to 7 days pre-harvest.

Roundup Ready 2 Xtend® Soybean: In-crop applications of this product may be made from emergence (cracking) up to and including beginning bloom (R1 growth stage of soybeans). Do not make in-crop applications of this product after beginning bloom (R1 growth stage of soybeans).

DIRECTED LAYBY APPLICATIONS:

Applications may be made with any standard spray tip as long as the system generates droplets that are coarse or larger in size (>341 microns vmd). Contact UGA Extension or your local Bayer representative for questions.

Application Requirements:

1. Release point for the herbicide must not be more than 10 inches from the soil and the cotton or soybean must be at least 20 inches in height. Spray tip must be angled downward to the soil making sure no spray droplets remain in the air.
2. A maximum of 6 mph is required to ensure herbicide does not lift above the release point.

Application Timing:

Cotton with XtendFlex® Technology: In-crop application of this product may be made from emergence (cracking) up to 7 days pre-harvest.

Roundup Ready 2 Xtend® Soybean: In-crop applications of this product may be made from emergence (cracking) up to and including beginning bloom (R1 growth stage of soybeans). Do not make in-crop applications of this product after beginning bloom (R1 growth stage of soybeans).

©2019
MONSANTO COMPANY
800 N. LINDBERGH BLVD.
ST. LOUIS, MISSOURI 63167 USA

24(c) Registrant:
Georgia Department of Agriculture
Plant Industry Division
19 Martin Luther King, Jr. Drive, Room 429
Atlanta, GA 30334

1/2/2019

35008T3-31

12/20/2020