

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF TEXAS

THIS FIFRA SECTION 2(EE) EXPIRES DECEMBER 31, 2020

EPA Reg. No. 279-3594

To permit the use of Topguard® Terra fungicide in cotton as preplant injection, preemergence, or early postemergence application for control of cotton root rot (*Phymatotrichopsis omnivora*).

Directions for Use

- **IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**
- **ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.**
- **This recommendation must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Topguard® Terra fungicide before applying.**

Crop

- Cotton

Use Rate and Method of Application

- Topguard® Terra fungicide may be injected into row up to 30 days prior to planting. Fungicide should be placed 2-3 inches directly below seed placement zone.
- Topguard® Terra fungicide may be applied in 3 – 4 inch band on top of row prior to cotton emergence.
- Topguard® Terra fungicide may be applied to emerged cotton either as a 4 inch band over-the-top of cotton not larger than 2 true leaves or as a directed spray at the base of cotton not larger than 4 true leaves. Directed spray should be concentrated in band no wider than 4 inches at base of plants.

Rates and Timing

Application Timing	Use Rate (fl. oz/A) (lb a.i./A)	Comments
Postemergence OTT of cotyledon to 2 leaf cotton. Directed-postemergence 2-4 leaf cotton. Injected into row up to 30 days prior to planting. Preemergence in 3-4 inch band behind the planter press wheel.	4-8 fl oz/A (0.13 – 0.26 lb ai/A)	Use higher rates on irrigated fields or where high infestations of cotton root rot are common. Rainfall or irrigation is required to move Topguard Terra into infection zone.