

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA



EPA Reg. No. 279-3557

2(ee) Recommendation
FOR DISTRIBUTION AND USE ONLY IN WASHINGTON, OREGON, & IDAHO
THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)
IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

DIRECTIONS FOR USE: HOPS, GRAPES AND STONE FRUIT

When applied as directed through drip irrigation and micro sprinklers, Topguard Fungicide Specialty Crops will provide control of powdery mildew on the foliage of hops, grapes and stone fruit.

Crop	Pest	Rate (fl. oz/A)	Application Directions	Retreatment Interval (Days)
Hops	Powdery Mildew (<i>Podosphaera macularis</i> , <i>Sphaerotheca macularis</i>)	14	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	14
Grapes (fresh, table, wine, raisin)	Powdery Mildew (<i>Uncinula necator</i>)	10	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	14
Stone Fruit (See label for list of stone fruit)	Powdery Mildew (<i>Podosphaera clandestina</i>)	14	Start applications at petal fall. Repeat as necessary if conditions are favorable for disease development.	7

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE TOPGUARD FUNGICIDE SPECIALTY CROPS LABEL (EPA REG. NO. 279-3557) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.