FIFRA Section 24(c) Special Local Need



FOR DISTRIBUTION AND USE ONLY WITHIN FLORIDA



FOR CONTROL OF ASIAN CITRUS PSYLLID (ACP) ON CITRUS DURING BLOOM PERIODS

EPA Reg. No. 59639-243 EPA SLN No. SLNFL210001

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2026

Active Ingredients	By Wt
Spirotetramat*	11.47%
Pyriproxyfen**	4.01%
Other Ingredients	84.52%
Total	100.00%

^{*}cis-3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4.5]dec-3-en-4-yl ethyl carbonate

Senstar Insecticide is a suspo-emulsion (SE) formulation that contains 1 pound spirotetramat and 0.35 pound pyriproxyfen per gallon.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Persons using this product must comply with all applicable directions, restrictions, and precautions found on this labeling and that of the label of the federally registered product upon which this amendment is based. Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label. This labeling must be in the possession of the user at the time of application.

Citrus Fruit (Crop Group 10-10)

Australian Desert Lime; Australian Finger-lime; Australian Round Lime; Brown River Finger Lime; Calamondin; Citron; Citrus hybrids; Grapefruit; Japanese Summer Grapefruit; Kumquat; Lemon; Lime; Mediterranean Mandarin; Mount White Lime; New Guinea Wild Lime; Orange, Sour; Orange, Sweet; Pummelo; Russell River Lime; Satsuma Mandarin; Sweet Lime; Tachibana Orange; Tahiti Lime; Tangelo; Tangerine (Mandarin); Tangor; Trifoliate Orange; Uniq Fruit; Cultivars, varieties, and/or hybrids of these

	PRODUCT RATES PER ACRE		
Pest Controlled	Fl. oz/ Acre	Lb.ai/ Acre	
	Formulated Product	Spirotetramat	Pyriproxyfen
Asian Citrus Psyllid	20	0.156	0.055

Restrictions

- Do not apply more than 20 fl oz of Senstar Insecticide (0.156 lb spirotetramat and 0.055 lb pyriproxyfen) per acre per application.
- Do not make more than one application during the primary citrus bloom period.
- Do not apply more than 40 fl oz of Senstar Insecticide (0.313 lb spirotetramat and 0.109 lb pyriproxyfen) per acre per year.
- Minimum interval between applications is 21 days.
- Do not apply within 1 day of harvest.
- Do not apply nonionic surfactants in tank mix combination with Senstar Insecticide on white grapefruit.

Please contact Valent U.S.A. LLC at 800-6-VALENT (682-5368) if you have questions regarding use of this product.

FIFRA 24(c) Registrant: Valent U.S.A. LLC

P.O. Box 5075

San Ramon, CA 94583

^{**4-}phenoxyphenyl (RS)-2-(2-pyridyloxy)propyl ether