UPBEET®

HERBICIDE

FIFRA 24(c) Special Local Need Label (SLN)

For Distribution and Use only in the State of New York
For Postemergence Weed Control in Garden Beets

EPA REG No. 279-9584 SLN No. NY-190008

(REGISTRATION TRANSFERRED FROM EPA REG. NO. 352-569 ON MAY 25, 2018)

This label expires and must not be distributed or used in accordance with this SLN registration after April 30, 2024

DIRECTIONS FOR USE:

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions, and precautions including statements pertaining to Worker Protection Standards on the EPA registered UPBEET® herbicide (UPBEET herbicide) label.
- Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

PRODUCT INFORMATION

UPBEET herbicide may be used in Garden Beets for selective postemergence control of broadleaf weeds including wild mustard, shepherd's purse, and velvetleaf. This product is a water dispersible granule containing 50% active ingredient by weight.

APPLICATION INFORMATION

Apply UPBEET herbicide by ground only at broadcast rate of 0.5 ounces per acre, starting when garden beets are at the 2 to 4 leaf stage. Additional applications may be made at the 4 to 6 leaf stage and at the 6 to 8 leaf stage. The total amount of UPBEET herbicide applied must not exceed 1.5 ounces per acre per growing season. For best results apply UPBEET herbicide to small actively growing weeds when the temperature is between 40F and 75F. If high temperatures are expected during the day, treatment should be applied early in the morning or in the evening, so that a period of moderate temperatures (lower than 75F) will follow the application. In some cases, discoloration and even slowed growth may be observed. These symptoms are usually temporary.

WEED STAGE AT APPLICATION

Weeds should be actively growing and not under stress. Applications made to larger weeds or to weeds under stress may result in unsatisfactory control. Since UPBEET herbicide has little to no soil activity, only weeds that have emerged above the soil surface will be controlled. Use sequential applications to control new weed flushes. The growth stage of weeds at application is very important for satisfactory control. For best results, apply to small, emerged weeds between the cotyledon and two true leaf stage

R-4295 102219 10-22-19

TANK MIX OPTIONS

UPBEET herbicide may be tank mixed with other suitable registered herbicides to broaden the weed control. FMC recommends that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or FMC representative as to the potential for crop injury before using the mixture. If no information is available, limit the initial use of UPBEET herbicide and the tank mix herbicide to a small area.

UPBEET herbicide may also be mixed with other suitable registered fungicides or insecticides labelled for use on garden beets. Read all label precautions for tank mix partners prior to use. Follow all manufacturers' label directions for the companion product. If these directions conflict with this label, do not tank mix with UPBEET herbicide.

PRECAUTIONS/RESTRICTIONS

- **Do Not** use this product until you have read the entire label.
- **Do Not** apply this product in a way that will contact workers or other persons either directly or through drift.
- **Do Not** use in Suffolk and Nassau Counties, Long Island, New York.
- **Do Not** treat when frost is expected in the hours following application.
- **Do Not** apply this product by aerial application methods in New York state
- **Do Not** Apply more than 1.5 ounces of product per acre per year in New York state.
- Apply no later than 30 days before harvest.
- Refer to the Section 3 Federal Product label for additional application instructions and use precautions.

SPRAY ADDITIVES

Nonionic Surfactant (NIS) and Crop Oil Concentrate (COC)

- A tank mix with 0.25% volume/volume (2 pt per 100 gal of spray solution) of non-ionic surfactant or crop oil concentrate may be added. Addition of surfactant may increase crop response. First time users should limit the use of surfactant to a small test area to determine if crop response is acceptable
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

This SLN contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully. This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal Product label.

FMC, and UPBEET, are trademarks of FMC Corporation or an affiliate

Sold By:

FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

R-4295 102219 10-22-19