



## FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF  
ALABAMA, FLORIDA, AND GEORGIA

### Roundup PowerMAX® 3 Herbicide

EPA Reg. No. 524-659

#### Recommendations Prior to Transplanting

#### Brassica Vegetables, Cucurbits and Fruiting Vegetables in Sandy Soils

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Read the product labeling affixed to the container of Roundup PowerMAX® 3 Herbicide before applying. Use of Roundup PowerMAX® 3 Herbicide according to this product bulletin is subject to the use precautions and limitations imposed by the labeling affixed to the container.

#### CROPS

**Brassica Vegetables:** broccoli, Chinese broccoli (gai lon), broccoli raab (rapini), brussels sprouts, cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), cauliflower, cavalo broccolo, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens.

**Cucurbit Vegetables and Fruits:** cantaloupe, casaba melon, crenshaw melon, cucumber, gherkin, gourds, honeydew melon, honey ball melon, mango melon, melons (all), muskmelon, Persian melon, pumpkin, squash (summer, winter), and watermelon.

**Fruiting Vegetables:** eggplant, groundcherry, okra, pepino, pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato.

**Root and Tuber Vegetables:** turnip

When applying Roundup PowerMAX® 3 Herbicide prior to transplanting these crops in bare ground production soil with over 85% sand and/or less than 0.5% organic matter, if the soil is not tilled after application and before planting, apply no more than 30 fluid ounces of this product per acre in a single application, and allow for a minimum accumulation of 0.5 inch of rainfall or overhead irrigation and wait 7 or more days between application and transplanting.

When applying Roundup PowerMAX® 3 Herbicide prior to transplanting these crops in mulch production where soil is over 85% sand and/or less than 0.5% organic matter, wait 3 more days before transplanting following a single application of this product up to 60 fluid ounces per acre, or wait 10 or more days following single application between 30 and 60 fluid ounces per acre, and allow for a single rainfall or irrigation event of at least 0.5 inch between application of this product at any rate and transplanting. Punch new transplant holes and place plants a minimum of 3 inches from old holes or torn mulch.

Single Application Rate of Roundup PowerMAX® 3 Herbicide	Interval Between Application and Transplanting	Rainfall/Irrigation Before Transplanting
Up to 30 fluid ounces	3 or more days before transplanting	0.5 inch or more
30 to 60 fluid ounces	10 or more days before transplanting	0.5 inch or more

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

**For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).**

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

**Bayer CropScience LP**  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167