

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

FIFRA 24(c) Special Local Need Label (SLN)

For distribution and use only in the state of CALIFORNIA

For use on citrus tree trunks or bare soil beneath citrus tree for control of Fuller's Rose Beetle



EPA REG. No. 279-3108

SLN#: CA-130011

This label expires and must not be distributed or used in accordance with this SLN registration after September 30, 2028.

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA registered label for Brigade WSB insecticide/miticide, EPA Reg. No. 279-3108, and this label.

Crop	Pest	Rate of Application
Citrus tree trunks and/or bare soil beneath citrus Trees including: oranges, lemons, mandarins, and grapefruit	Fuller's Rose Beetle (<i>Pantomorus cervinus</i>)	For a 2 Spray Program use: 5 lbs product/A (0.5 lb ai/A) For a 4 Spray Program use: 2.5 lbs product/A (0.25 lb ai/A)

Directions for use: as a Soil Treatment use 5 lbs. Product/A (0.5 lb ai/A):

Treatments should be initiated August - September

Apply Brigade WSB Insecticide/Miticide by ground equipment to bare soil beneath skirt pruned citrus trees. Brigade WSB Insecticide/Miticide must be uniformly applied from the trunk to the drip line of the tree. Brigade WSB Insecticide/Miticide forms an insecticide barrier on the soil so disturbance of the soil beneath trees should be minimized to keep the barrier intact.

Apply in a minimum of 40 gallons of spray per acre however, greater spray volume may achieve more uniform results. Use the full per acre rate concentrated in a band in the drip line of the citrus trees. Do not allow Brigade WSB Insecticide/Miticide to be directly sprayed on or allow spray drift to contact the fruit or foliage of citrus trees.

Directions for Use: trunk spray

Treatments should be initiated starting June 1st. Apply Brigade WSB Insecticide/Miticide only by ground using a hand wand sprayer or shielded sprayer to the lower 18 inches of tree trunks. Do not allow spray to contact the foliage or fruit



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

Brigade and FMC -FMC Trademarks
©2012 FMC Corporation
R-4800 060223 08-08-23

of the tree. For a 2-spray program, make applications 12 to 16 weeks apart. For a 4-spray program, make applications 6 to 8 weeks apart.

Apply to the lower trunk as a directed spray. Uniform coverage of the trunk is critical for pest control. Spray the trunks of 10 trees with water to determine the amount of spray that is needed for thorough coverage of the trunk, without runoff, from the soil line to approximately 18 inches up the trunk. Measure the amount of water needed to refill the spray tank to its initial level. Then calculate the amount of spray needed per acre, e.g., if it took 0.5 gallons to spray 10 trees and there are 180 trees per acre, then you will need $0.5 \times (180/10) = 9$ gal./A of spray mix. Based on the number of acres to be sprayed, put that amount of water per acre in the tank along with the appropriate amount of Brigade WSB Insecticide/Miticide.

The Preharvest Interval (PHI) is 63 days for the 2 spray program and 28 days for the 4 spray program.

Restrictions / Precautions:

- For use under this SLN label, do not apply this product through any type of irrigation system.
- Do not apply more than 10 pounds of product (1.0 lb. active ingredient) per acre, per season.
- Apply only to skirt-pruned citrus trees. Tree skirts must be pruned to a minimum height of 2 feet above the ground



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

Brigade and FMC -FMC Trademarks
©2012 FMC Corporation
R-4800 060223 08-08-23