

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND
HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA Reg. No. 279-3108

BIFENTHRIN	GROUP	3A	INSECTICIDE
------------	-------	----	-------------

2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

This Product Bulletin Contains New or Additional Directions For Use Which Are Recommended By FMC Corporation Under FIFRA Section 2(ee)

This recommendation for BRIGADE WSB Insecticide/Miticide is valid until 03/16/2026,
or until withdrawn, cancelled or suspended.

For Agricultural or Commercial Use Only

CROP	PEST	RATE OF APPLICATION
Citrus	Fuller Rose Beetle <i>Asynonychus godmani</i>	40 – 80 oz product/A (0.25 - 0.5 lb ai/A)

DIRECTIONS FOR USE: A single or split application as a soil or trunk treatment may be made depending on pest emergence.

Single Application: A single application of 80 oz product/A (0.5 lb ai/A) may be made before pest emergence.

Split Application: A split application using one application of 40 oz product/A (0.25 lb ai/A) followed later by a second application of 40 oz product/A (0.25 lb ai/A) may be made. Applications should be made prior to peak pest emergence.

Soil Treatment Recommendations: Make a uniform application to the soil beneath trees from the trunk to the drip line in a minimum of 40 gallons finished spray solution per acre. See product label for full soil application details. Initiate application in June-July if you are targeting larval and pupal stage. For adults, apply from July- September.

Trunk Treatment Recommendations: Application should be initiated starting July 1st to the lower trunk as a directed spray. Uniform coverage of the trunk is critical for pest control. Spray the trunks of 10 trees with water to determine the amount of spray that is needed for thorough coverage of the trunk, without runoff, from the soil line to approximately 18 inches up the trunk. Measure the amount of water needed to refill the spray tank to its initial level. Then calculate the amount of spray needed per acre, e.g. if it took 0.5 gallons to spray 10 trees and there are 180 trees per acre then you will need 0.5 X (180/10) = 9 gal/A of spray mix. Based on the number of acres to be sprayed, put that amount of water per acre in the tank along with the appropriate amount of BRIGADE WSB Insecticide/Miticide (80 oz/A for a single application or 40 oz/A for a split application)

RESTRICTIONS:

- Do not apply through irrigation systems.
- Do not allow BRIGADE WSB Insecticide/Miticide to contact fruit or foliage. Contact with fruit may result in illegal residues.
- Do not apply more than 80 oz product (0.5 lb ai) per acre per year.
- Do not apply within 1 day of harvest.
- Do not apply by air.
- Weed control around trees is essential for better performance.

All Applicable Directions, Restrictions, Worker Protection Standards Requirements and Precautions on the EPA Registered BRIGADE® WSB INSECTICIDE/MITICIDE (EPA REG. NO. 279-3108) label MUST be followed.
This Recommendation MUST be in the Possession of the USER at the time of Pesticide Application.