



**FIFRA 24(c)
Special Local Need
Label**

EPA SLN No.: LA-210002
EPA Reg. No. 279-3158

**FOR DISTRIBUTION AND USE
ONLY IN LOUISIANA**

This label for Command® 3ME Microencapsulated Herbicide is valid until October 5, 2027, or until withdrawn, canceled, or suspended.

For Resistant Italian Ryegrass Management in Fallow Fields

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FALLOW APPLICATION:

Command® 3ME microencapsulated herbicide may be used as a residual preemergence treatment to control resistant Italian ryegrass during the fallow period following crop harvest and before the following crop is planted.

Application use rates by soil texture:

Fine or heavy textured soils: 21.3 to 28 fl. oz./A (0.50 to 0.66 lbs. ai/A)

Medium textured soils: 16 to 24 fl. oz./A (0.375 to 0.56 lbs. ai/A)

Application timing:

Apply prior to Italian ryegrass emergence in the fallow period from October 1st through November 30th. If Italian ryegrass is already present, include an effective postemergence grass herbicide with the application.

Application volume:

Ground: Apply in a minimum of 10 gallons of finished spray per acre. **Air:** Apply in a minimum of 5 gallons of finished spray per acre.

PRECAUTIONS

Application of Command® 3ME microencapsulated herbicide to fields which have been precision leveled with deep cuts may result in rice crop injury including stand loss. Consult with rice specialists for soil amending practices which can reduce potential for herbicide injury in precision leveled fields.

24(c) Registrant:



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104

062722

Command and FMC – Trademarks of FMC or an affiliate
©2022 FMC Corporation

Some conditions may lead to temporary whitening or yellowing of leaves in approved rotational crops where soil residues of Command® 3ME microencapsulated herbicide exist. Factors that contribute to carryover include, but are not limited to, over-application, soil pH ≤ 5.9, extreme dryness in the four months following application, and/or choice of crop cultivar/hybrid.

RESTRICTIONS:

- **DO NOT** apply on coarse or medium textured soils when planting to rice the following spring.
- **DO NOT** apply to fallow fields in which concurrent crayfish or catfish farming is included in the cultural practices.
- **DO NOT** apply more than a maximum cumulative amount of Command® 3ME microencapsulated herbicide or any product containing clomazone in a 12-month period per crop referenced in the table below. For crops not listed, refer to the Section 3 label:

Crop	Rate (fl oz) Per Acre	Rate (lb ai) Per Acre
Cotton	53.3	1.3
Edamame	21.3	0.5
Rapeseed	4.5	0.1
Rice	34.1	0.8
Soybean	53.3	1.25
Sugarcane	53.3	1.25
Sweet Potato	53.3	1.3

Observe all buffer restrictions noted in the Command® 3ME microencapsulated herbicide in the Application Precautions section below.

APPLICATION PRECAUTIONS:

Caution must be taken to minimize off target drift of Command® 3ME microencapsulated herbicide as off-site movement can cause foliar whitening or yellowing of some plants.

DO NOT apply within 300 feet of downwind crops and desirable non-target plants.

DO NOT apply within 1,200 feet of Towns and Housing Developments, Commercial Fruit, Nut, or Vegetable Production¹, Commercial Greenhouses or Nurseries.

¹Except for vegetable crops registered for use on the Command® 3ME microencapsulated herbicide label.

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below, otherwise crop injury may occur.

Note: When using Command® 3ME microencapsulated herbicide with other registered herbicides always refer to rotational restrictions and precautions on the other product's label.

ANYTIME - Cotton*, Peas, Peppers, Pumpkins (processing), **Rice, Soybeans, Squash, Sweet Potatoes**, Tobacco and Tuberous and Corm Vegetables

9 MONTHS - Cotton, Corn (Field, Pop, Seed, Sweet), Cucurbits, Dry Beans, Peanuts, Potatoes, Snap Beans, Sorghum, Sugar Beets, Tomatoes (Transplanted)

12 MONTHS - All other crops

* Refer to "REQUIREMENTS FOR PLANTING TIME APPLICATIONS" and "REPLANTING INSTRUCTIONS" in the Command® 3ME microencapsulated herbicide federal label.

24(c) Registrant:



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104

062722

Command and FMC – Trademarks of FMC or an affiliate
©2022 FMC Corporation