



**FIFRA 24(c)
Special Local Need
Label**

EPA SLN No.: AR-260001
EPA Reg. No. 279-3158

**FOR DISTRIBUTION AND USE
ONLY IN ARKANSAS**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

**FOR APPLICATION OF COMMAND 3ME MICROENCAPSULATED
HERBICIDE POSTEMERGENCE OVER SOYBEAN**

Directions For Use

Command 3ME Microencapsulated Herbicide may be applied postemergence over-the-top of soybean at 12.8 to 32 fl oz/A (0.3 to 0.75 lb ai/A) from emergence up to the V6 soybean growth stage for preemergence and residual control of prickly sida (teaweed) and other labeled grass and broadleaf weed species. Command 3ME Microencapsulated Herbicide may be tank mixed with other soybean herbicides to broaden the weed control spectrum. To control emerged weeds at the time of application, Command 3ME Microencapsulated Herbicide must be tank mixed with another herbicide labeled for control of the emerged weed(s) present.

Use higher use rates (20 to 32 fl oz/A) for high weed pressure locations, when soil texture is medium to fine and/or when extended residual weed control is needed for soybeans to reach canopy closure.

SOIL TEXTURE CLASSIFICATION.

Coarse	Medium	Fine
Sand, Loamy sand and Sandy loam	Loam, Silt loam, Sandy clay loam and Silt	Sandy clay, Silty clay loam, Silty clay, Clay loam and Clay

24(c) Registrant:



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

ROTATIONAL CROP RESTRICTIONS

Rotate to crops as listed under Table 4. Rotational Crop Restrictions.

Cover crops may be planted anytime but stand reductions may occur in some areas. Refer to the ROTATIONAL CROP RESTRICTIONS table footnote for more information.

RESTRICTIONS

- **DO NOT** allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.
- **DO NOT** graze or harvest for food or feed cover crops planted less than 9 months after Command 3ME Microencapsulated Herbicide treatment.
- **DO NOT** apply after V6 soybean growth stage.
- **DO NOT** replant treated fields with any crop at intervals which are inconsistent with the Rotational Crop Instructions on this label.
- **DO NOT** apply more than 53.3 fl oz/A (1.25 lb ai/A) Command 3ME Microencapsulated Herbicide per application.
- **DO NOT** apply more than 53.3 fl oz/A (1.25 lb ai/A) Command 3ME Microencapsulated Herbicide per year.
- **DO NOT** soil incorporate herbicide tank mixtures of Command 3ME Microencapsulated Herbicide and other herbicides.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Aircraft

Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

PRECAUTIONS:

Caution must be taken to minimize off target drift of Command 3ME Microencapsulated Herbicide as off-site movement can cause foliar whitening or yellowing of some plants.

24(c) Registrant:



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

RESTRICTIONS:

**Do not apply within 300 feet of downwind crops and other desirable non-target plants.
Do not apply within 1,200 feet of Towns and Housing Developments, Commercial Fruit, Nut, or Vegetable Production¹, Commercial Greenhouses or Nurseries.**

¹ Except for vegetable crops registered for use on the Command 3ME Microencapsulated Herbicide label (including peppers, pumpkins (processing), succulent peas sweet potato and winter squash).

With split applications do not apply more than a total of 53.3 fl oz/A (1.25 lb ai/A) Command 3ME Microencapsulated Herbicide per year.

Do not use water containing Command 3ME Microencapsulated Herbicide residues from soybean cultivation to irrigate food or feed crops, which are not registered for use with Command 3ME Microencapsulated Herbicide.

REPLANTING INSTRUCTIONS

If initial seeding of soybean fails to produce a stand, soybean may be replanted in fields treated with Command 3ME Microencapsulated Herbicide alone (or with specified tank mixtures). When tank mixing with a labeled product, refer to the soybean replant instructions for that product.

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed under Table 4. Rotational Crop Restrictions on the Command 3ME Microencapsulated Herbicide label.

Cover crops may be planted anytime but stand reductions may occur in some areas. Refer to the ROTATIONAL CROP RESTRICTIONS table footnote for more information.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

R-4990 010926 09-22-25

24(c) Registrant:



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000