RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN Arizona and California

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

EPA Reg. No. 279-3126

For Celery and Lettuce to be Exported to Canada

CROP	PEST	RATE OF APPLICATION
Celery Lettuce (Head and Leaf)	See Product Label	2.4 - 4.3 fl oz/A (0.028 to 0.05 lb ai/A)

DIRECTIONS FOR USE: This use pattern is designed to meet the Canadian Default MRL. Follow all instructions, recommendations and restrictions on the Mustang Insecticide label with the following modifications for produce to be exported to Canada.

- PHI: Do not apply within 30 days of harvest.
- **Maximum Amount per Season:** Do not exceed 3 applications of Mustang Insecticide or 12.9 fl oz of product (0.15 pounds of active ingredient) per acre per season.
- Do not treat vegetables that will be harvested as "baby vegetables".
- **Do not** treat spinach
- **Do not** use Mustang® Max or Mustang® Maxx or products containing cypermethrin or alpha-cypermethrin on crops treated with Mustang Insecticide (combined residues may exceed the Canadian Default MRL).

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED MUSTANG INSECTICIDE LABEL (EPA REG. NO. 279-3126) MUST BE FOLLOWED. THIS RECOMMENDATION LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

