## THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



For Agricultural or Commercial Use Only EPA Registration No. 279-3241

## 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF GEORGIA

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

CROP	PEST	RATE OF APPLICATION (For control of weeds)
Cucurbits (transplanted only), Fruiting Vegetables (transplanted only) planted into plastic mulch	Annual and perennial broadleaf weeds (see section 3 label for specific weeds controlled)	0.5 to 2.0 fl. ozs. (0.088 to 0.031 pounds active ingredient) per acre

## DIRECTIONS FOR USE:

Apply Aim EC Herbicide alone or with other registered herbicides as a burn-down treatment prior to transplant of labeled crops into plastic mulch to control or suppress problem weeds. Aim EC Herbicide may be used as a burn-down treatment for previous crops prior to new planting. Apply Aim EC Herbicide alone at rates up to 2.0 fl. ozs. (0.031 pound active ingredient) per acre or at 0.5 to 1.0 fl. ozs. (0.088 to 0.016 pound active ingredient) per acre in combination with labeled glyphosate products.

For optimum performance, make applications to actively growing weeds up to 4 inches in height or rosettes less than 3 inches across. Coverage is essential for good control. When applying Aim EC Herbicide into plastic mulch prior to transplant, care must be taken to remove residues from the plastic prior to transplant. A single 0.5 inch water event, either by natural rainfall or sprinkler irrigation and a 7 day interval between application and transplant is required to remove residues. Care must be taken to insure that wash water flushed off the plastic mulch does not enter the transplant holes.

## RESTRICTIONS:

Do not apply more than 0.096 pounds active ingredient per acre per season (6.1 fl. ozs. per acre) of Aim EC.

Refer to respective tank-mix partner labels and follow the most restrictive labeling regarding plant back restrictions, rotational guidelines, methods of applications, and surfactant requirements of the tank mixture components.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED AIM EC HERBICIDE LABEL MUST BE FOLLOWED.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

