

**INTENDED FOR AGRICULTURAL OR COMMERCIAL USE**

EPA Reg. No. 279-3241

EPA Est 279-IL-1

**Active Ingredient:****By Wt**

Carfentrazone-ethyl ..... 22.3%

**Other Ingredients** ..... 77.7%

Total ..... 100.0%

This product contains 2.0 pounds active ingredient per gallon.

Contains Petroleum Distillates

U.S. Patent No. 5,125,958

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**FIRST AID**

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**If in Eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If Swallowed:** Call a poison control center or doctor immediately for treatment advice. **DO NOT** give any liquid to the person. **DO NOT** induce vomiting unless told to do so by the poison control center or doctor. **DO NOT** give anything by mouth to an unconscious person.

**HOTLINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

**Note to Physician:** Carfentrazone-ethyl is expected to have low oral and dermal toxicity, and moderate inhalation toxicity. It is expected to be slightly irritating to the skin and minimally irritating to the eyes. Treatment is otherwise controlled removal of exposure followed by symptomatic and supportive care.

See other panels for additional precautionary information.

**ACTIVE INGREDIENT MADE IN CHINA, FORMULATED AND PACKAGED IN USA.**

**Sold By**



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

**Net Contents: 1 Quart**

Label Code SL-4868A 022124 2-28-24

## PRECAUTIONARY STATEMENTS

### Hazards to Humans (and Domestic Animals)

#### CAUTION

Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

#### Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

#### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Carfentrazone-ethyl is very toxic to algae and moderately toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark, except as specified on this label. **DO NOT** contaminate water when disposing of equipment washwaters.

#### Fish Advisory Statement:

This product may be hazardous to aquatic organisms, particularly in clear, shallow water bodies that are adjacent to treated areas. Transport to water by runoff or spray drift of this product in areas where surface water is present, or intertidal areas below the mean high water mark, should be avoided. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

#### For ground water:

Residues of this chemical have properties and characteristics associated with chemicals detected in ground water. Residues of this chemical may leach into ground water if the chemical is used in areas where soils are permeable, particularly where the water table is shallow.

#### For surface water:

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of carfentrazone-ethyl residues from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

#### Non-Target Organism Advisory Statement:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by minimizing spray drift.

#### Physical/Chemical Hazards

**DO NOT** use or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Use Restrictions:

Only use for sites, pests and applications methods specified on this labeling.

**DO NOT** apply this product through any type of irrigation system.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### Endangered Species:

It is a Federal offense to use any pesticide in a manner that results in the death of an endangered species. Use of this product may pose a hazard to endangered or threatened species. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county in which you are applying the product. To obtain Bulletins, no more than six months before using this product, consult <http://www.epa.gov/espp/> or call 1-800-447-3813. You must use the Bulletin valid for the month in which you will apply the product.

## WEED RESISTANCE MANAGEMENT

For resistance management, AIM EC HERBICIDE is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to AIM EC HERBICIDE and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of AIM EC HERBICIDE or other Group 14 herbicides within a growing cycle sequence or among growing cycles with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact FMC Corporation at 1-800-331-3148

## SPRAY DRIFT

### Aerial Applications:

- For aerial applications, the distance of the outer most nozzles on the boom must not exceed 75% of the length of the wingspan or 90% of rotor diameter. To further reduce drift, use on half of the length of the wingspan or rotor diameter at the edge of the field.
- Applicators must only spray when wind speed is 10 miles per hour or less.
- Applicators must not spray during temperature inversions.
- For aerial applications, the release height must be no higher than 10 feet from the top of the crop canopy, unless a greater application height is required for pilot safety.
- For aerial applications, select nozzle and pressure that produce medium to coarse spray droplets as indicated in nozzle manufacturer's catalogues and in accordance with ASABE Standard 572.1.

### Ground Boom Applications:

- For ground boom applications, apply with the nozzle height no more than 4 feet above the ground or crop canopy. For all other ground applications, the nozzle must be no more than 4 feet from the target vegetation.
- For ground applications, select nozzle and pressure that produce medium to coarse spray droplets as indicated in nozzle manufacturer's catalogues and in accordance with ASABE Standard 572.1.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

### Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is: Long sleeve shirt and pants, waterproof gloves, and shoes plus socks.

## Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Re-entry Statement: DO NOT** allow people (other than applicator) or pets on treatment area during application. **DO NOT** enter treatment area until spray has dried.

## STORAGE AND DISPOSAL

**DO NOT contaminate water, food, or feed by storage and disposal.**

### Pesticide Storage

Not for use or storage in or around the home.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. **DO NOT** put formulated or dilute material into food or drink containers. **DO NOT** contaminate other pesticides, fertilizers, water, food, or feed by inappropriate storage or disposal.

In case of spill, avoid contact, isolate area and keep out unprotected persons and animals. Confine spills. **Call CHEMTREC (Transportation and spills): (800) 424-9300.**

To confine spill: Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a larger holding container. Identify contents per required hazardous waste labeling regulations.

### Pesticide Disposal

Waste resulting from the use of this product may be disposed of at an approved waste disposal facility.

### Container Handling

Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: (For containers greater than 5 gallons) Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. (For containers 5 gallons or less) Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Triple rinse (or equivalent). Then offer for recycling if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Returnable/Refillable Containers** - Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. If unable to return or refill, offer for recycling if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## PRODUCT INFORMATION

AIM EC HERBICIDE is an emulsifiable concentrate formulation. AIM EC HERBICIDE is to be mixed with water, liquid fertilizer or mixtures of water and liquid fertilizer and adjuvants and applied to labeled crops and non-crop areas for selective postemergence control of broadleaf weeds, for sucker control, for burndown prior to planting, as a harvest aid and to defoliate/desiccate labeled crops.

Weed control is optimized when the product is applied to actively growing weeds. AIM EC HERBICIDE is a contact herbicide. Within a few hours following application, the foliage of susceptible weeds shows signs of desiccation.

Extremes in environmental conditions including temperature, moisture, soil conditions, and cultural practices may affect the activity of AIM EC HERBICIDE. Herbicide symptoms may be accelerated under moist conditions. Weed control may be reduced when weeds are hardened off by drought and become less susceptible to AIM EC HERBICIDE.

AIM EC HERBICIDE is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications must not be made within 6 to 8 hours of either rain or irrigation or when heavy dew is present on the crop. Environmental conditions and certain spray tank additives may increase herbicidal symptoms on the crop.

## TANK MIXTURES

(See specific crop application instructions for additional tank mixture restrictions and information)

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

AIM EC HERBICIDE may be tank-mixed with other registered herbicides for controlling a broader spectrum of weeds. Refer to this and other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank mix partner. When preparing a new tank mix, conduct an appropriate compatibility test by mixing proportional amounts of all spray ingredients in a test vessel (jar) prior to tank mixing with other products. Shake the mixture vigorously and allow it to stand for five to ten minutes. Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and must not be applied. Provided the jar test indicates the mixture to be compatible, prepare the tank mixture as follows: Fill the tank one fourth full with water. With the agitator operating, add the specified amounts of ingredients using the following order: dry granules first and liquid suspensions (flowables) second. As the agitation continues and the tank is filled with water add emulsifiable concentrate products third followed by the addition of water soluble products.

## ADJUVANT USE REQUIREMENTS

The use of a quality spray adjuvant is required for optimum performance. Refer to the individual crop sections of this label for specific adjuvant type and use rates.

## ON-FARM TESTING

Not all varieties or cultivars of labeled crops have been fully evaluated under all environmental and soil conditions. Consult with your local seed company for additional information.

It may also be beneficial to conduct small on-farm trials under actual conditions with specific varieties or cultivars before treating large acreage.

## APPLICATION INFORMATION

### Mixing and Loading Instructions:

Start by filling the tank with  $\frac{3}{4}$  of the desired volume of clean water and, with agitation, add the proper amount of AIM EC HERBICIDE. Complete filling the spray tank to the desired volume. Maintain sufficient agitation to keep materials in solution during both mixing and application and until the spray tank has been emptied. For tank mixtures, follow your local extension guidelines for mixing order. General guidelines are: add dry materials first and agitate until mixed; then EW or water soluble liquids; then EC formulations; then, add adjuvants last. Ensure the compatibility of other products and/or liquid fertilizers with AIM EC HERBICIDE before mixing them together in the spray tank.

### Mixing Precautions

Avoid the overnight storage of AIM EC HERBICIDE spray mixtures. If spray solution is stored overnight or longer, thoroughly agitate spray mixture before applying the solution. Premixing AIM EC HERBICIDE spray solutions in nurse tanks is not specified. Maintain continuous and adequate spray solution agitation until all the spray solution has been used. **DO NOT** use with tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer spray solutions if necessary to maintain the pH between 5-8.

## SPRAY EQUIPMENT CLEAN-OUT

**Many pesticides are very active at low rates, especially to sensitive crops.** Residues left in mixing equipment, spray tanks, hoses, spray booms and nozzles can cause crop effects if they are not properly cleaned. As soon as possible after spraying AIM EC HERBICIDE and before using the sprayer equipment for any other applications, the sprayer equipment must be thoroughly cleaned using the following procedure. In addition, users must take appropriate steps to ensure proper equipment clean-out for any other products mixed with AIM EC HERBICIDE as required on the other product labels. More complete cleaning can be achieved if the spray system is cleaned immediately following the application.

1. Drain sprayer tank, hoses, spray boom and spray nozzles. Use a high-pressure detergent wash to remove physical sediment and residues from the inside of the sprayer tank and thoroughly rinse. Then, thoroughly flush sprayer hoses, spray boom and spray nozzles with a clean water rinse. Remove and clean spray tips and all filters and screens (tank, spray hose and spray tips) separately in the ammonia solution of Step 2.
2. Next, prepare a sprayer cleaning solution by adding three gallons of ammonia (containing at least 3% active) per 100 gallons of clean water. Prepare sufficient cleaning solution to allow the operation of the spray system for a minimum of 15 minutes to thoroughly flush hoses, spray boom and spray nozzles.
3. Convenient and thorough cleaning of the sprayer can be achieved if the ammonia solution or fresh water is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.
4. Before using the sprayer, completely drain the sprayer system. Rinse the tank with clean water and flush through the hoses, spray boom, and spray nozzles with clean water.
5. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State, and local regulations and guidelines.

**DO NOT** apply sprayer cleaning solutions or rinsate to sensitive crops.

**DO NOT** store the sprayer overnight or for any extended period of time with AIM EC HERBICIDE spray solution remaining in the tank, spray lines, spray boom plumbing, spray nozzles or strainers.

If the sprayer has been stored or idle, purge the spray boom and nozzles with clean water before beginning any application.

Should small quantities of AIM EC HERBICIDE remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. FMC accepts no liability for any effects due to inadequately cleaned equipment.

## APPLICATION METHODS

### GROUND APPLICATION

Use ground sprayers designed, calibrated and operated to deliver uniform spray droplets to the targeted plant or plant parts. Adjust sprayer nozzles to achieve uniform plant coverage. Overlaps and slower ground speeds (caused by continuing to spray while starting, stopping or turning) may result in higher application rates and possible crop response.

### Spray Buffer for Ground Application

Spray buffer zones for ground applications, listed in chart below, are required near desirable perennial vegetation or crops before blossom and after total leaf drop, and/or near other desirable or annual crops.

Buffers		
AIM EC HERBICIDE USE RATE (lbs ai/A)	Low Spray Boom Buffer (ft.)	High Spray Boom Buffer (ft.)
0.024	20	33
0.031	26	46

### **Broadcast Boom Sprayers**

Use a broadcast boom sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Use nozzles that produce minimal amounts of fine spray droplets. **DO NOT** exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a minimum of 10 gallons of finished spray per acre. Use higher spray volumes when there is a dense weed population or crop canopy. Adjust sprayers to position spray tips no lower than 12-18 inches above the crop or weed canopy depending on the nozzle specification. Operate the sprayer to avoid the application of high herbicide rates directly over the rows or into the whorl of treated crop plants.

### **Directed Sprayers**

For directed sprayers apply AIM EC HERBICIDE with drop nozzles or other post direct spray equipment.

**Post directed** applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. **DO NOT** apply when conditions favor drift or when wind speed is above 10 MPH.

Use drop nozzles or other spray equipment capable of directing the spray to target weeds and away from sensitive plant parts. Apply when labeled crops have reached minimum growth stages described in specific crop sections of this label and when spray will not be deposited on green stems, foliage, blossoms or fruit.

### **Hooded Sprayers**

To apply AIM EC HERBICIDE using a hooded sprayer, refer to the Hooded Sprayer Section for specific adjustment and operation instructions. For additional information, refer to the individual crop sections of this label.

### **Hand held or high volume orchard gun sprayers**

AIM EC HERBICIDE may be applied to certain labeled crops and non-crop areas with hand operated sprayers including backpack sprayers, compression sprayers, knapsack sprayers, or high volume orchard gun sprayers. Directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Refer to individual crop sections of this label.

### **AERIAL APPLICATION**

Use nozzle types and arrangements that will provide optimum coverage while producing a minimal amount of fine droplets. Apply at a minimum of 3 gallons of finished spray per acre. Spray volumes greater than 3 GPA may be needed for harvest aid and defoliation treatments, or for dense weed populations or with dense crop canopies.

#### **For Aerial Application in California:**

#### **Refer to individual crop sections to see if application is permitted by air**

For applications near desirable perennial vegetation or crops before blossom and after total leaf drop, and/or near other desirable or annual crops:

- **DO NOT** apply within 100 feet of all desirable vegetation or crops.
- If wind up to 10 miles per hour is blowing toward desirable vegetation or crops, **DO NOT** apply within 500 feet of the desirable vegetation or crops.
- **DO NOT** apply when winds are in excess of 10 mph or when inversion conditions exist.

### **SPRAY DRIFT MANAGEMENT**

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

AIM EC HERBICIDE is a contact PPO (protoporphyrinogen oxidase) herbicide. Avoid any drift conditions that would allow the product to contact desirable vegetation. AIM EC HERBICIDE is not volatile, however; mist from spray drift may cause injury to sensitive plants.

The interaction of equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from applications to agricultural field crops. These requirements **DO NOT** apply to forestry applications, public health uses or to applications of dry materials.

Where states have more stringent regulations, they must be observed.

## INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The optimum drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift when applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

For all non-aerial applications, wind speed must be measured adjacent to the application site, on the upwind side, immediately prior to application.

### Controlling Spray Droplet Size

**Volume** – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows usually produce larger droplets.

**Pressure** – **DO NOT** use pressures greater than that specified by the nozzle manufacturer. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

**Number of Nozzles** – Use the minimum number of nozzles that provide uniform coverage.

**Nozzle Orientation** – For aerial application, orient nozzles so that the spray is released parallel to the airstream. A parallel orientation results in larger droplets than other orientations and reduces air turbulence and the production of small droplets. Significant deflection from horizontal will reduce droplet size and increase drift potential.

**Nozzle Type** – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low drift nozzles. For aerial applications, solid stream nozzles oriented straight back produce the largest droplets and potentially the least drift.

**Boom Length** – For some aerial use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

**Application Height** – Making applications at the lowest height that is safe reduces exposure of spray droplets to evaporation and wind movement. Aerial applications must not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety.

**Swath Adjustment** – Swath adjustment distance must increase, with increasing drift potential (higher wind, smaller drops, etc.).

**Drift Reduction Technology (DRT)** – The EPA Drift Reduction Technology (DRT) Program was developed to encourage the manufacture, marketing, and use of spray technologies scientifically verified to significantly reduce pesticide drift. The use of DRTs should result in significantly less pesticide from spray applications drifting and being deposited in areas not targeted by those applications, compared to spray technologies that **DO NOT** meet the minimum DRT standard. EPA-verified drift reduction technologies (DRTs) and their ratings will be added to the following webpage as they become available: <https://www.epa.gov/reducing-pesticide-drift/epa-verified-and-rated-drift-reduction-technologies>

**Wind** – Drift potential is lowest between winds speeds of 3 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. Applications shall be avoided below 3 mph due to variable wind direction and high inversion potential. **DO NOT** apply AIM EC HERBICIDE when wind speed exceeds 10 mph. NOTE: Local terrain can influence wind patterns. Every applicator shall be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity** – When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions** – **DO NOT** apply AIM EC HERBICIDE during a temperature inversion because the drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the following morning. Their presence can be indicated by ground fog. However, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Shielded Sprayers** – Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

**Sensitive Areas** – AIM EC HERBICIDE shall only be applied when the wind is blowing away from adjacent sensitive areas (e.g. residential areas, bodies of water, known habitats for threatened or endangered species and non-target crops).

# ALLOWABLE AIM EC HERBICIDE USE INFORMATION

Refer to the crop section of this label for specific product use directions.

Table 1:

Maximum Allowable AIM EC HERBICIDE Use Per Acre Per Year* for Crop or Subgroups		
Crop Group/Subgroup	Maximum Rate AIM EC HERBICIDE (fl oz/A) Per Year	Maximum Rate AIM EC HERBICIDE (lb ai/A) Per Year
Alfalfa and Clover (Group 18)	2.5	0.04
Alfalfa and Clover (Group 18) harvest aid only	3.8	0.06
Asparagus	3.8	0.06
Banana	7.9	0.123
Berry, low growing (Subgroup 13-07G)	6.1	0.095
Bushberry (Subgroup 13-07B)	6.1	0.095
Cacao	7.9	0.123
Caneberry (Subgroup 13-07A)	25.6	0.4
Citrus fruit (Group 10-10)	7.9	0.123
Coconut	7.9	0.123
Coffee	7.9	0.123
Corn	2.0	0.031
Cotton	7.9	0.123
Cotton, harvest aid only	3.2	0.05
Date	7.9	0.123
Fig	7.9	0.123
Fruit, small vine climbing – except fuzzy kiwifruit (Subgroup 13-07F)	7.9	0.123
Globe Artichoke	6.1	0.095
Grass (Group 17)	5.9	0.092
Guayule	7.9	0.123
Herbs and Spices (Group 19)	6.1	0.095
Hops	7.6	0.12
Horseradish	6.1	0.095
Indian Mulberry	7.9	0.123
Kiwifruit	7.9	0.123
Mint	1.92	0.030
Nut, Tree (Group 14-12)	7.9	0.123
Oil Seed – except cottonseed (Group 20)	6.1	0.095
Olive	7.9	0.123
Palm Heart	7.9	0.123
Peanut	6.1	0.095
Peanut (harvest aid)	2.0	0.031
Persimmon	7.9	0.123
Pome fruit (Group 11-10)	7.9	0.123
Pomegranate	7.9	0.123
Rice (Southern Use Only)	8.8	0.138
Rice, harvest aid only (not permitted in California)	1.47	0.023
Small Grains	1.0	0.016
Small Grains (except winter wheat)	2.0	0.031
Small Grains (winter wheat)	2.0	0.031
Sorghum (harvest aid)	1.0	0.016
Sorghum (grown for seed and grain)	1.0	0.016
Soybeans (preplant and in-season and harvest aid)	1.47	0.023
Stone fruit (Group 12-12)	7.9	0.123
Sugarcane	6.1	0.095
Sugarcane (harvest aid)	2.0	0.031
Tea	7.9	0.123
Teff	2.0	0.031
Tobacco	3.2	0.05
Tropical fruit Trees	6.1	0.095
Vanilla	7.9	0.123
Vegetable, brassica (Group 5)	6.1	0.095
Vegetable, bulb (Group 3-07)	6.1	0.095
Vegetable, cucurbit (Group 9)	6.1	0.095
Vegetable, foliage of legume (Group 7)	6.1	0.095
Vegetable, fruiting (Group 8-10)	6.1	0.095
Vegetable, leafy except Brassica (Group 4)	6.1	0.095
Vegetable, leaves of root and tuber (Group 2)	6.1	0.095
Vegetable, legume (Group 6 - except soybean)	6.1	0.095
Vegetable, root (Subgroups 1A and 1B)	6.1	0.095
Vegetable, tuberous and corm (Subgroups 1C and 1D)	11.6	0.181
Wild Rice	19.2	0.3

\*The total allowable usage includes all applications made to the field per calendar year. This includes fallow treatments, burndown treatments and all in-crop treatments, including harvest aid.

## PREHARVEST INTERVALS

Refer to the crop section of this label for specific product use directions.

Table 2:

Preharvest Intervals (PHI) or Maximum Growth Stage for AIM EC HERBICIDE Applications	
Crop Group/Subgroup	PHI(Days Before Harvest) or Growth Stage
Alfalfa and Clover (Group 18) grown for Forage and /or Hay	21
Alfalfa and Clover (Group 18) grown for Seed	3
Asparagus	5
Banana	3
Berry, low growing (Subgroup 13-07G)	0
Bushberry (Subgroup 13-07B)	0
Cacao	3
Caneberry (Subgroup 13-07A)	15
Citrus fruit (Group 10-10)	3
Coconut	3
Coffee	3
Corn	14 Leaf Collars
Corn, Sweet corn grown for seed, popcorn, field corn (harvest aid)	3
Cotton (harvest aid)	7
Cotton (preplant and in-season)	7
Date	3
Fruit, small vine climbing – except fuzzy kiwifruit (Subgroup 13-07F)	3
Fig	3
Globe Artichoke	0
Grass (Group 17)	0
Guayule	3
Herbs and Spices (Group 19)	0
Hops	7
Horseradish	0
Indian Mulberry	3
Kiwifruit	3
Mint	5
Nut, Tree (Group 14-12)	3
Oil Seeds (Group 20 – except cottonseed)	0
Olive	3
Palm Heart	3
Peanut	7
Persimmon	3
Pome fruit (Group 11-10)	3
Pomegranate	3
Rice (southern use only)	60
Rice, harvest aid only (not permitted in California)	3
Small Grains (Except winter wheat)	Jointing Stage
Small Grains (harvest aid) – include Winter Wheat	7
Sorghum (harvest aid)	3
Sorghum (preplant and in-season)	14 Leaf Collars Stage
Soybean (harvest aid)	3
Soybeans (grown for seed and grain)	V10
Stone fruit (Group 12-12)	3
Sugarcane	7
Tea	3
Teff	Jointing Stage
Teff (forage - harvest aid)	7
Teff (grain - harvest aid)	3
Tobacco	6
Tropical fruit	0
Vanilla	3
Vegetable, brassica (Group 5)	0
Vegetable, bulb (Group 3-07)	0
Vegetable, cucurbit (Group 9)	0
Vegetable, foliage of legume (Group 7)	0
Vegetable, fruiting (Group 8-10)	0
Vegetable, leafy except brassica (Group 4)	0
Vegetable, leaves of root and tuber (Group 2)	0
Vegetable, legume (Group 6 – except soybean)	0
Vegetable, root (Subgroups 1A and 1B)	0
Vegetable, tuberous and corm (Subgroups 1C and 1D)	7
Wild Rice	60

## CROP ROTATIONAL RESTRICTIONS

Following an application of AIM EC HERBICIDE, a treated field may be rotated to a registered crop at any time, subject to specific crop restrictions that may be found in the individual crop sections. All other crops may be planted after 12 months.

Up to 12 months following application to cotton, potato and the non-grass animal feed crop group 18, the subsequent planted crop may only be a registered crop.

## WEED CONTROL

When used as directed, AIM EC HERBICIDE will provide control of the listed weeds up to four (4) inches in height, or as specified.

Table 3:

Weeds Controlled	AIM EC HERBICIDE Use Rate fl oz/A (lb ai/A)
Lambsquarters, common (up to 3 inches tall)	0.5 fl oz/A (0.008 lb ai/A)
Morningglory, ivyleaf (up to 3 leaves)	
Morningglory, pitted (up to 3 leaves)	
Nightshade, Eastern black	
Pigweed, redroot	
Velvetleaf	
Waterhemp (up to 2 inches tall)	
<b>Weeds Controlled</b>	<b>AIM EC HERBICIDE Use Rate fl oz/A (lb ai/A)</b>
All the weeds controlled at 0.5 fl oz/A (0.008 lb ai/A) plus the weeds listed below:	0.8 fl oz/A (0.013 lb ai/A)
Cheeseweed	
Filaree, redstem	
Flixweed	
Lambsquarters, common	
Mallow, common	
Morningglory, entireleaf	
Morningglory, ivyleaf	
Morningglory, pitted	
Morningglory, scarlet	
Nightshade, hairy	
Pennycress, field	
Pigweed, prostrate	
Pigweed, smooth	
Pigweed, tumble	
Purslane, common	
Sesbania, hemp	
Smartweed, PA (seedling)	
Spurge, prostrate	
Tansymustard	
Velvetleaf (24")	
Waterhemp, common & tall	
<b>Weeds Controlled</b>	<b>Use Rate fl oz/A (lb ai/A)</b>
All the weeds controlled at 0.8 fl oz/A (0.013 lb ai/A) plus the weeds listed below:	1.0 fl oz/A (0.016 lb ai/A)
Amaranth, spiny	
Anoda, spurred	
Bedstraw, catchweed	
Buffalobur	
Carpetweed	
Cocklebur	
Copperleaf, hophornbeam	
Cotton, GMO Varieties	
Cotton, volunteer	
Eclipta	
Fiddleneck, coast	
Groundcherry, smooth (seedling)	
Groundcherry, Wright's	
Jimsonweed	
Kochia (Non PPO resistant)	
Lettuce, Prickly 2-3 leaf	
Nettle, burning	
Nightshade, American black	
Nightshade, black	
Rocket, London	
Shepherdspurse	
Speedwell, Virginia	
Spiderwort, tropical	
Thistle, Russian (up to 2 inches tall)	
Wallflower, bushy	

Weeds Controlled	Use Rate fl oz/A (lb ai/A)
All the weeds controlled at 1.1 fl oz/A (0.016 lb ai/A) plus the weeds listed below:	1.6 fl oz/A (0.025 lb ai/A)
Amaranth, Palmer	
Corn Spurry	
Filaree, broadleaf	
Filaree, white	
Lettuce, prickly	
Mallow, Venice (up to 2 inches tall)	
Meadowfoam	
Redmaids	

**Burndown of top growth**

Weed List	AIM EC HERBICIDE Use Rate fl oz /A (lb ai/A)
Bindweed, field	1.0 - 2.0 fl oz/A (0.016 – 0.032 lb ai/A)
Burclover	
Dayflower	
Sage, lanceleaf	
Sowthistle	

**AGRICULTURE FARM AND FARMSTEAD USE – NON-CROP**

Timing and Method of Application

AIM EC HERBICIDE may be used for broadleaf weed control on farms and farmsteads in areas outside of crop growing areas. See the rate and weed table 3 to determine the proper rate for areas including grass waterways, field edges, terraces, equipment storage areas, shelter belts, fence lines, farm buildings, dry ditch, canal banks etc. AIM EC HERBICIDE is a contact herbicide and coverage is essential for good weed control. AIM EC HERBICIDE will control emerged weeds only. Weeds that germinate after application will require repeat treatments.

**BOOM EQUIPMENT**

**AIM EC HERBICIDE use Rate – Boom Equipment:**

Apply AIM EC HERBICIDE at up to 2.0 fl oz/A (0.031 lb ai/A).

Adjuvant Requirements – Boom Equipment

A nonionic surfactant, crop oil concentrate or methylated seed oil is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed based crop oil concentrate (COC) at 1.0 to 2 % v/v (1.0 to 2.0 gallons per 100 gallons of spray solution) or a methylated seed oil (MSO). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons) or ammonium sulfate at 2 to 4 pounds per acre in addition to the selected NIS, MSO or COC is allowed.

Tank Mixes – Boom Equipment

AIM EC HERBICIDE may be mixed with other herbicides labeled for this method of application in non-crop areas for broader spectrum weed control. See Mixing and Loading Instructions under the PRODUCT INFORMATION section of this label for specific mixing instructions. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner.

**SPOT TREATMENTS (Applications with hand operated sprayer including backpack sprayers, compression sprayers, knapsack sprayers.)**

Mix the amount of AIM EC HERBICIDE for the desired percent spray solution from the table below. These mixtures are based on 1 gallon of solution evenly covering 1000 square feet. Applications must be made on a spray-to-wet basis. Spray coverage must be uniform and complete. **DO NOT** spray to runoff. See Table 4 for weeds controlled at specific concentrations. Use lower concentrations for small seedling weeds at the 2-3 leaf stage. Higher concentrations are needed for larger weeds up to the 6-leaf stage. Applications beyond the 6-leaf stage may result in only partial control. AIM EC HERBICIDE may be mixed with other labeled herbicides including glyphosate, glufosinate, and paraquat for broader spectrum weed control.

**Table 4:**

Desired Volume	Amount AIM EC HERBICIDE				
	0.5 fl oz/A	0.8 fl oz/A	1.0 fl oz/A	1.6 fl oz/A	2.0 fl oz/A
1 Gal	0.4 ml	0.6 ml	0.7 ml	1.1 ml	1.4 ml
5 Gal	1.7 ml	2.7 ml	3.4 ml	5.4 ml	6.8 ml
25 Gal	8.5 ml	13.6 ml	17.0 ml	27.2 ml	34.0 ml

**Adjuvant Requirements:**

A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v having at least 80% active ingredient, or a methylated seed oil (MSO), or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v. A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v or ammonium sulfate (AMS) at the rate of .75 to 1.5 ounces per gallon in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

**Table 5:**

Desired Volume	Adjuvants Rate Conversion				
	NIS	COC or MSO		Liquid Nitrogen	
	0.25%v/v	1.5% v/v	2.0% v/v	2.0% v/v	4.0% v/v
1 Gal	0.35 fl oz	1.9 fl oz	2.5 fl oz	2.5 fl oz	5.0 fl oz
5 Gal	1.6 fl oz	9.6 fl oz	12.8 fl oz	12.8 fl oz	25.6 fl oz
25 Gal	8.0 fl oz	47 fl oz	2 qt	2 qt	4 qt

**PREPLANT BURNDOWN**

Apply AIM EC HERBICIDE alone or with other herbicides or liquid fertilizers as a burn-down treatment to control or suppress weeds. AIM EC HERBICIDE is effective as a burndown treatment for crops prior to new plantings. Apply AIM EC HERBICIDE up to 2.0 fl oz/A (0.031 lb ai/A). **DO NOT** exceed the applicable amounts as listed for the specific crop in the MAXIMUM ALLOWABLE AIM EC HERBICIDE USE in Table 1. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. **Coverage is essential for good control.** Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with labeled burndown herbicides including glyphosate, glufosinate, paraquat, 2,4-D, or dicamba.

<b>Apply AIM EC HERBICIDE as a burndown treatment no later than one (1) day after planting by seed to any of the following crops. (See specific crop section for other timings)</b>
Alfalfa and Clover (Crop Group 18)
Cereal grains (Crop Group 15)
Grasses (Crop Group 17)
Oil Seed (Crop Group 20 – except cottonseed)
Peanut
Soybean
Sugarcane
Vegetables, legume (succulent or dried) (Crop Group 6)
Vegetable, tuberous and corm (Subgroup 1C)
<b>Apply AIM EC HERBICIDE as a burndown treatment no later than one (1) day before transplanting any of the following crops.</b>
Avocado
Banana
Berry, low growing subgroup 13-07G
Cacao
Coconut
Coffee
Date
Fig
Fruit, citrus (Crop Group 10-10)
Fruit, pome (Crop Group 11-10)
Fruit, stone (Crop Group 12-12)
Globe Artichoke
Guayule
Hops
Horseradish
Indian Mulberry
Kiwifruit
Nuts, tree (Crop Group 14-12)
Olive
Palm Heart
Persimmon
Pomegranate
Small Fruit Vine, Climbing – except fuzzy kiwifruit (Subgroup 13-07F)
Tea
Tobacco
Vanilla
Vegetable, fruiting (Crop Group 8-10)

<b>For transplants (not seeded) of the following crops</b>
Vegetables, brassica (Crop Group 5)
Vegetable, cucurbit (Crop Group 9)
Vegetable, fruiting (Crop Group 8-10)
Vegetables, leafy except brassica (Crop Group 4)
<b>Apply AIM EC HERBICIDE as a burndown treatment no later than seven (7) days before planting by seed any of the following crops.</b>
Vegetables, brassica (Crop Group 5)
Vegetable, cucurbit (Crop Group 9)
Vegetable, fruiting (Crop Group 8-10)
Vegetable, leafy except brassica (Crop Group 4)
Vegetable, tuberous and corm (Crop Subgroups 1C and 1D)
<b>Apply AIM EC HERBICIDE as a burndown treatment no later than thirty (30) days before planting by seed any of the following crops.</b>
Sugarbeet
Vegetable, bulb (Crop Group 3-07)

#### Adjuvant Requirements for Preplant burndown

A nonionic surfactant, crop oil concentrate or methylated seed oil is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed based crop oil concentrate (COC) at 1.0 to 2 % v/v (1.0 to 2.0 gallons per 100 gallons of spray solution) or a methylated seed oil (MSO). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons) or ammonium sulfate at 2 to 4 pounds per acre in addition to the selected NIS, MSO or COC is allowed.

#### AIM EC HERBICIDE Plus Glyphosate or Glufosinate

Apply AIM EC HERBICIDE at 0.5 to 1.0 fl oz/A (0.008 to 0.016 lb ai/A) in combination with glyphosate or glufosinate products at their labeled rates for increased speed of activity and improved control of weeds e.g. those listed as follows.

**When applied as directed, AIM EC HERBICIDE plus labeled herbicides e.g. glyphosate, glufosinate, or paraquat will provide increased speed of activity and improved control of weeds listed in Table 6 plus weeds listed in Table 3 for the rate of AIM EC HERBICIDE used.**

**Table 6:**

Buttercup, smallflower	Morningglory, spp.
Chickweed	Pennycress, field
Curled Dock	Prostrate knotweed
Cutleaf Evening Primrose	Purslane, common
Bindweed, field	Smartweed, PA
Dandelion, common	Star-of-Bethlehem
*Fleabane	Shepherdspurse
Groundsel	Tansymustard
Henbit	Thistle, Russian
Kochia (Non PPO resistant)	Thistles, annual & biennial
Lambsquarters, common	Wild buckwheat
*Marestail	Wild hemp

\*glyphosate susceptible marestail and fleabane

**When tank mixing with fertilizer solutions, be sure to prepare an AIM EC HERBICIDE premixture of AIM EC HERBICIDE and clean water.**

For other specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section.

## HOODED SPRAYER APPLICATIONS

Apply AIM EC HERBICIDE to the row middles of the following emerged crops using hooded sprayers in accordance with specific use information in the following **Directions for Use** section.

Apply AIM EC HERBICIDE with hooded sprayers to control labeled weeds between the rows of the below listed emerged crops. This treatment is for crops grown in rows, and includes crops grown in rows where mulch or plastic barriers are used as a weed control tool in the drill or plant line.

**Hooded sprayers must be designed, adjusted and operated in such a manner to totally enclose the spray pattern and to prevent any spray deposition to green stem tissue, foliage, blooms or fruit of the crop.**

Sprayers shall not be operated at more than five (5) miles per hour in order to minimize vertical movement of the sprayer during application, including the bouncing or raising of the equipment. Use extreme care in applying to fields where the soil surface is uneven, has deep furrows, drains or other contours that would disturb the adjustment and positioning of the spray equipment and/or the spray pattern. Applications must not be made when wind conditions may disturb the spray patterns and result in spray deposition to sensitive plants or plant parts.

For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. **Coverage is essential for good control.**

#### Crops Labeled for Use with Hooded Sprayers:

Hooded Spray application can be used for all crops listed on this AIM EC HERBICIDE label.

#### Note:

**Crop injury will occur when spray is allowed to come in contact with the green stem tissue, leaves, blooms or fruit of the crop.**

**See listing for individual commodities contained within the respective Crop groups:**

**Vegetable, Root and Tuber (Group 1) including:** Arracacha, Arrowroot, Chinese artichoke, Jerusalem artichoke, Garden Beet, Sugar beet, Edible Burdock, Edible Canna, Carrot; Bitter and Sweet Cassava, Celeriac, Chayote (root), Turnip-rooted Chervil, Chicory, Chufa, Dasheen (taro), Ginger, Ginseng, Horseradish, Leren, Turnip-rooted Parsley, Parsnip, Potato, Radish, Oriental (daikon) Radish, Rutabaga, Salsify, Black Salsify, Spanish Salsify, Skirret, Sweet Potato, Tanier, Turmeric, Turnip, Yam bean; True Yam

**Vegetable, leaves of root and tuber (Group 2) including:** Garden Beet, Sugar Beet, Edible Burdock, Carrot, Bitter and Sweet Cassava, Celeriac, Chervil, Turnip-rooted, Chicory, Dasheen (taro), Parsnip, Radish, Oriental (daikon) Radish, Rutabaga, Black Salsify, Sweet Potato, Tanier, Turnip, True Yam

**Vegetable, bulb (Group 3-07) including:** Fresh Leaves Chive, Chinese Fresh Leaves Chive, Bulb Daylily, Elegans Hosta; Bulb Fritillaria, Leaves Fritillaria, Bulb Garlic, Great-headed Garlic, Serpent Bulb Garlic, Kurrat; Lady's Leek, Leek, Wild Leek, Bulb Lily, Beltsville Bunching Onion, Bulb Onion, Chinese Bulb Onion, Fresh Onion, Green Onion, Macrostem onion, Pearl onion, Potato Bulb Onion, Tree Tops Onion, Welsh Tops Onion, Bulb Shallot, Fresh Leaves shallot, and cultivars, varieties, and/or hybrids of these

**Vegetable, leafy except brassica (Group 4) including:** Amaranth (Chinese Spinach), Arugula (Roquette), Cardoon, Celery, Chinese Celery, Celtuce, Chervil, Edible-Leaved Chrysanthemum, Garland Chrysanthemum, Corn Salad, Cress, Garden, Upland Cress, Dandelion, Dock (Sorrel), Endive (Escarole), Florence Fennel, Head And Leaf Lettuce, Orach, Parsley, Garden Purslane, Winter Purslane, Radicchio (Red Chicory), Rhubarb, Spinach, New Zealand Spinach, Vine Spinach, Swiss Chard

**Vegetable, brassica (Group 5) including:** Broccoli; Chinese Broccoli, (gai lon), Broccoli Raab (rapini), Brussels Sprouts, Cabbage, Chinese Cabbage, (bok choy); Chinese Cabbage (napa), Chinese Mustard Cabbage (gai choy), Cauliflower, Cavalo Broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens

**Vegetable, legume, except soybean (succulent or dried) (Group 6) including:** Bean (Lupinus spp.) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (Phaseolus spp.) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); bean (Vigna spp.) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); broad bean (fava); chickpea (garbanzo); guar; jackbean; lablab bean (hyacinth bean); lentil; pea (Pisum spp.) (includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea); pigeon pea; soybean (immature seed); sword bean

**Vegetable, foliage of legume (Group 7) including:** Plant parts of any legume vegetable included in the legume vegetables group that will be used as animal feed

**Vegetable, fruiting (Group 8-10) including:** African eggplant, Bush Tomato, Bell Pepper, Cocona, Currant Tomato, Eggplant, Garden Huckleberry, Goji Berry, Groundcherry, Martynia, Naranjilla, Okra, Pea Eggplant, Pepino, Non-Bell Pepper, Roselle, Scarlet Eggplant, Sunberry, Tomatillo, Tomato, Tree Tomato, and cultivars, varieties, and/or hybrids of these

**Vegetable, cucurbit (Group 9) including:** Chayote (fruit), Chinese Waxgourd (Chinese Preserving Melon), Citron Melon, Cucumber, Gherkin, Edible Gourd (includes Hyotan, Cucuzza, Hechima, Chinese Okra), Momordica spp. (includes Balsam Apple, Balsam Pear, Bittermelon, Chinese Cucumber), Muskmelon (includes Cantaloupe), Pumpkin, Summer Squash, Winter Squash (includes Butternut Squash, Calabaza, Hubbard Squash, Acorn Squash, Spaghetti Squash), Watermelon

**Citrus Fruit (Group 10-10) including:** Australian Desert Lime, Australian Finger-Lime, Australian Round Lime, Brown River Finger Lime, Calamondin, Citron, Citrus hybrids, Grapefruit, Japanese Summer Grapefruit, Kumquat, Lemon, Lime, Mediterranean Mandarin, Mount White Lime, New Guinea Wild Lime, Sour Orange; Sweet Orange, Pummelo, Russell River Lime, Satsuma Mandarin, Sweet Lime, Tachibana Orange, Tahiti Lime, Tangelo, Tangerine (mandarin), Tangor, Trifoliate Orange; Uniq Fruit, and cultivars, varieties, and/or hybrids of these

**Pome Fruit (Group 11-10) including:** Apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these

**Stone Fruit (Group 12-12) including:** Apricot, Japanese Apricot, Capulin, Black Cherry, Nanking Cherry, Sweet Cherry, Tart Cherry, Chinese Jujube, Nectarine, Peach, Plum, American Plum, Beach Plum, Canada Plum, Cherry Plum, Chickasaw Plum, Damson Plum, Japanese Plum, Klamath Plum, Prune Plum, Plumcot, Sloe and cultivars, varieties, and/or hybrids of these

**Caneberry (subgroup 13-07A) including:** Blackberry, Loganberry, Black and Red Raspberry, Wild Raspberry, and cultivars, varieties, and/or hybrids of these

**Bushberry (subgroup 13-07B) including:** Aronia Berry, Highbush Blueberry, Lowbush Blueberry, Buffalo Currant, Chilean Guava, Highbush Cranberry, Black Currant, Red Currant, Elderberry, European Barberry, Gooseberry, Edible Honeysuckle, Huckleberry, Jostaberry, Juneberry (Saskatoon Berry), Lingonberry, Native Currant, Salal, Sea Buckthorn and cultivars, varieties, and/or hybrids of these

**Fruit, small vine climbing – except fuzzy kiwifruit (subgroup 13- 07F) including:** Amur river grape; gooseberry; grape; hardy kiwifruit; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these

**Berry, low growing (subgroup 13-07G) including:** Bearberry, Bilberry, Lowbush Blueberry, Cloudberry, Cranberry, Lingonberry, Muntries, Partridgeberry, Strawberry, and cultivars, varieties, and/or hybrids of these.

**Tree Nuts (Group 14-12) including:** African Nut-Tree, Almond, Beechnut, Brazil Nut; Brazilian Pine, Bunya, Bur Oak, Butternut, Cajou Nut, Candlenut, Cashew, Chestnut, Chinquapin, Coconut, Coquito nut, Dika Nut, Ginkgo, Guiana Chestnut, Hazelnut (filbert); Heartnut, Hickory Nut, Japanese Horse-Chestnut, Macadamia Nut, Mongongo Nut, Monkey-Pot, Monkey Puzzle Nut, Okari Nut, Pachira Nut, Peach Palm Nut, Pecan, Pequi, Pili Nut, Pine Nut, Pistachio, Sapucaia Nut, Tropical Almond, Black Walnut, English Walnut, Yellowhorn and cultivars, varieties, and/or hybrids of these

**Cereal Grains (Group 15) including:** Barley, Buckwheat, Corn, Millet (Pearl and proso), Oats, Popcorn, Rice, Rye, Sorghum (milo), Teosinte, Triticale, Wheat, and Wild Rice

**Forage, Fodder and Straw of Cereal Grains (Group 16) including:** Forage, fodder and straw of all commodities included in the cereal grains (Group 15)

**Grasses (Group 17) including:** Any grass, Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage

**Non-grass Animal Feed (Group 18) including:** Alfalfa, Velvet Bean, Clover (Trifolium spp., Melilotus spp.), Kudzu, Lespedeza, Lupin, Sainfoin, Trefoil, Vetch, Crown Vetch, Milk Vetch.

**Herbs and Spices (Group 19) including:** Allspice, Angelica, Anise (seed), Star Anise, Annatto (seed), Balm (Lemon Balm), Basil (Fresh and Dried), Borage, Burnet, Camomile, Caper Buds, Caraway, Black Caraway, Cardamom, Cassia Bark, Cassia Buds, Catnip, Celery Seed, Chervil (dried), Chive, Chinese Chive, Cinnamon, Clary, Clove Buds, Coriander Leaf (Cilantro or Chinese Parsley), Coriander Seed (Cilantro), Costmary, Culantro (Leaf), Culantro (Seed), Cumin, Curry (Leaf), Dill (Dillweed), Dill (Seed), Fennel (Common), Florence Fennel (seed), Fenugreek, Grains of Paradise, Horehound, Hyssop, Juniper Berry, Lavender, Lemongrass, Lovage (leaf), Lovage (seed), Mace, Marigold,

Marjoram (includes Sweet or Annual Marjoram, Wild Marjoram or Oregano, and Pot Marjoram), Mustard (Seed), Nasturtium, Nutmeg, Parsley (Dried), Pennyroyal, Black Pepper, White Pepper, Poppy (Seed), Rosemary, Rue, Saffron, Sage; Summer and Winter Savory, Sweet Bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.

**Oil Seeds (Subgroups 20A & 20B, except Cottonseed) including:** Borage, Crambe, Cuphea, Echium, Flax Seed, Gold of Pleasure, Hare's Ear Mustard, Lesquerella, Lunaria, Meadowfoam, Milkweed, Mustard Seed, Oil Radish, Poppy Seed, Rapeseed, Sesame, Sweet Rocket, Calendula, Castor Oil Plant, Chinese Tallowtree, Euphorbia, Evening Primrose, Jojoba, Niger Seed, Rose Hip, Safflower, Stokes Aster, Sunflower, Tallowwood, Tea Oil Plant, Vernonia and cultivars, varieties, and/or hybrids of these.

**Tropical fruit:** including Acerola, Atemoya, Avocado, Biriba, Black Sapote, Canistel, Cherimoya, Custard apple, Feijoa, Guava, Jaboticaba, Llama, Longan, Lychee, Mamey Sapote, Mango, Papaya, Passionfruit, Pawpaw, Pulasan, Rambutan, Sapodilla, Soursop Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu, Aloe vera, Cactus.

For additional information regarding crops within a group, refer to the Website:

<https://www.gpo.gov/fdsys/pkg/CFR-2010-title40-vol23/pdf/CFR-2010-title40-vol23-sec180-41.pdf>

# APPLICATION INSTRUCTIONS

## Alfalfa and Clover (Established Stands Only) Crop Group 18

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Postemergence Weed Control (Dormant, In-crop, and Stubble)</b>	<b>DO NOT</b> apply within 21 days of harvest for stands grown for forage and hay.	Refer to table 3	0.5-2.5 fl oz/A (0.008 – 0.04 lb ai/A)	<b>DO NOT</b> apply more than 2.5 fl oz/A (0.04 lb ai/A) per year for postemergence weed control applications in nongrass animal feeds.
<b>Harvest Aid</b>	<b>DO NOT</b> apply within 21 days of harvest for stands grown for forage and hay.  <b>DO NOT</b> apply within 3 days of harvest for stands grown for seed.	Refer to table 3	2.0 to 3.8 fl oz/A (0.031 – 0.06 lb ai/A)	After an application of this product to crop group 18 (nongrass animal feed crops), you may only rotate the field to a carfentrazone-ethyl registered crop.

### DIRECTIONS FOR USE

#### Postemergence Weed Control Treatment

##### Dormant Period (Fall or Winter Application Postemergence on Weeds)

AIM EC HERBICIDE may be applied on dormant crop stubble alone or in combination with other registered herbicides for the post emergence control of weeds in established nongrass animal feed stands during the dormant period (between growing periods). To control insect pests, AIM EC HERBICIDE may be tank mixed with insecticides, including Mustang Maxx.

##### Between Cutting In-Crop Application (Spring/Summer Applications Postemergence on Weeds)

AIM EC HERBICIDE may be applied alone or in combination with other registered herbicides between cuttings (in-crop) for the post emergence control of weeds in established crop stands. In-crop applications must be made as soon as possible after removal of the previous hay crop and prior to significant regrowth on stems and crowns. Applications may be made from hay removal up to 6 inches of new growth. To control insect pests, AIM EC HERBICIDE may be tank mixed with insecticides, including Mustang Maxx.

#### AIM EC HERBICIDE Use Rates

For optimum results, weeds must be treated when small. Applications shall be made in spray volumes sufficient to provide complete coverage of foliage.

Use a minimum of 10 gallons of finished spray per acre for ground application equipment, and a minimum of 3 gallons per acre of finished spray for aerial equipment. For optimum results, apply AIM EC HERBICIDE to weeds up to 4 inches tall and rosettes less than 3 inches across. Use a quality nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. For more active treatments, use a Crop Oil Concentrate (COC) at 0.5 to 1.0% v/v (one half to one gallon per 100 gallons). Some temporary leaf speckling and necrosis may occur on green alfalfa or clover tissue present with between cutting applications, which must be rapidly outgrown under good growing conditions. Adjuvant selection and high moisture environmental conditions will enhance this effect. A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed. Coverage is essential for satisfactory performance. Repeat application if necessary. **DO NOT** irrigate just prior to or just after application. Weed control under dry and hot conditions will be improved with COC or similar products.

#### Tank Mix

For tank mixture applications, refer to the use directions and restrictions of the mixture product.

AIM EC HERBICIDE may be tank mixed with other labeled herbicides to control weeds not listed on this label. Read and follow all manufacturers' label directions and label restrictions for the companion herbicide. When tank mixing AIM EC HERBICIDE with other products, be sure AIM EC HERBICIDE is mixed in the spray tank water first.

#### Harvest Aid Treatment

Apply AIM EC HERBICIDE to crops grown for forage, hay or seed alone or as a tank mixture with other harvest aids. Applications shall be made when the crop is mature, or according to Extension Service guidelines in the use area. Apply AIM EC HERBICIDE at 2.0 to 3.8 fl oz/A (0.031 to 0.06 lb ai/A), but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE AIM EC HERBICIDE USE RATE CHART and the PREHARVEST INTERVAL charts for additional application information. If treatments of AIM EC HERBICIDE have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment.

Applications must be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed. Coverage is essential for satisfactory performance. Repeat application if necessary.

#### Note

If applied as a tank mixture, refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

**ASPARAGUS**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Postemerge Weed Control</b>	<b>DO NOT</b> apply within 5 days of harvest.	Refer to table 3	Apply one to two applications of AIM EC HERBICIDE at 0.5 to 1.92 fl oz/A (0.008 to 0.03 lb ai/A)  Use higher rates when Asparagus tissues and weeds are under stress or are larger.	<b>DO NOT</b> apply more than 3.8 fl oz/A (0.06 lb ai/A) per year.  <b>DO NOT</b> make applications less than 20 days apart.

**DIRECTIONS FOR USE**

Apply AIM EC HERBICIDE as a broadcast application after harvest of Asparagus spears for control of broadleaf weeds and new existing Asparagus tissues.

**Coverage is essential for good control.**

**Adjuvant Requirements**

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed. Repeat application if necessary.

For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section.

**BUSHBERRY Subgroup 13-07B**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
Postemergence Weed Control	Can be applied up to harvest	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> apply more than 2 fl oz/A (0.031 lb ai/A) per application during the dormant period. <b>DO NOT</b> apply more than 6.1 fl oz/A (0.095 lb ai/A) per year.

**DIRECTIONS FOR USE**

AIM EC HERBICIDE applications will control susceptible emerged broadleaf weeds. Repeat applications may be necessary for weeds that emerge after an AIM EC HERBICIDE treatment.

**Equipment and Application**

Apply only by ground equipment including boom sprayers, shielded or hooded sprayers, hand-held or high-volume wands or orchard guns. Use a minimum of 20 gallons finished spray solution per broadcast acre.

**Dormant Applications**

Apply AIM EC HERBICIDE as a broadcast application to the base of the trunk to control emerged and actively growing weeds during the dormant stage of the crop.

**Post-directed Applications for Broadleaf Weed Control**

Apply AIM EC HERBICIDE as a directed spray avoiding contact with the berry plant but directed at actively growing weeds. AIM EC HERBICIDE is a contact herbicide and coverage is essential for good weed control. **DO NOT** allow AIM EC HERBICIDE spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage.

Newly planted bush berries must only be treated with shielded sprayers or hooded sprayers.

**AIM EC HERBICIDE Use Rates**

Apply up to 2 fl oz/A (0.031 lb ai/A) AIM EC HERBICIDE per broadcast acre. For best control, apply to seedling weeds in the 2 to 3-leaf stage. Use higher labeled rates of AIM EC HERBICIDE for larger weeds up to 6 leaves. Weeds greater than 6 leaves may be only partially controlled. See Table 3 for AIM EC HERBICIDE use rates and weeds controlled.

**Adjuvant Requirements**

A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

**Tank Mix**

AIM EC HERBICIDE may be mixed with other registered herbicides for broader spectrum weed control. When tank mixing with fertilizer solutions, be sure to prepare an AIM EC HERBICIDE premixture of AIM EC HERBICIDE and clean water.

See Mixing and Loading Instructions under the PRODUCT INFORMATION section of this label for specific mixing instructions. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner.

**Precautions**

Extreme caution must be taken during applications when desirable fruit, foliage and/or blooms are present in order to avoid spotting or necrosis. **DO NOT** allow AIM EC HERBICIDE spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage.

For seedling or newly transplanted bushes, **DO NOT** allow spray to contact green bark of trunk area. Use shielded sprayers only.

**Band Treatment Application**

For band treatment, apply the broadcast equivalent rate and volume per acre.

To determine these:

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$$

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Volume Per Acre} = \text{Band Volume}$$

CANEBERRY Subgroup 13-07A

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
Postemergence Weed Control	DO NOT apply within 15 days of harvest.	Refer to table 3	Apply 6.4 fl oz/A AIM EC HERBICIDE (0.1 lb ai/A) per broadcast acre as a directed spray when weeds and primocanes are approximately 6 inches tall. Apply up to 2 fl oz/A (0.031 lb ai/A) AIM EC HERBICIDE per broadcast acre. For best control, apply to actively growing weeds up to 4 inches tall or rosettes less than 3 inches across.	DO NOT apply more than 25.6 fl oz/A (0.4 lb ai/A) per year. DO NOT make applications less than 14 days apart.

**DIRECTIONS FOR USE**

**Equipment and Application**

Apply only by ground equipment including boom sprayers, shielded or hooded sprayers, hand-held or high-volume wands or orchard guns. **DO NOT** allow AIM EC HERBICIDE spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage. **DO NOT** apply when conditions favor drift or when wind is above 10 mph.

**Post-Directed Application for Primocane and Weed Control**

AIM EC HERBICIDE is a contact herbicide for directed application for the control of primocanes and weeds.

Use a minimum of 20 gallons finished spray per broadcast acre at intervals of 14 to 21 days. Direct spray to the bottom 18 inches of the canes and to the soil 24 inches from each side of the plant row. Refer to weed control list in Table 3 for appropriate weed control information.

**Adjuvant Requirements**

An adjuvant is required. See Adjuvant Requirements below under weed control.

**Post-Directed Application for Weed Control**

Apply AIM EC HERBICIDE to actively growing weeds. AIM EC HERBICIDE is a contact herbicide and coverage is essential for good weed control. Use a minimum of 20 gallons finished spray solution per acre.

**Band Treatment Application**

For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

**Band Treatment Application**

For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$$

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Volume Per Acre} = \text{Band Volume}$$

**Coverage is essential for good control.**

**Adjuvant Requirements**

A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

AIM EC HERBICIDE may be mixed with other herbicides registered in caneberrys for broader spectrum weed control.

AIM EC HERBICIDE must be the first product added to the spray tank water. See Mixing and Loading Instructions under the PRODUCT INFORMATION section of this label for specific mixing instructions. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner.

**Precautions**

Extreme caution must be taken during applications when desirable fruit, foliage and/or blooms are present in order to avoid spotting or necrosis. **DO NOT** allow AIM EC HERBICIDE spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage.

Newly planted caneberrys must only be treated with shielded sprayers or hooded sprayers.

**Restrictions**

**DO NOT** apply more than 25.6 fl oz/A (0.4 lb ai/A) per year.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 15 days of harvest.

**Corn (Field, Seed, Silage, Popcorn, Sweet Corn – Processing and Fresh Market)**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
Preplant Burndown	NA	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per year including all preplant, in-crop, and harvest aid applications.
Postemergence (Broadcast)	14 leaf collar	Refer to table 3	Up to 1.0 fl oz/A (0.016 lb ai/A)	
Postemergence (Hooded Sprayer and Directed Applications)	14 leaf collar	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	
Harvest Aid	<b>DO NOT</b> apply within 3 days of harvest	Refer to table 3	1.0 to 2.0 fl oz/A (0.016 to 0.031 lb ai/A)	

**DIRECTIONS FOR USE**

**Postemergence Weed Control Treatment**

Apply AIM EC HERBICIDE alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to corn in all tillage systems from prior to planting up to 14-leaf collar growth stage. When applying AIM EC HERBICIDE to corn greater than V8 stage, utilize drop nozzles aligned between the rows with directed application to reduce contact with the corn foliage and improve contact with the weeds. For optimum performance, make application to actively growing weeds up to 4 inches high and rosettes less than 3 inches across. Coverage is essential for good control.

**Broadcast Applications:**

Use AIM EC HERBICIDE at 0.5 to 1.0 fl oz/A as a broadcast application using a minimum of 10 gallons per acre of spray volume by ground or 3 gallons per acre by air. Broadcast applications may be applied through V8 stage corn.

**Tank Mix**

AIM EC HERBICIDE may be tankmixed with other corn herbicides to control weeds not listed on this label. Read and follow all manufacturers' label directions for the companion herbicides. When tank mixing AIM EC HERBICIDE with other labeled corn herbicides, use adjuvants as directed by the tank mix partner's label.

**AIM EC HERBICIDE plus Atrazine**

AIM EC HERBICIDE may be tank mixed at a rate of 0.5 fl oz/A (0.008 lb ai/A) with Atrazine 4L (16 fl oz/A) or Atrazine 90DF (0.6 -1.6 lb/A) to control the following weeds:

**When used as directed, AIM EC HERBICIDE + atrazine will provide control of listed weeds up to 4 inches tall.**

Amaranth, Palmer (not triazine resistant)	Copperleaf, hophornbeam	Mallow, Venice	Purslane, common
Amaranth, spiny	Croton, wooly	Morningglory spp.	Sesbania, hemp
Anoda, spurred	Devilsclaw	Nightshade, Eastern black	Thistle, Russian
Buckwheat, wild	Eveningprimrose, cutleaf	Nightshade, hairy	Velvetleaf
Buffalobur	Jimsonweed	Pigweed, redroot	Waterhemp, common
Carpetweed	Kochia*	Pigweed, smooth	Waterhemp, tall
Cocklebur	Lambsquarters, common	Potato, volunteer	Sesbania, hemp

\* Kochia control up to 2 inches tall with AIM EC HERBICIDE + Atrazine + COC only. Refer to the Atrazine labels for additional weed listings and for higher use rates.

**AIM EC HERBICIDE plus Dicamba**

AIM EC HERBICIDE at 0.5 fl oz/A (0.008 lb ai/A) plus 0.25% v/v nonionic surfactant (2 pints per 100 gallons) can be tank mixed with dicamba herbicides (8 -16 fluid ounces per acre) for control of broadleaf weeds including the following:

<b>When used as directed, AIM EC HERBICIDE + dicamba will provide control of listed weeds up to 4 inches tall.</b>			
Buckwheat, wild	Morningglory, spp.	Potato, volunteer	Thistle, Russian
Cocklebur, common	Nightshade, black	Ragweed, common	Velvetleaf
Jimsonweed	Pigweed, redroot	Ragweed, giant	Waterhemp, common
Kochia	Pigweed, smooth	Smartweed, PA (seedling)	Waterhemp, tall
Lambsquarters	Pigweed, triazine resistant	Sunflower, common	

Refer to the dicamba labels for additional weed listings and for higher use rates. Refer to the Tank Mixture Section for information on potential leaf injury.

**AIM EC HERBICIDE Plus Atrazine Plus Dicamba or 2,4-D**

For the control of additional or certain larger weeds up to 6 inches tall, Atrazine may be added to the tank mixtures of AIM EC HERBICIDE plus dicamba or AIM EC HERBICIDE plus 2,4-D (amine).

Add 2,4-D (amine) to the tank mix at 0.125 to 0.25 lb ai/A or dicamba at 3 to 8 fl oz/A. Higher rates of atrazine and dicamba herbicides are allowed, but **DO NOT** exceed the specific label use rates allowed by these labels. Add a 0.25% v/v nonionic surfactant (2 pints per 100 gallons) to the tank mixture. Under very dry soil moisture conditions, the use of crop oil concentrate at 1% v/v (1 gallon per 100 gallon spray solution) may improve weed control. The use of crop oil concentrate may increase leaf speckling. Refer to the Tank Mixture section for information on potential leaf injury.

**For control of the following weeds up to 6 inches in height, or as specified, add dicamba at 3 to 8 ounces per acre to AIM EC HERBICIDE tank mixes with atrazine or to AIM EC HERBICIDE tank mixes with other products that allow the use of dicamba on their labels.**

Amaranth, Palmer (up to 4 inches)	Nightshade, Eastern black	Smartweeds, annual (seedling)
Amaranth, Spiny (up to 4 inches)	Nightshade, hairy	Sunflower, common (up to 4 inches tall)
Cocklebur, common	Pigweed, redroot	Velvetleaf (up to 24 inches)
Kochia (up to 4 inches tall)	Pigweed, smooth	Waterhemp, common
Lambsquarters, common	Ragweed, common	Waterhemp, tall
Morningglory spp.	Ragweed, giant (up to 4 inches tall)	

**Adjuvant Requirements:**

Use a non-ionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution). Under dry conditions, the use of a crop oil concentrate (COC) at 1.0% v/v may improve weed control. The use of crop oil concentrate can increase leaf speckling and crop response on treated corn leaves.

**Directed Spray Applications:**

Apply AIM EC HERBICIDE with drop nozzles between the rows to the target weeds and away from the whorl of the corn plant. Directed spray applications must be used when corn is V8 to V14 stage. Apply AIM EC HERBICIDE up to 2.0 fl oz/A (0.031 lb ai/A). Be aware that weeds growing in and under the dense canopies may not receive adequate spray coverage and may require the use of higher spray volumes for acceptable control. Use appropriate rates of adjuvants including non-ionic surfactant (NIS), crop oil concentrate (COC), or methylated seed oil (MSO).

**Hooded Sprayer Applications:**

Apply AIM EC HERBICIDE with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

**Harvest Aid:**

Apply 1.0 to 2.0 fl oz/A AIM EC HERBICIDE, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE AIM EC HERBICIDE USE RATE and the PREHARVEST INTERVAL Table (Table 2) for additional application information. If treatments of AIM EC HERBICIDE have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment.

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use methylated seed oil, or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil is allowed.

**Coverage is essential for satisfactory performance.**

**Seed Corn Production:**

For seed production fields, apply AIM EC HERBICIDE using drop nozzles or other equipment to make a directed spray treatment. Avoid directing spray solution into the whorl.

Seed corn inbred lines have generally shown good tolerance to AIM EC HERBICIDE. However, all inbred lines have not been tested. Broadcast applications may result in spray being concentrated into the whorl of the plant that will increase leaf response. To minimize application into the whorl of the plants, drop nozzles or other type directed sprayers must be used to direct the spray to the targeted weeds.

**Sweet Corn Precaution:**

When applying AIM EC HERBICIDE to sweet corn; broadcast applications may result in spray being concentrated into the whorl of the plant that will increase leaf response. To minimize application into the whorl of the plants, drop nozzles or other type directed sprayers must be used to direct the spray to the targeted weeds.

Use only NIS as the spray adjuvant in sweet corn applications.

**Application Precautions:**

Leaf speckling can occur when AIM EC HERBICIDE is used with certain crop protection products and adjuvants. Refer to the Tank Mixtures and Adjuvants requirements sections under PRODUCT INFORMATION. Bromoxynil mixtures and bentazon mixtures may cause significant crop response when in contact with crop foliage.

**Crop Response**

The application of AIM EC HERBICIDE to corn may result in temporary crop response including speckling or necrosis of the leaves. Grain yields will not be affected. **DO NOT** make applications when air temperatures are abnormally cool or humidity is high or if the corn foliage is wet from dew, rainfall or irrigation. Users must be aware of these inherent risks and accept these risks prior to application of AIM EC HERBICIDE.

For additional information regarding potential crop response, refer to the PRODUCT INFORMATION section of the AIM EC HERBICIDE label.

**Coverage is essential for satisfactory performance.**

**Cotton:**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Removal of Failed Cotton Stands</b>	<b>DO NOT</b> apply within 7 days of harvest.	Failed Cotton (up to 3 leaf cotton)	1.0 to 1.6 fl oz/A (0.016 – 0.025 lb ai/A)	<b>DO NOT</b> apply when conditions favor drift or when wind is above 10mph. <b>DO NOT</b> apply more than 7.9 fl oz/A (0.123 lb ai/A) total for preplant, in-crop weed control and harvest aid. <b>DO NOT</b> apply more than 3.2 fl oz/A (0.05 lb ai/A) total for managed maturity and/or as a harvest aid.
<b>Pre Plant Burndown</b>	<b>DO NOT</b> apply within 7 days of harvest.	Refer to table 3	Up to 1.6 fl oz/A (0.025 lb ai/A)	
<b>Hooded Sprayer</b>	<b>DO NOT</b> apply within 7 days of harvest.	Refer to table 3	Up to 1.6 fl oz/A (0.025 lb ai/A)	
<b>Post-directed and Lay-by</b>	<b>DO NOT</b> apply within 7 days of harvest.	Refer to table 3	Up to 1.6 fl oz/A (0.025 lb ai/A)	
<b>Managed Maturity</b>	<b>DO NOT</b> apply within 7 days of harvest.	Manage unproductive terminal growth in cotton.	0.25 to 0.5 fl oz/A (0.004 – 0.008 lb ai/A)	
<b>Defoliation/ Harvest Aid</b>	<b>DO NOT</b> apply within 7 days of harvest.	Defoliate and desiccate cotton and troublesome weeds	Up to 1.6 fl oz/A (0.025 lb ai/A)	

**DIRECTIONS FOR USE**

**Removal of Failed Cotton Stands**

Apply 1.0 to 1.6 fl oz/A AIM EC HERBICIDE (0.016 to 0.025 lb ai/A) broadcast as a foliar spray over the top of the remaining cotton plants with sufficient spray volume to provide adequate coverage of the cotton plant, particularly the terminal area. Use higher rates on larger failed cotton. For best results **DO NOT** exceed 3 leaf cotton. **Coverage is essential for good control.**

**PRE PLANT BURNDOWN**

See instructions under the Pre Plant Burndown section of this label.

**Hooded Sprayer Applications**

Apply AIM EC HERBICIDE with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

**Post-directed and Lay-by Applications**

AIM EC HERBICIDE is a contact herbicide for postemergence directed sprayer or hooded/shielded sprayer applications for the control of broadleaf weeds in cotton. Apply AIM EC HERBICIDE alone or as a tank mixture with other herbicides to emerged and actively growing weeds. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. Applications of AIM EC HERBICIDE or AIM EC HERBICIDE tank mixes must be made with directed sprayers or hooded sprayers to prevent contact of spray solution with the cotton plant. **DO NOT** allow spray solution to contact cotton foliage, green stem tissue, or blooms. Directed spray equipment must position nozzles a minimum 3 to 4 inches above the soil, with nozzles directed beneath the crop canopy. AIM EC HERBICIDE or AIM EC HERBICIDE tank mix applications shall be made to cotton that is a minimum of 6 inches in height. Applications to cotton at 5 to 6 nodes or less must be made with hooded or shielded sprayer equipment to completely avoid contact with cotton plants. Apply lay-by applications of AIM EC HERBICIDE or AIM EC HERBICIDE tank mixtures at later growth stages of cotton when cotton plants have achieved a height of 12 inches or more with sufficient bark development and height differential between crop bottom leaves and the soil. Spray solution shall be directed at the base of cotton plants for minimal contact with green stem tissue or foliage while maintaining maximum contact with broadleaf weeds that are at appropriate treatment size.

For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. Coverage is essential for good control.

**Adjuvant Requirements**

A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

**AIM EC HERBICIDE Use Rates and Weeds Controlled**

Apply up to 1.6 fl oz/A (0.025 lb ai/A) AIM EC HERBICIDE as a post-directed treatment using a directed sprayer a hooded sprayer or lay-by sprayer delivering a minimum finished spray volume of 10 gallons per acre. **DO NOT** apply more than 3.2 fl oz/A (0.05 lb ai/A) AIM EC HERBICIDE per year by post-directed and lay-by applications. Refer to weed control list in Table 3 for appropriate weed control information.

For control of additional broadleaf weeds and grasses, AIM EC HERBICIDE may be tank mixed with other herbicides registered for cotton post-directed and/or lay-by applications. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

**Managed Maturity Application for Cotton**

Apply AIM EC HERBICIDE as an aid to remove undesirable top growth and reduce unproductive terminal growth. Use alone or as a tank mixture with other cotton insecticides and herbicides. Read all product labels and follow all directions and precautions when tank mixing with this product.

**Timing**

Apply AIM EC HERBICIDE when cotton is actively growing and the plants have 1% to 20% open bolls; with applications at 15% open bolls being optimum. When using the Cotman monitoring program, apply AIM EC HERBICIDE at NAWF5, plus 450 – 650 heat units. Avoid Managed Maturity treatments to fields, or areas of fields, that are stressed.

**AIM EC HERBICIDE Use Rates – Manage Maturity**

Apply AIM EC HERBICIDE as a broadcast spray at 0.25 fl oz/A (0.004 lb ai/A) to 0.5 fl oz/A (0.008 lb ai/A), targeting 0.375 fl oz/A (0.006 lb ai/A) in spray volume adequate to obtain upper canopy coverage of the plant foliage. In situations of extremely lush growth, apply up to 0.5 fl oz/A (0.008 lb ai/A). Make applications using a minimum of 10 gallons of finished spray per acre for ground application and a minimum of 5 gallons per acre by air. Good upper canopy coverage is essential for optimum performance.

Use a quality crop oil concentrate (COC) at the specified rate of 1% v/v.

**Defoliation / Harvest Aid Application**

Apply AIM EC HERBICIDE as a harvest aid to defoliate and desiccate cotton and troublesome weeds that may be present at harvest. Apply AIM EC HERBICIDE alone or as a tank mixture with other cotton harvest aids.

Use a quality spray adjuvant e.g. nonionic surfactant (NIS) or crop oil concentrate (COC) at the specified rates. NIS is the required adjuvant during warmer periods with COC being the better choice for applications during cooler periods. Make application when 60 to 70 percent of the bolls are open, or according to the State Agricultural Extension Service guidelines in the use area.

Apply up to 1.6 fl oz/A AIM EC HERBICIDE (up to 0.025 lb ai/A) in spray volume sufficient to provide complete coverage of cotton foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. **Coverage is essential for good defoliation.** Repeat application if necessary to remove remaining foliage or control regrowth. **DO NOT** apply more than 3.2 fl oz/A (0.05 lb ai/A) total as a harvest aid. Dense cotton canopy, large plant size, and environmental conditions not conducive to complete plant coverage may reduce initial application performance and increase the need for a second application.

Apply AIM EC HERBICIDE alone, as a tank mix, or as a sequential application alone or tank mixed with other registered cotton harvest aid products.

Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

**Dried shelled beans, peas Crop Subgroup 6-C: and flax (except soybean) and vegetable foliage of legume Group 7.**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Preplant Burndown</b>	NA	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> apply more than 6.1 fl oz/A (0.095 lb ai/A) per year.
<b>Harvest Aid Applications</b>	Can be applied up to 0 days before harvest.	Refer to table 3	1.0 to 4.0 fl oz/A (0.016 to 0.063 lb ai/A)	<b>DO NOT</b> apply more than 4.0 fl oz/A (0.063 lb ai/A) and for up to 12 months following harvest-aid application, the subsequent planted crop may only be a registered crop.

**DIRECTIONS FOR USE****Preplant Burndown:**

Refer to the preplant burn down section of this label.

**Harvest Aid Treatment:**

Apply AIM EC HERBICIDE as a harvest aid to dry beans and dry peas at maturity when 80 to 90% of seed pods are yellow or buck skin in color and only 30% of green leaves remain on the plant. Apply to flax when 75% of the bolls have turned brown. Thorough coverage is essential for harvest aid and multiple applications may be needed. For optimum performance use 15 to 30 gallons per acre finished sprayed with a methylated seed oil (MSO) type adjuvant to ensure thorough coverage and retention for harvest aid.

**AIM EC HERBICIDE Use rates:**

Apply AIM EC HERBICIDE alone or as a tank mixture with other harvest aids. Apply AIM EC HERBICIDE at 1.0 to 6.1 fl oz/A (0.016 to 0.095 lb ai/A) , but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE AIM EC HERBICIDE USE RATE CHART and the PREHARVEST INTERVAL charts for additional application information.

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A methylated seed oil (MSO) or crop oil concentrate (COC) is required at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). The addition of a high quality sprayable liquid nitrogen fertilizer at 2 to 4% v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil may enhance performance. If spraying dry beans before full maturity and pods are not all mature and turning color, a repeat application may be necessary.

**Note**

If applied as a tank mixture, refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

## Fallow Systems

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
Postemergence Weed Control	NA	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	For crop planting information following fallow treatments, refer to the Preplant Burndown section of this label for planting interval instructions.

### DIRECTIONS FOR USE

Apply AIM EC HERBICIDE by ground or air application alone or with other herbicides in the fallow period prior to planting or the emergence of any crop listed on this label to control or suppress weeds. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. Coverage is essential for good weed control.

AIM EC HERBICIDE may be utilized in Fallow Cropping Systems for chemical weed control to aid in moisture conservation between cropping periods.

### Adjuvant Requirements

A nonionic surfactant, crop oil concentrate or methylated seed oil is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed based crop oil concentrate (COC) at 1.0 to 2 % v/v (1.0 to 2.0 gallons per 100 gallons of spray solution) or a methylated seed oil (MSO). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons) or ammonium sulfate at 2 to 4 pounds per acre in addition to the selected NIS, MSO or COC is allowed.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide including glyphosate, glufosinate or paraquat. Refer to Table 3 for proper use rate for weed spectrum. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section.

### Tank Mix

For all products used in tank mixes, refer to the specific product labels for all restrictions on tank mixing and observe all label precautions, instructions and rotational cropping restrictions.

## Fruit, small vine climbing - except fuzzy kiwifruit (subgroup 13-07F)

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
Postemerge Weed Control	<b>DO NOT</b> apply within 3 days of harvest.	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per application (including preplant site preparation treatments). <b>DO NOT</b> apply more than 7.9 fl oz/A (0.124 lb ai/A) per year. <b>DO NOT</b> make applications less than 14 days apart.

### DIRECTIONS FOR USE

AIM EC HERBICIDE may be applied for postemergence weed control or for sucker control.

### Weed Control

Apply AIM EC HERBICIDE alone or as a tank mixture with other herbicides as a postemergence directed spray treatment or as a hooded spray treatment to control emerged and actively growing weeds. Apply AIM EC HERBICIDE to middles (between rows of plants) and in strips (in row of plants). Apply AIM EC HERBICIDE at any time during the growing period (see precautions). AIM EC HERBICIDE may be mixed with other herbicides that have pre-emergence or post-emergence activity. Any pre-emergence activity must rely on activity from other herbicides as directed on their labels. Herbicides including glyphosate may be tank mixed with AIM EC HERBICIDE for broader spectrum weed control. If AIM EC HERBICIDE is used in a tank mixture, observe the other product's label for restrictions, precautions and rotational cropping instructions.

### Sucker Management

AIM EC HERBICIDE is effective as an aid in the management of undesirable sucker growth from the base of vine trunks or root sprouts. Suckers and other undesirable growth must be treated when the tissue is young and not mature and/or hardened off. Care must be taken not to allow spray mist to contact desirable fruit or foliage or green stem tissue (see precautions). Application of AIM EC HERBICIDE with other sucker control herbicides is allowed.

### Hooded Sprayer Applications

Apply AIM EC HERBICIDE with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the **Hooded Sprayer Applications** section of this label for additional specific use directions.

### Equipment and Application

Coverage is essential for good control. Use a spray volume adequate to obtain thorough coverage with a minimum of 10 gallons of finished spray per acre. Apply only with ground equipment. Apply AIM EC HERBICIDE with hooded sprayers, boom equipment, shielded sprayers, hand-held and high-volume wands or orchard guns. Always add AIM EC HERBICIDE to the spray tank first. See "Mixing and Loading Instructions" under PRODUCT INFORMATION.

### Adjuvant Requirements

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a quality nonionic surfactant (NIS) containing at least 80% active at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate (COC) at 1% v/v (one gallon COC per 100 gallons), or a methylated seed oil (MSO). The use of a high quality sprayable liquid nitrogen fertilizer at 2 to 4% v/v or ammonium sulfate (AMS) used at 2 to 4 pounds per acre in addition to the NIS, or MSO or COC is allowed.

**Application Precautions: Extreme caution must be used during applications when desirable fruit or foliage is present in order to avoid fruit spotting or leaf necrosis.**

**DO NOT** allow AIM EC HERBICIDE spray mist to come in contact with desirable fruit, green stem tissue, foliage or blooms.

**DO NOT** use on seedling or newly transplanted vines **DO NOT** allow spray to contact green bark of trunk area.

**Fruit Tree, Tree Nut and Other Crops**

<b>CROP GROUP</b>	<b>PHI</b>	<b>Target Weeds</b>	<b>Rates</b>	<b>Restrictions</b>
<b>Citrus Fruits:</b> including Calamondin, Citrus Citron, Chironja, Tangelo, Tangor, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (sour), Orange (Sweet), Pummelo, Satsuma and Mandarin	<b>DO NOT</b> apply within 3 days of harvest	Refer to table 3	Apply up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> make applications with airblast sprayers. <b>DO NOT</b> make applications less than 14 days apart. <b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per application and 7.9 fl oz/A (0.123 lb ai/A) per year, including preplant site preparation.
<b>Pome Fruits:</b> including Apple, Crabapple, Loquat, Mayhaw, Pear, Pear (Oriental) and Quince	<b>DO NOT</b> apply within 3 days of harvest	Refer to table 3	Apply up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> make applications with airblast sprayers. <b>DO NOT</b> make applications less than 14 days apart. <b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per application and 7.9 fl oz/A (0.123 lb ai/A) per year, including preplant site preparation.
<b>Stone Fruits:</b> including Apricot, Cherry (Sweet), Cherry (Tart), Nectarine, Peach, Plum, Plum (Chickasaw), Plum (Damson), Plum (Japanese), Prune and Plumcot	<b>DO NOT</b> apply within 3 days of harvest	Refer to table 3	Apply up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> make applications with airblast sprayers. <b>DO NOT</b> make applications less than 14 days apart. <b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per application and 7.9 fl oz/A (0.123 lb ai/A) per year, including preplant site preparation.
<b>Tree Nuts:</b> including Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut (Bush Nut), Pecan, Pistachio and Walnut (Black and English)	<b>DO NOT</b> apply within 3 days of harvest	Refer to table 3	Apply up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> make applications with airblast sprayers. <b>DO NOT</b> make applications less than 14 days apart. <b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per application and 7.9 fl oz/A (0.123 lb ai/A) per year, including preplant site preparation.
<b>Tropical fruit:</b> including Papaya, Avocado, Black Sapote, Canistel, Mamey Sapote, Mango, Sapodilla, Star apple, Guava, Feijoa, Jaboticaba, Wax jambu, Starfruit, Passionfruit, Acerola, Lychee, Longan, Spanish lime, Rambutan, Pulasan, Sugar apple, Atemoya, Custard apple, Cherimoya, Llama, Soursop, and Biriba	Can be applied up to harvest	Refer to table 3	Apply up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> make applications with airblast sprayers. <b>DO NOT</b> make applications less than 14 days apart. <b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) in a single application and 6.1 fl oz/A (0.095 lb ai/A) per year, including preplant site preparation.
<b>Other Crops:</b> including Banana, Cacao, Coconut, Coffee, Date, Fig, Guayule, Indian Mulberry, Olive, Palm Heart, Persimmon, Pomegranate, Tea, and Vanilla	<b>DO NOT</b> apply within 3 days of harvest	Refer to table 3	Apply up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> make applications with airblast sprayers. <b>DO NOT</b> make applications less than 14 days apart. <b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per application and 7.9 fl oz/A (0.123 lb ai/A) per year, including preplant site preparation.

**DIRECTIONS FOR USE**

**PRODUCTION SYSTEMS**

Different production systems dictate different application techniques. Skirted trees are those allowing the lower branches of the trees to grow to the ground line. Non-skirted trees are grown in production systems where branches are pruned allowing access to the trunk area.

**Equipment and Application Skirted Orchards and Groves**

Hooded sprayers are required for AIM EC HERBICIDE applications in skirted trees. Refer to the HOODED SPRAYER APPLICATIONS section of this label.

**Non-Skirted Orchards and Groves**

Apply only by ground equipment including boom sprayers, shielded or hooded sprayers, hand-held or high-volume wands or orchard guns. Use a minimum of 20 gallons finished spray solution per broadcast acre.

**Weed Control**

Apply AIM EC HERBICIDE alone or as a tank mix with other registered herbicides to actively growing weeds. AIM EC HERBICIDE is a contact herbicide and coverage is essential for good weed control. Use a minimum of 20 gallons finished spray solution per broadcast acre.

**DO NOT** allow AIM EC HERBICIDE spray solution to contact green stem tissue, leaves, fruit or blooms of trees.

**AIM EC HERBICIDE Application Rates**

Apply AIM EC HERBICIDE up to 2 fl oz/A (0.031 lb ai/A) for postemergence control of susceptible broadleaf weeds. Refer to weed control list in Table 3 for appropriate weed control information. For best control, apply to seedling weeds in the 2 to 3-leaf stage. For larger weeds up to 6 leaves, use higher labeled rates of AIM EC HERBICIDE. Weeds greater than 6 leaves may be only partially controlled.

**Adjuvant Requirements**

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate at 1% v/v (one gallon COC per 100 gallons). AIM EC HERBICIDE may also be applied with labeled rates of MSO or silicone adjuvants.

**Tank Mix**

AIM EC HERBICIDE may be mixed with other herbicides that have preemergence or postemergence activity. AIM EC HERBICIDE only controls emerged vegetation. Any preemergence activity must rely on activity from registered preemergence herbicides mixed with AIM EC HERBICIDE. Contact herbicides e.g. glyphosate, glufosinate, and paraquat may be tank mixed with AIM EC HERBICIDE for broader spectrum weed control. See Mixing and Loading Instructions under the PRODUCT INFORMATION section of this label for specific mixing instructions. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner.

**Sucker Management**

AIM EC HERBICIDE is effective as an aid in the management of undesirable sucker growth from the base of the trunks or root sprouts. Apply AIM EC HERBICIDE at 2.0 fl oz/A (0.031 lb ai/A). Suckers and other undesirable growth must be treated when the tissue is young and not mature and/or hardened off. Care must be taken not to allow spray mist to contact desirable fruit, foliage or green stem tissue (see Precautions).

**Adjuvant Requirements**

Refer to adjuvant section of this label.

**Chemical Mowing**

Apply AIM EC HERBICIDE alone or in tank mixtures with other herbicides in chemical mowing practices for orchard vegetation management.

**Hooded Sprayer Application**

Apply AIM EC HERBICIDE with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

**Precautions**

Extreme caution must be used during applications when desirable fruit and/or foliage are present in order to avoid fruit spotting and/or leaf necrosis. **DO NOT** allow spray mist of AIM EC HERBICIDE to come in contact with green stem tissue, foliage, blooms or desirable fruit. On seedling or newly transplanted trees **DO NOT** allow spray to contact green bark of trunk area. For new seedlings up to 2 year old trees, the trunk base must be wrapped to help prevent chemical contact with the bark.

**Tank Mixture**

If AIM EC HERBICIDE is used in a tank mixture, observe the other product's label for restrictions, precautions, and rotational cropping instructions.

**Grasses: (Forage, Fodder, Hay, Seed and Sod)**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Postemerge Weed Control</b>	When AIM EC HERBICIDE is applied alone, grazing and hay operations may proceed with no restrictions.	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> make applications less than 7 days apart. <b>DO NOT</b> apply more than 5.9 fl oz/A (0.093 lb ai/A) per year. <b>DO NOT</b> make more than three applications per year.

**DIRECTIONS FOR USE**

Apply AIM EC HERBICIDE alone or in combination with other registered pesticides for the control of weeds in rangeland, pastures, hay, grasses grown for hay or silage and grass seed production and grass grown in Conservation Reserve Programs (CRP). Note that CRP usage must be in compliance with Federal, State, and local use guidelines.

Apply AIM EC HERBICIDE at use rates up to 2.0 fl oz/A (0.031 lb ai/A) per broadcast acre. For optimum results, weeds must be treated when small. Applications shall be made with ground equipment delivering a minimum of 10 gallons of finished spray per acre or aerial delivering a minimum of 3 gal/A of finished spray. Adjust sprayers to provide optimum coverage of the target weeds. Refer to weed control list in Table 3 for appropriate weed control information.

**Adjuvant Requirements**

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a quality nonionic surfactant (NIS) containing at least 80% active at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate (COC) at 1% v/v (one gallon COC per 100 gallons), or a methylated seed oil (MSO). The use of a high quality sprayable liquid nitrogen fertilizer at 2 to 4% v/v or ammonium sulfate (AMS) used at 2 to 4 pounds per acre in addition to the NIS, or MSO or COC is allowed.

When AIM EC HERBICIDE is applied alone, grazing and hay operations may proceed with no restrictions.

**Tank Mix**

AIM EC HERBICIDE may be tank mixed with other labeled herbicides to control weeds not listed on this label. Read and follow all manufacturers' label directions for the companion herbicide.

For tank mixture applications, refer to the use directions and restrictions of the mixture product.

**HOPS**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Post-Directed for Sucker Management</b>	<b>DO NOT</b> apply within 7 days of harvest.	Refer to table 3	2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> apply AIM EC HERBICIDE using air blast or air assisted sprayers. <b>DO NOT</b> apply through any type of irrigation system.
<b>Postemergence Weed Control</b>	<b>DO NOT</b> apply within 7 days of harvest.	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> apply more than 7.6 fl oz/A (0.12 lb ai/A) per year. Allow a minimum of 14 days between treatments of AIM EC HERBICIDE.

**DIRECTIONS FOR USE**

**Post-Directed Application for Sucker Management.**

AIM EC HERBICIDE is a contact herbicide for directed spray application to the basal portion of the hop plant for the management of sucker growth. Apply AIM EC HERBICIDE at 2.0 fl oz/A (0.031 lb ai/A) per application in a minimum of 20 gallons of spray solution by boom-type ground application equipment only to the basal portion of the hop plant (approximately the lower 1.5 feet) and to the sucker mat which extends from the base of the plant to approximately 1.5 to 2 feet into the row. An alternate row treatment program may be followed to avoid the removal of excessive photosynthetic capacity from the crown area by treating alternate rows on different days. Applications timing and techniques may vary from region to region. Please consult local university extension personnel for local management practices.

**Postemergence Control of Broadleaf Weeds**

Apply AIM EC HERBICIDE using shielded sprayers or hooded sprayers to control emerged and actively growing broadleaf weeds within or between the rows of the crop. Refer to Table 3 for appropriate weed control information.

**Adjuvant Requirements**

Coverage is essential to obtain good basal growth management. Use a nonionic surfactant (NIS) having at least 80 percent active ingredient at 0.25 % v/v (2 pints of NIS per 100 gallons of spray volume) or a quality crop oil concentrate (COC) at labeled rates. If AIM EC HERBICIDE is used in a tank mixture, refer to the other product labels for all restrictions on tank mixing and observe all label precautions, instructions and rotational cropping restrictions. For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$$

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Volume Per Acre} = \text{Band Volume}$$

**Application Precautions**

Extreme caution must be taken during application to avoid upward drift of the spray solution and contact with the highly susceptible new growth. Avoid applications until newly trained vines have developed sufficient barking to avoid damage to the stem and are high enough up the string to avoid contact with the apical bud.

**LOW GROWING BERRY Subgroup 13-07G**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Postemergence Weed Control</b>	Can be applied up to harvest	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> apply more than 2 fl oz/A (0.031 lb ai/A) during the dormant period. <b>DO NOT</b> apply more than 6.15 fl oz/A (0.096 lb ai/A) per year.
<p><b>DIRECTIONS FOR USE</b> AIM EC HERBICIDE applications will control susceptible emerged broadleaf weeds. Repeat applications may be necessary for weeds that emerge after an AIM EC HERBICIDE treatment.</p> <p><b>Equipment and Application</b> Apply only by ground equipment including boom sprayers, shielded or hooded sprayers, hand-held or high-volume wands or orchard guns. Use a minimum of 20 gallons finished spray solution per broadcast acre.</p> <p><b>Dormant Applications</b> Apply AIM EC HERBICIDE as a broadcast application to the base of the trunk to control emerged and actively growing weeds during the dormant stage of the crop.</p> <p><b>Post-directed Applications for Broadleaf Weed Control</b> Apply AIM EC HERBICIDE as a directed spray avoiding contact with the berry plant but directed at actively growing weeds. AIM EC HERBICIDE is a contact herbicide and coverage is essential for good weed control. <b>DO NOT</b> allow AIM EC HERBICIDE spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage.</p> <p>Newly planted bush berries must only be treated with shielded sprayers or hooded sprayers.</p> <p><b>AIM EC HERBICIDE Use Rates</b> Apply up to 2 fl oz/A (0.031 lb ai/A) AIM EC HERBICIDE per broadcast acre. For best control, apply to seedling weeds in the 2 to 3-leaf stage. Use higher labeled rates of AIM EC HERBICIDE for larger weeds up to 6 leaves. Weeds greater than 6 leaves may be only partially controlled. See Table 3 for AIM EC HERBICIDE use rates and weeds controlled.</p> <p><b>Adjuvant Requirements</b> A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.</p> <p>AIM EC HERBICIDE may be mixed with other registered herbicides for broader spectrum weed control. When tank mixing with fertilizer solutions, be sure to prepare an AIM EC HERBICIDE premixture of AIM EC HERBICIDE and clean water. See Mixing and Loading Instructions under the PRODUCT INFORMATION section of this label for specific mixing instructions. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner.</p> <p><b>Precautions</b> Extreme caution must be taken during applications when desirable fruit, foliage and/or blooms are present in order to avoid spotting or necrosis. <b>DO NOT</b> allow AIM EC HERBICIDE spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage. For seedling or newly transplanted bushes, <b>DO NOT</b> allow spray to contact green bark of trunk area. Use shielded sprayers only.</p> <p><b>Band Treatment Application</b> For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:</p> $\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$ $\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Volume Per Acre} = \text{Band Volume}$				

**MINT**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Broadcast</b>	<b>DO NOT</b> apply within 5 days of harvest.	Refer to table 3	Apply one application of AIM EC HERBICIDE at 0.5 to 1.92 fl oz/A (0.008 to 0.03 lb ai/A). Use higher rates when weeds are under stress or are larger.	<b>DO NOT</b> apply to actively growing crop. <b>DO NOT</b> apply more than 1.92 fl oz/A (0.03 lb ai/A) of AIM EC HERBICIDE per year.
<p><b>DIRECTIONS FOR USE</b> Apply AIM EC HERBICIDE as a broadcast application before Mint break dormancy for control of existing broadleaf weeds.</p> <p><b>Coverage is essential for good control.</b></p> <p><b>Adjuvant Requirements</b> Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed. Repeat application if necessary.</p> <p>For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section.</p>				

**PEANUT**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Postemergence Weed Control</b>	<b>DO NOT</b> apply within 7 days of harvest.	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> apply more than 6.1 fl oz/A (0.095 lb ai/A) per year. <b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per year as a harvest aid treatment.
<b>Harvest Aid</b>	<b>DO NOT</b> apply within 7 days of harvest.	Refer to table 3		<b>DO NOT</b> apply more than one harvest aid treatment per year. <b>DO NOT</b> feed immature peanut plant or peanut hay to livestock.

**DIRECTIONS FOR USE**

**Weed Control**

Apply AIM EC HERBICIDE alone or as a tank mixture with other herbicides as a postemergence treatment or as a hooded/directed spray treatment to control emerged and actively growing weeds. Apply hooded/directed applications of AIM EC HERBICIDE to middles (between rows of plants) and in strips (in row of plants). Apply AIM EC HERBICIDE at any time during the growing period (see precautions). AIM EC HERBICIDE may be mixed with other herbicides that have pre-emergence or post-emergence activity. Any pre-emergence activity must rely on activity from other herbicides as directed on their labels. Herbicides including glyphosate may be tank mixed with AIM EC HERBICIDE for broader spectrum weed control. If AIM EC HERBICIDE is used in a tank mixture, observe the other product's label for restrictions, precautions and rotational cropping instructions.

**Harvest Aid Application**

Apply AIM EC HERBICIDE as a harvest aid to defoliate and desiccate troublesome weeds that may be present at harvest. Apply AIM EC HERBICIDE alone or as a tank mixture with other peanut harvest aids.

**Adjuvant Requirements**

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a quality nonionic surfactant (NIS) containing at least 80% active at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate (COC) at 1% v/v (one gallon COC per 100 gallons), or a methylated seed oil (MSO). The use of a high quality sprayable liquid nitrogen fertilizer at 2 to 4% v/v or ammonium sulfate (AMS) used at 2 to 4 pounds per acre in addition to the NIS, or MSO or COC is allowed.

**Harvest Aid**

Apply AIM EC HERBICIDE as a harvest aid to defoliate and desiccate troublesome weeds that may be present at harvest. Apply AIM EC HERBICIDE alone or as a tank mixture with other peanut harvest aids.

Coverage is essential for satisfactory performance.

**Crop Rotation Restriction:**

After an application of this product to peanuts, you may only rotate the field to a carfentrazone-ethyl registered crop.

RICE: (Southern Use Only)

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Pre-flood Applications to Dry Seeded Rice</b>	<b>DO NOT</b> apply within 60 days of harvest once field is flooded.	See weed list in table 10 below	1.25 to 3.2 fl oz/A (0.0195 to 0.05 lb ai/A)	<b>DO NOT</b> apply when conditions favor drift or when wind is above 10 mph. Pre-flood treatment, once field is flooded, water must be held for at least 23 days following treatment before release. <b>DO NOT</b> apply more than 8.8 fl oz/A (0.138 lb ai/A) per year including fallow/preplant burndown and other labeled in-crop applications.
<b>Post Flood Applications to Exposed Weeds</b>		See weed list in table 11 below	1.6 to 6.4 fl oz/A (0.025 to 0.10 lb ai/A)	<b>DO NOT</b> apply when conditions favor drift or when wind is above 10 mph. <b>DO NOT</b> apply more than 8.8 fl oz/A (0.138 lb ai/A) per year including fallow/preplant burndown and other labeled in-crop applications. <b>DO NOT</b> release water for at least 23 days following a Post Flood treatment in the water.
<b>Harvest Aid</b> (not permitted in California)	<b>DO NOT</b> apply no earlier than soft dough up to the 3 days of harvest.	Desiccate troublesome broadleaf weeds e.g. hemp sesbania, Indian and northern jointvetch, morningglories, and pigweeds.	1.25 to 1.6 fl oz/A (0.0195 to 0.025 lb ai/A)	<b>DO NOT</b> apply when conditions favor drift or when wind is above 10 mph. <b>DO NOT</b> apply more than 1.47 fl oz/A (0.023 lb ai/A) as a harvest aid.

**DIRECTIONS FOR USE**

Apply AIM EC HERBICIDE alone or as a tank mixture with other rice herbicides to emerged and actively growing weeds. Apply AIM EC HERBICIDE with either ground or aerial spray equipment.

**DO NOT** apply when conditions favor drift.

To control weeds not listed on this label, AIM EC HERBICIDE may be tank mixed with other herbicides registered for use on rice. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions, and rotational cropping restrictions.

**Postemergence Pre-flood Applications to Dry Seeded Rice**

Apply AIM EC HERBICIDE at 1.25 to 3.2 fl oz/A (0.0195 to 0.05 lb ai/A). Use a minimum of 10 gallons of finished spray per acre for ground application equipment, and a minimum of 3 gallons per acre of finished spray for aerial equipment. For optimum results, apply AIM EC HERBICIDE to weeds up to 4 inches tall. Use a quality nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. For more active treatments, use a Crop Oil Concentrate (COC) at 0.5 to 1.0% v/v (one half to one gallon per 100 gallons). Apply when the rice is at the 2 leaf stage or larger, but prior to flooding. Some leaf speckling may occur. Once field is flooded, water must be held for at least 23 days following treatment before release.

**When used as directed AIM EC HERBICIDE will provide Control of listed weeds up to 4 inches tall.**

**Table 10:**

Cocklebur, common	Morningglory, spp.
Copperleaf, hophornbeam	Pigweed spp.
Dayflower, spreading	Purslane, common
Groundcherry, cutleaf	Redweed
Hyssop, water	Sesbania, hemp
Jointvetch, Indian	Smartweed, PA (seedling)
Jointvetch, northern	

**Suppression of listed weeds**

Alligatorweed	Flatsedge, rice
Ducksalad	Redstem
Eclipta	Texasweed

**Tank Mix**

For control of weeds listed as suppressed or not listed on this label, apply AIM EC HERBICIDE following a preemergence grass herbicide or tank with other rice herbicides for broad spectrum weed control. Use tank mix applications when rice is well established and in the appropriate stage of growth for treatment with AIM EC HERBICIDE and the tank mix partner. For optimum results, weed species must also be in the proper stage of growth as specified on the AIM EC HERBICIDE and tank mix partner label. Read and follow all manufacturers' label directions for the companion herbicide except for specific directions on this label. **DO NOT** add a surfactant or crop oil concentrate when tank mixing herbicides formulated as emulsifiable concentrates unless required by the tank mix partners label. For other herbicide tank mix partners that are not Emulsifiable concentrates refer to their label for specific adjuvant specifications. Observe all applicable directions, restrictions and precautions on the partner herbicide labels.

**Post Flood Applications to Exposed Weeds**

For post flood applications apply AIM EC HERBICIDE to rice and weeds after the establishment of the permanent flood and when 80% of the foliage of the weeds are exposed. Apply AIM EC HERBICIDE at 1.6 to 6.4 fl oz/A (0.025 to 0.10 lb ai/A) to actively growing weeds. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. For more active treatments, use a Crop Oil Concentrate (COC) at 1.0% v/v (one gallon per 100 gallons). Apply when the rice is at the 2-leaf stage or later. Use a minimum of 10 gallons of finished spray per acre for ground application equipment and a minimum of 3 gallons of finished spray per acre for aerial application equipment. For optimum results, make applications to small rather than larger weeds. If water level has been lowered to allow this treatment, it must be returned to normal levels 24 hours following treatment. Users of AIM EC HERBICIDE must hold the water on the rice fields for 23 days following treatment.

**When used as directed, AIM EC HERBICIDE will provide control of listed weeds.**

**Table 11:**

Arrowhead, annual	Morningglory spp.
Jointvetch, Indian	Sesbania, hemp
Jointvetch, northern	

**Suppression of listed weeds up to 4 inches.**

Alligatorweed	Ducksalad
Ammannia, purple	Flatsedge, rice
Dayflower, spreading	Texasweed

**Harvest Aid Application:**

AIM EC HERBICIDE is effective as a harvest aid to defoliate and desiccate troublesome weeds that may be present at harvest. Apply AIM EC HERBICIDE alone or as a tank mixture with other rice harvest aids. Harvest aid treatment applications may be made no earlier than soft dough up to the 3 day PHI. Refer to Table 1 for maximum use rate as harvest aid.

**Harvest Aid Restriction: not permitted in California****Crop Rotation Restriction:**

After an application of this product to rice, you may only rotate the field to a carfentrazone-ethyl registered crop.

**Rice, Wild: Wild rice grown in cultivated fields where the water discharge / release can be controlled.**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Postemerge Weed Control</b>	<b>DO NOT</b> apply within 60 days of harvest.	See weed list in table 14.	6.4 to 12.0 fl oz/A (0.1 to 0.19 lb ai/A)	<p><b>DO NOT</b> apply when conditions favoring drift exist.</p> <p><b>DO NOT</b> apply when winds exceed 10 mph.</p> <p><b>DO NOT</b> apply more than 12.0 fl oz/A (0.19 lb ai/A) per application.</p> <p><b>DO NOT</b> apply more than 19.2 fl oz/A (0.3 lb ai/A) per year, including fallow/preplant, burndown, and labeled in-crop applications.</p> <p><b>DO NOT</b> apply during the floating leaf stage when exposed wild rice leaves are most susceptible to injury.</p> <p><b>DO NOT</b> apply to wild rice when there is heavy dew on the leaves or under high humidity conditions.</p> <p><b>DO NOT</b> release flood water off wild rice field(s) for a minimum of 23 days after application of this product.</p> <p><b>DO NOT</b> apply during the floating leaf stage when exposed wild rice leaves are most susceptible to injury. (Specific to California)</p> <p><b>DO NOT</b> apply within 0.5 mile of sensitive crops (for California wild rice).</p> <p><b>DO NOT</b> apply to wild rice by air in California</p> <p>Do not release flood water off wild rice field(s) for a minimum of 23 days after application of this product in California.</p>

**DIRECTIONS FOR USE**

Apply AIM EC HERBICIDE alone or as a tank mixture with other rice herbicides to emerged and actively growing weeds. Wild rice must be well rooted and vigorously growing at the time of application. Earlier applications may cause unacceptable crop response. Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons per acre.

Apply AIM EC HERBICIDE to weeds at the rate of 6.4 to 12.0 fl oz/A (0.1 to 0.19 lb ai/A) to the foliage of exposed weeds above the water surface. Make applications after the floating leaf stage through tillering. The water in paddies may be lowered if practical. Smaller weeds with more leaf area exposed will give better control. If water is lowered for application, it may be re-flooded to normal depths 24 hours after the application

**When used as directed AIM EC HERBICIDE will provide control or suppression of the following weeds.**

**Table 14:**

Ammannia, purple (Suppression only)	Burrweed, giant (Suppression only)
Ammannia, redstem (Suppression only)	Umbrellaplant, smallflower (Suppression only)
Arrowhead, California	Waterplantain, common (Suppression only)
Bulrush, ricefield	

**Crop Response**

Some temporary leaf specking may occur following application.

**Tank Mix**

AIM EC HERBICIDE may be tank mixed with other herbicides to control weeds not listed on this label. Not all combinations of AIM EC HERBICIDE and other formulated herbicides and adjuvants have been tested. In general, EC formulations, nonionic and silicone based surfactants, and crop oil concentrates, will increase leaf speckling on the wild rice leaves. These tank mixes must be tested on a small portion of the field to ensure crop safety prior to general use.

**Precautions**

Wet leaf surfaces at the time of application can cause unacceptable injury.

**Crop Rotation Restriction**

After an application of this product to wild rice, you may only rotate the field to a carfentrazone-ethyl registered crop.

**SMALL GRAINS**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
Preplant Burndown	NA	Refer to table 3	Up to 1.0 fl oz/A (0.016 lb ai/A)	<b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per application. <b>DO NOT</b> apply when conditions favor drift.
Postemergence	Except Winter Wheat – jointing stage Winter Wheat – boot stage	Refer to table 3	0.5 to 1.0 fl oz/A (0.008 to 0.016 lb ai/A)	<b>DO NOT</b> harvest for forage within 7 days of application. <b>DO NOT</b> apply more than 1.0 fl oz/A of AIM EC HERBICIDE (0.016 lb ai/A) per year including fallow, preplant burndown, or labeled in-crop applications.
Harvest Aid Applications	<b>DO NOT</b> apply within 7 days of harvest.	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	

**DIRECTIONS FOR USE**

**Timing and method of application:**

AIM EC HERBICIDE may be applied preplant (up to 1 day before seeding), postemergence or harvest aid. For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. For dense weed pressure, use the higher labeled application rate plus tank mix combinations. **Coverage is essential for good control.** Refer to Table 3 for weeds controlled at labeled rates of AIM EC HERBICIDE. For broader spectrum weed control, AIM EC HERBICIDE may be tank mixed with other herbicides registered for use in small grains.

**Pre Plant Burndown:**

Refer to the pre plant burndown section of this label.

**Postemergence Application:**

Postemergence application may be made from 4-inches tall to just prior to the boot stage, **DO NOT** apply more than 0.016 lb ai/A including preplant and postemergent application (not including harvest aid). **DO NOT** apply more than 0.016 lb ai/A as a harvest aid treatment.

**AIM EC HERBICIDE Use Rate**

Apply from 0.5 to 1.0 fl oz/A AIM EC HERBICIDE (0.008 – 0.016 lb ai/A). Use a minimum finished spray solution of 10 gallons per acre by ground or 3 gallons per acre by air. Up to half of the spray volume (by air or ground) may be liquid nitrogen fertilizer.

**Adjuvant Requirements**

Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. The use of a high quality sprayable liquid nitrogen fertilizer (2 to 4% v/v or 2 to 4 gallons per 100 gallon spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant is allowed.

**DO NOT** use AIM EC HERBICIDE with crop oil concentrates (COC), methylated seed oils (MSO) or silicone based adjuvants for postemergence applications.

**Tank Mix**

To control weeds not listed on this label, AIM EC HERBICIDE may be tank mixed with other registered herbicides.

For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions, and rotational cropping restrictions. Use aerial or ground equipment for AIM EC HERBICIDE applications. **Coverage is essential for good control.** Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre. Applications made by air shall utilize a minimum finished spray volume of 3 gallons per acre. Up to half of the spray volume (by air or ground) may be liquid nitrogen fertilizer. Refer to Table 3 for appropriate weed control information.

**AIM EC HERBICIDE Plus 2,4-D (amine or ester) or MCPA (amine or ester)**

AIM EC HERBICIDE may be tank mixed at a rate of 0.5 to 1.0 fl oz/A (0.008-0.016 lb ai/A) with 2,4-D (amine or ester) or MCPA (amine or ester) for use on small grains. For optimum results add 2,4-D (amine or ester) to the tank at 0.25 lb acid equivalent per acre or MCPA (amine or ester) at 0.375 lb acid equivalent per acre. Higher rates of these herbicides are allowed, but **DO NOT** exceed the label use rates allowed by these labels. Add nitrogen fertilizer (2 to 4% v/v) 2 to 4 gallons per 100 gallons or ammonium sulfate 4 lbs. per acre) to the tank mixture.

**When applied as directed, AIM EC HERBICIDE in tank mixtures with 2,4-D (amine or ester) or MCPA (amine or ester) herbicides will provide control of listed weeds up to 4 inches tall.**

**Table 15:**

Amaranthus spp.	Nightshade, black
Bedstraw, catchweed	Pennycress, field **
Buckwheat, wild	Pepperweed, greenflower**
Cocklebur	Pigweed, prostrate
Croton, woolly	Pigweed, redroot
Fiddleneck	Pigweed, smooth
Filaree, redstem	Primrose, cutleaf
Flixweed**	Primrose, tumble
Gromwell, common	Radish, wild
Groundsel, common	Ragweed, common
Knotweed, prostrate*	Ragweed, giant
Kochia	Rocket, London
Lambsquarters, common	Sowthistle, annual
Lettuce, miners	Speedwell, ivyleaf
Lettuce, prickly	Sunflower, wild
Mustard, blue***	Tarweed, coast
Mustard, tansy***	Thistle, Russian
Mustard, tumble**	Wallflower, bushy
Mustard, wild**	Waterhemp, tall

\*For Knotweed control, use AIM EC HERBICIDE + 2,4-D (amine or ester) only.

\*\*These weeds can be treated from the rosette through bolting growth stages.

\*\*\*Apply to rosette growth stage (before bolting) of blue mustard.

#### **Harvest Aid**

Apply up to 2.0 fl oz/A AIM EC HERBICIDE, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE AIM EC HERBICIDE USE RATE and the PREHARVEST INTERVAL Table (Table 2) for additional application information. If treatments of AIM EC HERBICIDE have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application.

#### **Adjuvant Requirements – Harvest Aid**

A methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use methylated seed oil, or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil is allowed.

**Coverage is essential for satisfactory performance.**

**SORGHUM (Grown for Grain and Seed)**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Pre Plant Burndown</b>	NA	Refer to table 3	Up to 1.0 fl oz/A (0.016 lb ai/A)	<b>DO NOT</b> make foliar broadcast applications to forage sorghum or sorghum grown for seed.
<b>Foliar Broadcast Application (Grain Sorghum Only)</b>	<b>DO NOT</b> apply past pre-boot stage	Refer to table 3 for weeds controlled at 0.5 fl oz/A per acre rate.	Up to 0.5 fl oz/A (0.008 lb ai/A)	<b>DO NOT</b> apply more than 1.0 fl oz/A (0.016 lb ai/A) per year including fallow, preplant burndown and labeled applications to the growing crop (not including Harvest Aid treatments). See Table 1.
<b>Directed or Shielded Spray Applications.</b>	<b>DO NOT</b> apply past pre-boot stage (Grain)	Refer to table 3	Up to 1.0 fl oz/A (0.016 lb ai/A)	
<b>Harvest Aid</b>	<b>DO NOT</b> apply within 3 days of harvest.	Desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf.	Up to 1.0 fl oz/A (0.016 lb ai/A)	<b>DO NOT</b> apply more than 1.0 fl oz/A (0.016 lb ai/A) per year as a Harvest Aid treatment. See Table 1.

**DIRECTIONS FOR USE**

AIM EC HERBICIDE may be applied to grain and forage sorghum as a pre plant burndown; a hooded or shielded spray; and a post directed spray. In addition to these applications methods, AIM EC HERBICIDE may be applied to grain sorghum (sorghum grown for grain but not for seed production) as a foliar broadcast and harvest aid treatment. See Table 1 for Maximum Allowable AIM EC HERBICIDE Use Rate and Table 3 for weeds controlled at labeled rates of AIM EC HERBICIDE on sorghum.

**PRE PLANT BURNDOWN**

See instructions under the Pre Plant Burndown section of this label.

**FOLIAR BROADCAST (Grain Sorghum Only)**

Apply to grain sorghum from 4 inches tall to just prior to the boot stage. AIM EC HERBICIDE may be applied alone or as a tank mixture with other herbicides labeled for use on sorghum. Broadcast applications of AIM EC HERBICIDE to sorghum with wet foliage or application during periods of adverse environmental conditions including cool, cloudy, wet, or high humidity may cause increased crop response. Directed sprays are suggested under these conditions. For additional information on crop response, refer to the PRODUCT INFORMATION section of the AIM EC HERBICIDE label.

**AIM EC HERBICIDE Use Rates – Foliar Grain Only**

**DO NOT** exceed 0.5 fl oz/A (0.008 lb ai/A) AIM EC HERBICIDE. See Table 3 for weeds controlled at 0.5 fl oz/A of AIM EC HERBICIDE. Rates below 0.5 fl oz/A may not fully control weeds.

**Adjuvant Requirements – Foliar Grain Only**

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. **DO NOT** use crop oil concentrates or methylated seed oils for broadcast applications on emerged sorghum.

**Tank Mix –Foliar Grain Only**

For control of additional broadleaf weeds and grasses, AIM EC HERBICIDE may be tank mixed with 2,4-D (amine), atrazine, dicamba, quinclorac, prosulfuron, halosulfuron, or fluroxypyr products. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner.

Leaf speckling can occur when AIM EC HERBICIDE is used with certain formulations of crop protection products and adjuvants.

**DIRECTED OR SHIELDED SPRAY APPLICATIONS**

Apply AIM EC HERBICIDE when the sorghum is at least 4 inches tall to prior to the boot stage. Use drop nozzles or other sprayers capable of directing the spray to the target weeds and away from the whorl and leaves of the sorghum plant. Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons per acre. Refer to Table 3 for weeds controlled at labeled rates of AIM EC HERBICIDE. Coverage is essential for good control. Directed, shielded, or hooded sprayers are required for post emergence treatments to forage sorghum and sorghum grown for seed.

**AIM EC HERBICIDE Use Rates – Directed or Shielded Spray**

Apply up to 1.0 fl oz/A AIM EC HERBICIDE (0.016 lb ai/A) using directed or shielded sprayers.

**Adjuvant Requirements – Directed or Shielded Spray**

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. Crop oil concentrates or methylated seed oils may increase crop injury on sorghum.

**Tank Mix – Directed or Shielded Spray**

For control of additional broadleaf weeds and grasses, AIM EC HERBICIDE may be tank mixed with 2,4-D (amine), atrazine, dicamba, quinclorac, prosulfuron, halosulfuron, or fluroxypyr products. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner.

**HOODED SPRAYER APPLICATION**

Apply AIM EC HERBICIDE with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

**HARVEST AID (WEED CONTROL)**

Apply AIM EC HERBICIDE to defoliate and/or desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf that may be present at harvest. Apply up to 1.0 fl oz/A AIM EC HERBICIDE, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE AIM EC HERBICIDE USE RATE and the PREHARVEST INTERVAL Table (Table 2) for additional application information. If treatments of AIM EC HERBICIDE have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment.

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application.

**Adjuvant Requirements – Harvest Aid**

A methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use methylated seed oil, or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil is allowed.

**Coverage is essential for satisfactory performance.**

**PRECAUTIONS**

**DO NOT** use crop oil concentrates or methylated seed oils for broadcast applications on emerged sorghum.

Leaf speckling can occur when AIM EC HERBICIDE is used with certain formulations of crop protection products and adjuvants.

**SOYBEANS**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
Preplant Burndown	NA	Refer to table 3	Up to 1.5 fl oz/A (0.023 lb ai/A)	<b>DO NOT</b> apply more than 1.5 fl oz/A (0.023 lb ai/A) per year.
Postemergence (Broadcast)	V10	Refer to table 3	0.25 to 0.5 fl oz/A (0.004 to 0.008 lb ai/A). (See Directions for Use below for details).	<b>DO NOT</b> feed treated soybean forage or hay to livestock. <b>DO NOT</b> use with diphenylether herbicides.
Postemergence (Directed Spray and Hooded Sprayer Applications)	V10	Refer to table 3	Up to 1.5 fl oz/A (0.023 lb ai/A)	<b>DO NOT</b> apply when conditions favoring drift exist. <b>DO NOT</b> apply when crop foliage is wet from dew, rainfall or irrigation.
Harvest Aid	<b>DO NOT</b> apply within 3 days of harvest	Refer to table 3	Up to 1.5 fl oz/A (0.023 lb ai/A)	

**DIRECTIONS FOR USE**

Apply AIM EC HERBICIDE alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to soybeans in all tillage systems from prior to planting up to prior to emergence. **DO NOT** apply AIM EC HERBICIDE during a period from emergence to V2. After plants have reached V3, applications are allowed up to V10.

For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. Use the higher rates when treating more mature weeds or dense vegetative growth. **Coverage is essential for good control.** Refer to weed control list in Table 3 for appropriate weed control information.

**Broadcast Postemergence Application**

Apply AIM EC HERBICIDE at 1.5 fl oz/A (0.023 lb ai/A) for the control of velvetleaf. **DO NOT** apply AIM EC HERBICIDE to soybeans with maturities less than Group 2.0. For soybeans of maturity Group 2.1 to 3.4, apply AIM EC HERBICIDE at rates up to 1.5 fl oz/A (0.023 lb ai/A). Use caution when making applications when making these treatments.

For soybeans maturing later than Group 3.5, apply AIM EC HERBICIDE at rates up to 1.5 fl oz/A (0.023 lb ai/A).

**Adjuvant Requirements**

Use NIS only as the adjuvant for this treatment at the rate of 0.25% v/v (2 pints per 100 gallons of spray solution).

**Broadcast Application Precaution**

The application of AIM EC HERBICIDE to soybeans may result in crop response. Soybeans may show some burn, speckling or necrosis of crop leaves. Soybeans quickly outgrow initial herbicide effects and yields are not affected. **DO NOT** make applications during conditions of abnormal cool, high humidity or if foliage is wet from dew, rainfall or irrigation. Users must be aware of these potential effects prior to making applications. If the user is not willing to accept these risks, applications must not be made.

For additional information on crop response, refer to the PRODUCT INFORMATION section of this label.

**Tank Mix**

AIM EC HERBICIDE may be tankmixed with other herbicides to control weeds not listed on this label. **Do not use with diphenylether herbicides.** Read and follow all manufacturers' label directions for the mixture herbicide except for specific directions on this label. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. For control of additional broadleaf weeds and grasses, AIM EC HERBICIDE may be tankmixed with glyphosate or glufosinate products for use on GMO soybeans. Leaf injury can occur when AIM EC HERBICIDE is used with certain formulations of crop protection products and adjuvants. Refer to the Tank Mixtures and Required Adjuvants sections under PRODUCT INFORMATION.

**When used as directed AIM EC HERBICIDE at 0.25 fl oz/A (0.004 lb ai/A) will provide:**

**Control of listed weeds up to 4 inches tall.**

Velvetleaf	
------------	--

**When used as directed, AIM EC HERBICIDE at 0.5 fl oz/A (0.008 lb ai/A) will provide:**

**Control of weeds up to 4 inches tall, or as specified.**

Lambsquarters, common	Nightshade, black
Morningglory, Pitted (up to 3 true leaves)	Pigweed, redroot
Morningglory, Ivyleaf (up to 3 true leaves)	Waterhemp, spp. (up to 3 inches tall)

**Hooded Sprayer Application**

Apply AIM EC HERBICIDE with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications of this label for additional specific use directions.

**Directed Sprayer Application**

Use AIM EC HERBICIDE at 0.5 to 1.5 fl oz/A (0.008 to 0.023 lb ai/A). Applications must be made by ground equipment using a finished volume of 10 to 20 gallons of spray per acre. When soybeans are grown under very dry soil moisture conditions, the use of a high quality sprayable liquid nitrogen fertilizer (2 to 4% v/v) or 2 to 4 gallons per 100 gallon spray solution) used in addition to the nonionic surfactant is allowed. Apply as a post-directed treatment with spray directed toward the base of the plant and avoid contact with soybean foliage. The use of spray shields may reduce spray contact with soybean foliage. AIM EC HERBICIDE contact with soybean foliage can result in significant crop response.

**SUGARCANE:**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Postemergence Treatment or Hooded/directed Spray</b>	<b>DO NOT</b> apply within 7 days of harvest.	Refer to table 3	Up to 2.0 fl oz/A (0.031 lb ai/A)	<b>DO NOT</b> apply more than 6.1 fl oz/A (0.095 lb ai/A) per year. <b>DO NOT</b> apply more than one harvest aid treatment per year.
<b>Harvest Aid</b>	<b>DO NOT</b> apply within 7 days of harvest.	Desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf.	1.0 - 2.0 fl oz/A (0.016 0.031 lb ai/A)	<b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per year as a harvest aid treatment.

**DIRECTIONS FOR USE**

**Postemergence/Hood Spray Application**

Apply AIM EC HERBICIDE alone or as a tank mixture with other herbicides as a postemergence treatment or as a hooded/directed spray treatment to control emerged and actively growing weeds. Apply AIM EC HERBICIDE up to 2.0 fl oz/A (0.031 lb ai/A). Apply hooded/directed applications of AIM EC HERBICIDE to middles (between rows of plants) and in strips (in row of plants). Apply AIM EC HERBICIDE at any time during the growing season (see precautions). AIM EC HERBICIDE may be mixed with other herbicides that have pre-emergence or post-emergence activity. Any preemergence activity must rely on activity from other herbicides as directed on their labels. Herbicides including glyphosate may be tank mixed with AIM EC HERBICIDE for broader spectrum weed control. If AIM EC HERBICIDE is used in a tank mixture, observe the other product's label for restrictions, precautions and rotational cropping instructions.

**Harvest Aid Application**

AIM EC HERBICIDE is effective as a harvest aid to defoliate and desiccate troublesome weeds that may be present at harvest. Apply AIM EC HERBICIDE alone or as a tank mixture with other sugarcane harvest aids.

**Adjuvant Requirements**

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a quality nonionic surfactant (NIS) containing at least 80% active at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate (COC) at 1% v/v (one gallon COC per 100 gallons), or a methylated seed oil (MSO). The use of a high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v or ammonium sulfate (AMS) used at 2 to 4 pounds per acre in addition to the NIS, or MSO or COC is allowed.

**Crop Rotation**

After an application of AIM EC HERBICIDE to sugarcane, you may only rotate the field to a carfentrazone-ethyl registered crop.

**TEFF (Grain and Forage)**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Pre Plant Burndown</b>	NA	Refer to table 3.	Up to 1.0 fl oz/A (0.016 lb ai/A)	<b>DO NOT</b> make foliar broadcast applications to forage teff or teff grown for seed.
<b>Foliar Broadcast Application (GrainTeff Only).</b>	Jointing Stage	Refer to table 3 for weeds controlled at 0.5 fl oz/A per acre rate.	Up to 0.5 fl oz/A (0.008 lb ai/A)	<b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per year including fallow, preplant burndown and labeled applications to the growing crop (not including Harvest Aid treatments). See Table 1.
<b>Directed or Shielded Spray Applications.</b>	Jointing Stage	Refer to table 3	Up to 1.0 fl oz/A (0.016 lb ai/A) per acre	
<b>Harvest Aid - Forage</b>	<b>DO NOT</b> apply within 7 days of harvest.	Desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf.	Up to 2.0 fl oz/A (0.031lb ai/A)	<b>DO NOT</b> apply more than 2.0 fl oz/A (0.031 lb ai/A) per year as a Harvest Aid treatment. See Table 1.
<b>Harvest Aid - Forage</b>	<b>DO NOT</b> apply within 3 days of harvest.	Desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf.	Up to 2.0 fl oz/A (0.031 lb ai/A)	

**DIRECTIONS FOR USE**

AIM EC HERBICIDE may be applied to grain and forage teff as a pre plant burndown; a hooded or shielded spray; and a post directed spray. In addition to these applications methods, AIM EC HERBICIDE may be applied to grain teff (teff grown for grain but not for seed production) as a foliar broadcast and harvest aid treatment. See Table 1 for Maximum Allowable AIM EC HERBICIDE Use Rate and Table 3 for weeds controlled at labeled rates of AIM EC HERBICIDE on teff.

**PRE PLANT BURNDOWN**

See instructions under the Pre Plant Burndown section of this label.

**FOLIAR BROADCAST (Grain Teff Only)**

Apply to grain teff from 4 inches tall to just prior to the boot stage. AIM EC HERBICIDE may be applied alone or as a tank mixture with other herbicides labeled for use on teff. Broadcast applications of AIM EC HERBICIDE to teff with wet foliage or application during periods of adverse environmental conditions including cool, cloudy, wet, or high humidity may cause increased crop response. Directed sprays are suggested under these conditions. For additional information on crop response, refer to the PRODUCT INFORMATION section of the AIM EC HERBICIDE label.

**AIM EC HERBICIDE Use Rates – Foliar Grain Only**

**DO NOT** exceed 0.5 fl oz/A (0.008 lb ai/A) AIM EC HERBICIDE. See Table 3 for weeds controlled at 0.5 fl oz/A of AIM EC HERBICIDE. Rates below 0.5 fl oz/A may not fully control weeds.

**Adjuvant Requirements – Foliar Grain Only**

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. **DO NOT** use crop oil concentrates or methylated seed oils for broadcast applications on emerged teff.

**Tank Mix –Foliar Grain Only**

For control of additional broadleaf weeds and grasses, AIM EC HERBICIDE may be tank mixed with 2,4-D (amine), atrazine, dicamba, quinclorac, prosulfuron, halosulfuron, or fluroxypyr products. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner. Leaf speckling can occur when AIM EC HERBICIDE is used with certain formulations of crop protection products and adjuvants.

**DIRECTED OR SHIELDED SPRAY APPLICATIONS**

Apply AIM EC HERBICIDE when the teff is at least 4 inches tall to prior to the boot stage. Use drop nozzles or other sprayers capable of directing the spray to the target weeds and away from the whorl and leaves of the teff plant. Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons per acre. Refer to Table 3 for weeds controlled at labeled rates of AIM EC HERBICIDE.

**Coverage is essential for good control.** Directed, shielded, or hooded sprayers are required for post emergence treatments to forage teff and teff grown for seed.

**AIM EC HERBICIDE Use Rates – Directed or Shielded Spray**

Apply up to 1.0 fl oz/A AIM EC HERBICIDE (0.016 lb ai/A) using directed or shielded sprayers.

**Adjuvant Requirements – Directed or Shielded Spray**

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. Crop oil concentrates or methylated seed oils may increase crop injury on teff.

**Tank Mix – Directed or Shielded Spray**

For control of additional broadleaf weeds and grasses, AIM EC HERBICIDE may be tank mixed with 2,4-D (amine), atrazine, dicamba, quinclorac, prosulfuron, halosulfuron, or fluroxypyr products. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner.

**HOODED SPRAYER APPLICATION**

Apply AIM EC HERBICIDE with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

**HARVEST AID (WEED CONTROL)**

Apply AIM EC HERBICIDE to defoliate and/or desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf that may be present at harvest. Apply up to 1.0 fl oz/A AIM EC HERBICIDE, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE AIM EC HERBICIDE USE RATE and the PREHARVEST INTERVAL Table (Table 2) for additional application information. If treatments of AIM EC HERBICIDE have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment.

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application.

**Adjuvant Requirements – Harvest Aid**

A methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use methylated seed oil, or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil is allowed.

**Coverage is essential for satisfactory performance.**

**PRECAUTIONS**

**DO NOT** use crop oil concentrates or methylated seed oils for broadcast applications on emerged teff.

Leaf speckling can occur when AIM EC HERBICIDE is used with certain formulations of crop protection products and adjuvants.

**TOBACCO**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Postemerge Weed Control</b> (pre-transplant, shielded/hooded spray, directed spray)	<b>DO NOT</b> apply within 6 days of harvest.	Refer to table 3	Up to 1.5 fl oz/A (0.023 lb ai/A)	<b>DO NOT</b> apply more than 3.2 fl oz/A (0.05 lb ai/A) per year.

**DIRECTIONS FOR USE**

Apply AIM EC HERBICIDE alone or as a tank mixture with other registered herbicides to emerged and actively growing weeds. For optimum performance, make applications to weeds up to 4 inches tall and rosettes less than 3 inches across. Use higher rates when treating more mature weeds or dense vegetative growth.

**Coverage is essential for good control.**

**Adjuvant Requirements**

Use adequate spray volume to achieve thorough coverage, but a minimum of 10 gallons of finished spray per acre is required. Use a quality crop oil concentrate (COC) at 1% v/v (1 gallon of COC per 100 gallons of spray solution).

AIM EC HERBICIDE may be tank mixed with other herbicides registered for use on tobacco to provide additional weed control. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. Refer to the other product label for restrictions on tank mixing and observe all label precautions, instructions and rotational cropping restrictions.

For additional information refer to the PRODUCT INFORMATION section of the AIM EC HERBICIDE label.

**Pre-transplant burndown**

AIM EC HERBICIDE is a contact herbicide for pre-transplant burndown control of broadleaf weeds in tobacco. Apply AIM EC HERBICIDE as a broadcast application alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply AIM EC HERBICIDE up to one (1) day prior to transplanting.

**Shielded spray or Hooded spray**

Apply AIM EC HERBICIDE using shielded sprayers or hooded sprayers to emerged and actively growing broadleaf weeds in tobacco from transplanting until layby. Shielded spray or hooded spray applications of AIM EC HERBICIDE or AIM EC HERBICIDE tank mixtures must utilize application equipment that must prevent contact of spray solution with the tobacco plant. **DO NOT** allow spray solution to contact tobacco foliage or green stem tissue. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

**Directed spray after first priming (Flue Cured Tobacco Only)**

Apply AIM EC HERBICIDE as a directed spray application after the first priming in only flue cured tobacco only for the control of emerged and actively growing broadleaf weeds. Directed spray equipment must position nozzles a minimum of 3 to 4 inches above the soil, with nozzles directed underneath the crop canopy. Spray solution must be directed at the base of tobacco plants for minimal contact with foliage while maintaining maximum contact with broadleaf weeds that are at appropriate treatment size. **DO NOT** apply when conditions favor drift or wind is above 10 mph.

For control of additional broadleaf weeds and grasses, AIM EC HERBICIDE may be tank mixed with other herbicides registered for use in tobacco at the appropriate timing. Refer to weed control list in Table 3 for appropriate weed control information. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

**TUBEROUS AND CORM VEGETABLES (SUBGROUP 1C & 1D – EXCEPT POTATO)**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<p><b>Fallow Systems</b> See the <b>Fallow Systems</b> section for directions for application.</p> <p><b>Preplant Burndown</b> See the <b>Preplant Burndown</b> section for directions for application.</p>	<p><b>DO NOT</b> apply within 7 days of harvest.</p>	<p>Refer to table 3</p>	<p>Up to 2.0 fl oz/A (0.031 lb ai/A)</p>	<p><b>DO NOT</b> apply more than 11.6 fl oz/A of (0.181 lb ai/A) per crop year as a desiccant.</p> <p><b>DO NOT</b> apply when conditions favor drift or wind is above 10 mph.</p>
<p><b>Harvest Aid</b></p>	<p><b>DO NOT</b> apply within 7 days of harvest.</p>	<p>Refer to table 3</p>	<p>3.2 to 5.8 fl oz/A (0.05 to 0.09 lb ai/A)</p> <p>2.0 – 5.8 oz/A (0.031 to 0.09 lb ai/A) with other registered potato desiccants.</p>	

**DIRECTIONS FOR USE**

Apply AIM EC HERBICIDE alone or in a tank mix combination with other herbicides and insecticides as a fallow systems treatment, as a preplant burndown treatment and/or as a harvest aid to desiccate potatoes and those susceptible weeds that may be present.

**Fallow Systems**

Apply AIM EC HERBICIDE by ground or air alone or with other herbicides in the fallow period prior to planting or the emergence of any crop listed on this label to control or suppress weeds. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. **Coverage is essential for good weed control.**

AIM EC HERBICIDE may be utilized in Fallow Cropping Systems for chemical weed control to aid in moisture conservation between cropping periods.

**Preplant Burndown**

Apply AIM EC HERBICIDE alone or with other herbicides or liquid fertilizers as a burn-down treatment to control or suppress weeds. AIM EC HERBICIDE is effective as a burndown treatment for crops prior to new plantings. **DO NOT** exceed the applicable amounts as listed for the specific crop in the MAXIMUM ALLOWABLE AIM EC HERBICIDE USE in Table 1. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. Coverage is essential for good control. Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a labeled burndown herbicides including glyphosate, glufosinate, paraquat, 2,4-D, or dicamba.

**Harvest Aid Desiccation Application**

Apply AIM EC HERBICIDE foliar to potatoes in the later stages of senescence for desiccation of potato foliage and vines. AIM EC HERBICIDE will also desiccate late season susceptible broadleaf weeds to aid in tuber harvest. Adequate desiccation is achieved within 14 days after the initial treatment is applied. If the potato crop is in the active vegetative growth stage when desiccation is initiated, two applications may be required to provide desiccation of leaf and stem tissue. Dense potato canopy, large plant size and environmental conditions not conducive to product absorption or activity will reduce initial application efficacy and increase the need for a second application. If a second application is necessary, apply at 7 to 14 days after the first application. Thorough coverage of the potato plant to be desiccated is essential. Use a sufficient volume of water to obtain thorough coverage of the potato leaves and vines.

**Ground Application**

Apply AIM EC HERBICIDE in at least 20 gallons of water per acre. Vary the spray volume and spray pressure as indicated by the density of the potato canopy and vines to assure thorough spray coverage. Increase the spray volume and pressure if the potato canopy is dense or under cool, cloudy or dry conditions. Increased spray volumes will enhance performance.

**Aerial Application**

Apply AIM EC HERBICIDE with aerial equipment using 5 to 10 gallons of water per acre, using higher volumes when potato canopies and vines are dense. Adjust the nozzles to provide a uniform pattern and a droplet size of 350 to 450 microns.

**Adjuvant Requirements**

A nonionic surfactant (NIS), methylated seed oil (MSO), or crop oil concentrate (COC) or other suitable surfactant mixture is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). The use of a high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

Adjuvant rates must increase as spray volumes exceed 20 gallons per acre.

**Tank Mixtures**

Apply AIM EC HERBICIDE as a tank mix or as a sequential application with other potato desiccants. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

## Conditions of Sale and Limitation of Warranty and Liability:

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or FMC, and Buyer assumes the risk of any such use.**

### DISCLAIMER

FMC Corporation intends that AIM EC HERBICIDE be distributed only to end users and/or growers (and/or applicators acting on the behalf of growers), who agree to the terms and conditions as stated herewith and further agree to a waiver and release from any and all liability by the user and/or grower of FMC for failure to perform and/or crop damage resulting from the use of AIM EC HERBICIDE as specified on the labeled crops under the those specific sections of this label. If such terms and conditions are unacceptable, FMC requests the return at once of all products in unopened original containers. FMC considers the user to have accepted such terms and conditions upon the use of AIM EC HERBICIDE.

AIM EC HERBICIDE, when used as directed, may result in crop injury, crop loss or crop damage. FMC directs that the user and/or grower test AIM EC HERBICIDE in order to determine its suitability for the intended use. FMC makes AIM EC HERBICIDE available to the user and/or grower solely to the extent that the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of AIM EC HERBICIDE. The decision to use, or not to use, AIM EC HERBICIDE must be made by each individual user and/or grower on the basis of possible crop injury from AIM EC HERBICIDE, the severity of weed infestations, the cost of alternative weed control measures and other factors. Because of the risk of crop damage, all such use is at the user and/or grower's risk.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC OR SELLER SHALL NOT BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OF OR HANDLING OF THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

**This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.**

NOTICE TO BUYER -- Purchase of this material does not confer any rights under patents of countries outside of the United States. FMC, the FMC logo, and AIM EC are trademarks of FMC Corporation and/or an affiliate.

## LABEL TRACKING INFORMATION

Label Code: SL-4868A 022124 2-28-24