

COMPACT®

ACTIVE INGREDIENT:

Mepiquat Chloride: N, N-dimethylpiperidinium Chloride	4.2%
OTHER INGREDIENTS:	95.8%
TOTAL	100.0%

Contains 0.35 pounds active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: No known antidote, treat symptomatically.

EMERGENCY CONTACT: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-424-7452 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin and clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal area below the high water mark. Do not contaminate water when disposing of rinsate or equipment washwater. Do not contaminate water by cleaning of equipment or disposal of wastes.

SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS, COMPLETE DIRECTIONS FOR USE, WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY.

NET CONTENTS: 2.5 GALLONS (9.46 L)

EPA Reg. No. 1381-189
EPA Est. No. 42750-MO-002

45901A50022

DISTRIBUTED BY:
WINFIELD SOLUTIONS, LLC
PO BOX 64589
ST. PAUL, MN 55164-0589

2/1212/2

Non-target Organism Advisory: This product may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by minimizing spray drift.

Runoff Prevention Advisory: To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not store below 32° F, nor above 100° F. Store in a dry place away from heat or open flame. Store in locked area in original container only, with lid tightly closed. Store separately from other pesticides and fertilizers, food and feed to prevent contamination. Use care to avoid puncturing container during storage or transit.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Use the instructions appropriate to the container size and type.

Nonrefillable container equal to or less than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable container greater than 5 gallons: Nonrefillable containers. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Refillable container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call
CHEMTREC 1-800-424-9300**

ITEM # 10120837

PULL HERE TO OPEN ▲

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin and clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal area below the high water mark. Do not contaminate water when disposing of rinsate or equipment washwater. Do not contaminate water by cleaning of equipment or disposal of wastes.

Non-target Organism Advisory: This product may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by minimizing spray drift.

Runoff Prevention Advisory: To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

PRODUCT INFORMATION

COMPACT is a foliar applied plant growth regulator. It allows the grower to manage the cotton plant for short-season production leading to reduced risk of yield and quality loss due to delayed and prolonged harvest. Benefits derived from the use of COMPACT include increased early boll retention and/or larger bolls, reduced plant height which provides a more open canopy, less boll rot, improved defoliation, less trash and lower ginning costs, better harvest efficiency and a darker leaf color. These benefits can provide for earlier maturity and often result in improved yields.

COMPACT contains a dye and effectiveness is not related to the color of the spray solution.

SPRA
Thoro

In Wa
Areas
Groun
Aerial
Califo
Groun
Aerial

In Oil:
Use a
TN an
in the
shoul
volum

Mix u
3% of
and d

See th

TANK
It is t
direct
in the

The u

COMP
used v
fertil.

Comp
Add c

1. We
2. sou
3. Pr
4. W
5. En
6. W
7. Let
8. Ev
- (cl

SPRA
Avoid





USE INFORMATION

SPRAY VOLUMES

Thorough coverage is required

In Water:

Areas other than California:

Ground Application - Use a minimum of 2 gallons of water per acre.

Aerial Application - Use a minimum of 2 gallons/acre.

California Only:

Ground Application - Use a minimum of 5 gallons/acre.

Aerial Application - Use a minimum of 5 gallons/acre.

In Oil:

Use a minimum total oil volume of 2 pints/acre for ultra-low volume (ULV) aerial application. Application in oil is permitted only in AL, AR, FL, GA, LA, MO, MS, NC, OK, SC, TN and TX. Use a nonphytotoxic oil concentrate which contains either a petroleum or vegetable oil base, contains only EPA-exempt ingredients, provides good mixing quality in the jar test (see "Compatibility Jar Test for Mix Components" in the "TANK MIXING" section of the label), and has been used successfully in your locality. The oil diluent should contain emulsifiers which provide good mixing quality. If the oil does not contain an emulsifier, one must be added during mixing at a volume equal to 3% of the final volume of the mixing tank. Do not apply COMPACT as ULV without using emulsifiers. If using a vegetable oil-based product, only highly refined concentrates should be used.

Mix under constant agitation. Pour one-half of the required volume of oil into the spray tank, followed by the emulsifier (if the oil does not already contain one) at approximately 3% of the final spray tank volume, and then pour in the COMPACT while the remainder of the oil is added. Constant, moderate agitation is required during and after mixing and during application.

See the "TANK MIXING" section or the label for additional information.

TANK MIXING

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

The use of a high quality, EPA exempt surfactant will enhance the uptake of COMPACT into the plant.

COMPACT is water based, and is compatible with most insecticides and miticides. If compatibility is in doubt, perform a jar test to check for compatibility. COMPACT can be used with foliar fertilizers if your prior experience shows the combination is compatible and will not injure cotton under your conditions. Use caution when applying with foliar fertilizers under conditions of extreme heat.

Compatibility Jar Test for Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

1. Water: For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
2. Products in PVA (polyvinyl alcohol) Bags: Cap the jar and invert 10 cycles.
3. Water-Dispersible Products (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions): Cap the jar and invert 10 cycles.
4. Water-Soluble Products (such as this product): Cap the jar and invert 10 cycles.
5. Emulsifiable Concentrates (oil concentrates): Cap the jar and invert 10 cycles.
6. Water-Soluble Additives: Cap the jar and invert 10 cycles.
7. Let the solution stand for 15 minutes.
8. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. Do not use any spray solution that could clog spray nozzles.

SPRAY DRIFT MANAGEMENT

Avoid spray drift. Do not allow this product to drift on to non-target areas. Where states have more stringent requirements they must be observed.

Mandato

Aerial Ap

- App
- Do
- Do
- Do
- The
- or 5
- Wh
- Ori

Ground A

- App
- Do
- Do
- Wh

Spray Drif

THE APPL

Importanc

An effectiv
drift, the p

Controlli

- Volu
- grea
- Pres
- Spra

Controlli

- Adju
- with

Boom Heig

For grounc

Release H

Higher rele

Shielded 5

Shielding t
of the spr

Temperatu

When mak

Temperatu

Drift poter
limited clo
smoke ger
and rapidh

Wind

Drift poter

separately from

clean clothing.

sing of rinsate or

the treated site.

waters. Applying
ent area. Rinsing

g application. For

r the protection
contamination,
:tive equipment

treated such as

d and quality loss
t which provides
provide for earlier





Mandatory Spray Drift

Aerial Applications:

- Applicators are required to use a medium to coarser droplet size, as defined by ASABE Standard S572.1.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.
- Do not release spray at a height greater than 10 feet above the crop canopy, unless a greater application height is necessary for pilot safety.
- The spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
- When applying to crops via aerial application equipment, use 1/2 swath displacement upwind at the edge of the field.
- Orient nozzles so the spray is directed toward the back of the aircraft.

Ground Applications:

- Applicators are required to use a medium to coarser droplet size, as defined by ASABE Standard S572.1.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.
- When using ground application equipment, apply with nozzle height no more than 3 feet above the ground or crop canopy.

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Importance Of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide thorough coverage. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a nozzle type that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

Boom Height- Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

Release Height - Aircraft

Higher release heights increase the potential for spray drift.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Wind

Drift potential generally increases with wind speed. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Use Restriction

- Do not apply to
- **Maximum Se**
 - Do not appl
 - The sum of
- **Maximum Sir**
- **Pre-harvest I**
- **Grazing/Feed**
- **Rotational Co**
- Do not tank r
- **Rotational Co**
- See the "SPR
- Do not apply

Use Limitation

- Insect or mite

Early Applicati

Growers may use Do not apply COI with insecticides

In the absence of and timing of si instructions.

Use higher rates COMPACT per ac

If significant loss COMPACT is inc

Late Season Ap

Late application substitute for ea season vegetati ginning costs. S or nutrient stres combination of

Timing for Late

- On fields wh but not alwa
- On fields wh mainstem n the last node bloom. Durin soon enough

Use Rate for La

Apply COMPACT the higher rate r

NC, OK, SC, mixing quality e oil diluent , of the final .ld be used. proximately after mixing

tations and ach product

FACT can be g with foliar

1e intended

m, nor thick



APPLICATION INSTRUCTIONS

SHORT-STAPLE AND LONG-STAPLE (PIMA) COTTON

Use Restrictions:

- Do not apply COMPACT to cotton that is under stress. If using low-rate multiple applications, discontinue use until your crop has overcome any stress.
- Maximum Seasonal Rate:**
 - Do not apply more than 48 fl. oz. (0.132 lb. a.i.) of COMPACT per acre per season.
 - The sum of all products and formulations containing mepiquat chloride must not exceed 0.132 pounds (60 grams) of mepiquat chloride per acre per season.
- Maximum Single Application Rate:** Do not exceed the rates in "TABLE I" and "TABLE II".
- Pre-harvest Interval (PHI):** Do not apply COMPACT within 30 days of harvest.
- Grazing/Feeding Restriction:** Do not graze or feed cotton forage to livestock.
- Do not tank mix with other products other than those mentioned under "TANK MIXING".
- Rotational Crop Restriction:** Do not plant another crop within 75 days after last treatment.
- See the "SPRAY DRIFT MANAGEMENT" section of this label for mandatory and advisory spray drift mitigation measures.
- Do not apply this product through any type of irrigation system.

Use Limitations:

- Insect or mite damage to COMPACT treated crops can lead to yield decreases or other undesirable effects.

Early Application of COMPACT

Growers may use low-rate multiple applications, or higher, less frequent dosages which provide maximum flexibility under a wide range of growing conditions. Carefully scout fields. Do not apply COMPACT to plants under stress. If stress is alleviated, evaluate plants for vegetative growth before additional applications are made. COMPACT may be tank mixed with insecticides/miticides when application timing coincides. See the "TANK MIXING" section of the label for further information.

In the absence of stress, a maximum of five low-rate applications can be made each season. The first application can be applied at the matchhead square stage. The rate and timing of subsequent applications depend on growing conditions and desired benefits. See "TABLE II: LOW-RATE MULTIPLE APPLICATIONS" for specific application instructions.

Use higher rates of COMPACT if vegetative growth becomes excessive or a greater degree of height control is desired. Do not use more than a total of 48 fl. oz. (0.132 lb. a.i.) of COMPACT per acre in a growing season. See "TABLE I: HIGH RATE SINGLE OR MULTIPLE APPLICATIONS" for specific application instructions.

If significant loss of squares and/or young bolls has occurred earlier due to insect pressure or other stresses, but now these stresses have been alleviated, the need for COMPACT is increased - excess vegetative growth is likely because of poor fruit load.

Late Season Application of COMPACT

Late application of COMPACT (approximately during the fourth to sixth week of blooming) can provide certain benefits to cotton. However, it should not and does not substitute for early season use, the time of the greatest benefit from the use of COMPACT. Late season application can lead to one or more of the following: reduction in late season vegetative growth or regrowth after cutout or defoliation, more complete and manageable cutout, better defoliation, earlier maturity and reduction in trash and lower ginning costs. Some of these effects may favorably influence the yield potential and fiber quality. Only apply a late season application of COMPACT if fields are not drought or nutrient stressed; that is, those fields likely to experience additional vegetative growth or regrowth. However, fields that are very rank and extremely vigorous due to a combination of poor boll load and excellent growing conditions may not respond as much as desired to late season applications at the suggested rates.

Timing for Late Season Applications

- On fields where cotton cuts out and then starts regrowth: apply when regrowth begins, as evidenced by new leaves in the terminal and stem elongation. This would often, but not always, be in the period of 5-6 weeks after the first bloom.
- On fields where cotton never completely cuts out, apply COMPACT when there are 4-6 nodes above the white flower (NAWF). Measure NAWF by counting the number of mainstem nodes from the first position white bloom (the one closest to the mainstem) to the terminal. Count the node with the first position white bloom as zero and the last node in the terminal, which is counted, should have a leaf at least the size of a quarter. Generally, the NAWF first reaches 4-6 during the fourth to sixth week of bloom. During this time period, the NAWF should be decreasing about one node every 5-6 days - if its rate of decrease is less, this means that the plant is not cutting out soon enough (the crop is too vigorous). If the fifth week of bloom arrives and NAWF is still above 5-6, apply COMPACT.

Use Rate for Late Season Application

Apply COMPACT at a rate between 8 to 24 fl. oz. (0.022 to 0.066 lb. a.i.) per acre. Use the lower rate range on cotton with only moderate additional growth potential, and the higher rate range on fields likely to continue vigorous growth. Total use per season (early plus late application) must not exceed 48 fl. oz. (0.132 lb. a.i.) per acre.



TABLE I. HIGH RATE SINGLE OR MULTIPLE APPLICATIONS:

Use these instructions when you are not able to start growth regulation treatments early, or when you want to make the fewest number of applications.

LOCATION	APPLICATION INSTRUCTIONS	APPLICATION RATE PER ACRE
AL, AR, AZ, CA FL, GA, LA, MO, MS, NM, NC, SC, TN, VA	First Application: Apply when cotton is actively growing and is between 20" and 30" tall, but not more than 7 days beyond early bloom (5-6 blooms per 25 row feet). Also, apply if cotton is 24" tall, and has no blooms. Use the 8 fl. oz. rate on the cotton where excessive vegetative growth is not expected. Use 16 fl. oz. where excessive growth has historically occurred. See the "Use Restrictions" and "Use Limitations" sections of the label.	8 to 16 fl. oz. (0.022 to 0.044 lb. a.i.)
	Second Application: Make another application in 2 to 3 weeks if additional growth control is desired.	8 to 16 fl. oz. (0.022 to 0.044 lb. a.i.)
	Third Application for Control of Excessive Vegetative Growth: If the cotton field has a history of vigorous growth or if conditions continue to favor vigorous growth, apply a third application 1 to 2 weeks after the second application.	8 to 16 fl. oz. (0.022 to 0.044 lb. a.i.)
	Late Season Application: See section titled "Late Season Application of COMPACT".	8 to 24 fl. oz. (0.022 to 0.066 lb. a.i.)
Areas <i>without</i> a history of excessive vegetative growth: OK, TX, (except Rio Grande Valley)	First Application: Apply when cotton is in the early bloom stage (5-6 blooms per 25 row feet) and actively growing. Also apply if no blooms are present and the cotton is 20" tall and actively growing. See the "Use Restrictions" and "Use Limitations" sections of the label.	8 fl. oz. (0.022 lb. a.i.)
	Second Application: Make a second application in 2 to 3 weeks if additional growth control is desired.	8 fl. oz. (0.022 lb. a.i.)
	Third Application: If conditions after the second application of COMPACT continue to favor vigorous growth, apply a third application 1 to 2 weeks after the second application.	8 fl. oz. (0.022 lb. a.i.)
	Late Season Application: See section titled "Late Season Application of COMPACT".	8 to 24 fl. oz. (0.022 to 0.066 lb. a.i.)
Areas <i>with</i> a history of excessive vegetative growth: OK, TX	First Application: For best results, apply when plants are in early bloom stage (5-6 blooms per 25 row feet) and an average of 24" tall. Treatments can also be made when cotton height averages a minimum of 20" and a maximum of 30", provided cotton is not more than 7 days beyond early bloom. If cotton is 24" tall and has no blooms, apply COMPACT. See the "Use Restrictions" and "Use Limitations" sections of the label.	16 fl. oz. (0.044 lb. a.i.)
	Second Application: For fields with a history of excessive growth, or if conditions after the first application favor excessive growth, make a second application in 2 to 3 weeks.	8 to 16 fl. oz. (0.022 to 0.044 lb. a.i.)
	Third Application: If conditions after the second application of COMPACT continue to favor vigorous growth, apply a third application 1 to 2 weeks after the second application.	8 to 16 fl. oz. (0.022 to 0.044 lb. a.i.)
	Late Season Application: See section titled "Late Season Application of COMPACT".	8 to 24 fl. oz. (0.022 to 0.066 lb. a.i.)

TABLE II.

Use these

LOC

AL, AR,
GA, LA,
NC, NM,
TN, TX, **Use the
Matchh
the planDo not c
STORAC
Store sePESTICI
disposer
Regiona

CONTAI
Nonrefi
rinse as
full with
after th
other pr
Nonrefi
follows:
its side
the cont
or dispo
approve
Refillab
respons
the rem
water w
offer fo

**TABLE II. LOW-RATE MULTIPLE APPLICATIONS:**

Use these instructions when you want to maintain maximum flexibility in plant regulation treatments.

LOCATION	APPLICATION INSTRUCTIONS	EXCESSIVE GROWTH NOT EXPECTED OR LOWER RATES HAVE WORKED IN THE PAST APPLICATION RATE PER ACRE	EXCESSIVE GROWTH EXPECTED OR HIGHER RATES HAVE BEEN NECESSARY IN THE PAST APPLICATION RATE PER ACRE
AL, AR, AZ, CA, FL, GA, LA, MO, MS, NC, NM, OK, SC, TN, TX, VA	First Application: Apply at the matchhead square ¹ stage of growth.	2 fl. oz. (0.0055 lb. a.i.)	4 fl. oz. (0.011 lb. a.i.)
	Second Application: 7-14 days later, or when regrowth occurs.	2 fl. oz. (0.0055 lb. a.i.)	4 fl. oz. (0.011 lb. a.i.)
	Third Application: 7-14 days later, or when regrowth occurs.	2 to 4 fl. oz.* (0.0055 to 0.011 lb. a.i.)	4 to 8 fl. oz.* (0.011 to 0.022 lb. a.i.)
	Fourth Application: 7-14 days later, or when regrowth occurs.	2 to 8 fl. oz.* (0.0055 to 0.022 lb. a.i.)	4 to 12 fl. oz.* (0.011 to 0.033 lb. a.i.)
	Fifth Application: (If needed): 7-14 days later, or when regrowth occurs.	4 to 8 fl. oz.* (0.011 to 0.022 lb. a.i.)	4 to 12 fl. oz.* (0.011 to 0.033 lb. a.i.)
	Late Season Application: See section titled "Late Season Application of COMPACT".	8 to 16 fl. oz.* (0.022 to 0.044 lb. a.i.)	12 to 24 fl. oz.* (0.033 to 0.066 lb. a.i.)

*Use the higher rate if previous application was not made or if growing conditions favor excessive growth.

¹Matchhead square is when the first square of a typical cotton plant is about the size of a match head (about 1/8" in diameter). Make the first application when 50% of the plants have one or more matchhead squares.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not store below 32° F, nor above 100° F. Store in a dry place away from heat or open flame. Store in locked area in original container only, with lid tightly closed. Store separately from other pesticides and fertilizers, food and feed to prevent contamination. Use care to avoid puncturing container during storage or transit.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Use the instructions appropriate to the container size and type.

Nonrefillable container equal to or less than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable container greater than 5 gallons: Nonrefillable containers. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Refillable container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call
CHEMTREC 1-800-424-9300**

The directions FURNISHED " ANY KIND TO INCLUDING, B USAGE. UNIN ABSENCE OF ARE BEYOND AGREEMENTS REPRESENTA

TO THE EXTEN DAMAGES, OI MANUFACTUR ANY DAMAGE FOLLOW LABI THE BUYER, U NOT EXCEEDII

If you do not a the Seller, and Compact" is a



RE
OF
LD,
DE
OR
CH
ND
NY

IAL
AL
OR
TO
BY
IES

l to



COMPACT®

Plant Growth Regulator

ACTIVE INGREDIENT:

Mepiquat Chloride: N, N-dimethylpiperidinium Chloride 4.2%

OTHER INGREDIENTS: 95.8%

TOTAL 100.0%

Contains 0.35 pounds active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: No known antidote, treat symptomatically.

EMERGENCY CONTACT: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-424-7452 for emergency medical treatment information.

**SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS, COMPLETE DIRECTIONS FOR USE,
WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY.**

NET CONTENTS: 2.5 GALLONS (9.46 L)

EPA Reg. No. 1381-189
EPA Est. No. 42750-MO-002

45901A50022

DISTRIBUTED BY:
WINFIELD SOLUTIONS, LLC
PO BOX 64589
ST. PAUL, MN 55164-0589

2/1212/2



(01)00884464496728

ITEM # 10120837

PROOF

THIS PROOF IS TO BE CHECKED FOR ACCURACY

Please review and approve **Text, Spelling, Copy Placement, Size, Shape, Colors** and **Dieline**.

Authorized signature accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.

PLEASE NOTE: Due to color variance between printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors.

THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH.

Dieline does not print.



4201 North Westport Ave. • Sioux Falls, SD 57107
Phone: (605) 978-0451 • Fax: (605) 978-0463

DATE	JOB NUMBER	CUSTOMER
10-16-23	198563	Albaugh
LABEL SIZE	LEAFLET FOLDED SIZE	LEAFLET FLAT SIZE
6.75" x 6.75"	6.5" x 6"	6.5" x 23.75"
LABEL COLORS	LEAFLET "OUT" COLORS	LEAFLET "IN" COLORS
 BLK	 BLK  576	 BLK
PATTERN VARNISH: <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO		

Form: CS 006L - 11/8/2011

ARTWORK IS APPROVED

REVISED PROOF NEEDED

**WE CANNOT PROCESS
THIS ORDER WITHOUT AN
AUTHORIZED SIGNATURE**

Signed Lisa Pothan

Date 10/18/2023