THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



2(ee) Recommendation

FOR USE AND DISTRIBUTION IN THE STATE OF NEW MEXICO ONLY.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

This Product Bulletin Contains New or Additional Directions For Use Which Are Recommended By FMC Corporation Under FIFRA Section 2(ee)

50 WG Insecticide

EPA REG. NO. 71512-9-279

CROP	PEST	RATE OF APPLICATION
Pecan	Yellow Pecan Aphid Black Pecan Aphid Blackmargined Aphid	2.0-2.8 oz/A (0.062-0.089 lb ai/A)

DIRECTIONS FOR USE:

Begin applications before populations begin to build and before damage is evident according to local pest management guidelines. Use low rate for building populations and high rate for greater populations and/or dense foliage. Scout trees often and retreat as necessary to maintain populations below damaging levels. If identification of aphid species has not been confirmed use high rate.

RESISTANCE MANAGEMENT:

Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of Carbine 50WG Insecticide in programs that seek to minimize the occurrence of pest resistance.

RESTRICTIONS:

- Do not apply more than 2.8 oz/A of product per application.
- Do not apply more than 8.4 oz/A of product per season.
- Do not apply more than 3 applications per season. Allow a minimum of 7 days between applications.

PHI: Do not apply within 40 days of harvest.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED CARBINE® 50WG INSECTICIDE (EPA REG. NO. 71512-9-279) LABEL MUST BE FOLLOWED. THIS RECCOMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

