

RESTRICTED USE PESTICIDE

Due to Toxicity to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.



EPA REG. No. 279-3313

FIFRA Section 24(c)
SPECIAL LOCAL NEED LABEL FOR USE ON
ALFALFA AND CLOVER GROWN FOR SEED.

FOR DISTRIBUTION AND USE ONLY IN
WITHIN THE STATE OF WASHINGTON

EPA SLN No. WA -070015

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

THIS LABEL FOR BRIGADE 2EC INSECTICIDE/MITICIDE EXPIRES AND MUST NOT BE DISTRIBUTED OR USED IN ACCORDANCE WITH THIS SLN REGISTRATION AFTER DECEMBER 31, 2025.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

Crop	Pest	Rate of Application
Alfalfa Grown for Seed Clover Grown for Seed	Aphids, Lygus Bugs, Spider Mites, Weevils	6.4 fl. oz./A

Directions for Use:

Apply Brigade 2EC Insecticide/Miticide at the rate 6.4 fluid ounces (0.1 lb. active ingredient) per acre. Apply in a minimum of 5 gallons of water per acre by air equipment or in a minimum of 20 gallons of water by ground equipment.
Apply as a pre-bloom or post-bloom spray only.

Restrictions / Precautions:

For use in accordance with this SLN label do not apply this product through any type of irrigation system. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Do not apply more than 6.4 fluid ounces (0.1 lb. active ingredient) per acre per application. Do not make more than 2 applications of bifenthrin (all products) per season. Maintain a minimum of 21 days between applications. Rotational crops may be planted no sooner than 30 days after last application. Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Only apply products containing bifenthrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

WSDA Aquatic Advisory: This pesticide is extremely toxic to fish and aquatic invertebrates. Brigade 2EC Insecticide/Miticide should not be used in accordance with this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Natural Resources Assessment Section web site (<https://agr.wa.gov/departments/land-and-water/natural-resources/endangered-species-program>) for additional information related to pesticide use and endangered species protection. Information from the EPA about no-spray buffer zones is available at <https://agr.wa.gov/departments/land-and-water/natural-resources/endangered-species-program/buffers>. Pesticide applicators may use the Washington Department of Fish and Wildlife's SalmonScape mapping tool at <https://apps.wdfw.wa.gov/salmonscape> to determine if listed salmonid species occur in or near a proposed application site. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

All alfalfa/clover seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amounts of material disposed of, the grower(s), and the date of disposal.

No portion of the alfalfa/clover seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.

All alfalfa/clover seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.

Alfalfa/clover seed may not be distributed for human consumption or animal feed.

Pollinator Protection: This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area: apply as a pre-bloom or post-bloom spray only.

WSDA Container Disposal Guidance:

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <https://agr.wa.gov/departments/pesticides-and-fertilizers/pesticides/waste-pesticide>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.