

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

BIFENTHRIN | GROUP 3A | INSECTICIDE

Supplemental Labeling

This supplemental label expires on December 31, 2024 and must not be used or distributed after this date.



For use on Peach, Pome Fruit and Pomegranate

EPA Reg. No. 279-3313

Active Ingredient:	By Wt.
Bifenthrin: *	25.1%
Other Ingredients:	74.9%
	100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum

**Contains xylene range aromatic solvents.

This product contains 2 pounds active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER
INCONSISTENT WITH ITS LABELING.
ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA
REGISTERED LABEL ARE TO BE FOLLOWED.**

This Supplemental labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for BRIGADE 2EC Insecticide/Miticide before applying. Use of BRIGADE 2EC Insecticide/Miticide according to this supplemental labeling is subject to the use precautions and limitations imposed by the label affixed to the container of BRIGADE 2EC Insecticide/Miticide. Carefully follow all precautionary statements and application use directions.

PEACH Subgroup 12-12B

PEST	DOSAGE		REMARKS
	lb ai/A	fl oz/A	
Aphids Codling Moth Cutworms Green Fruitworm Leafhoppers Leafminers Leafrollers Lygus spp. Plant Bugs Plum Curculio San Jose Scale (Crawlers) Stink Bugs Tarnished Plant Bugs	0.04 - 0.2	2.6 - 12.8	<p>Application by ground: Apply as a dilute (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre) spray in sufficient water to provide thorough coverage.</p> <p>Application by air: Apply the specified dosage in a minimum of 10 gallons of finished spray per acre by air.</p>
Two spotted Spider Mite Yellow Mite	0.06 - 0.2	3.8 - 12.8	
European Red Mite	0.08 - 0.2	5.12 - 12.8	
<p>RESTRICTIONS</p> <ul style="list-style-type: none"> Do not apply more than 0.2 lb ai/A (12.8 fl oz/A) per application. Do not apply more than 0.5 lb ai/A (32 fl oz/A) per year with no more than 0.45 lb ai/A (28.8 fl oz/A) applied after petal fall. Do not make more than 3 applications per year. Do not make applications less than 30 days apart. Do not apply within 14 days of harvest. Do not graze livestock in treated orchards or cut treated cover crops for feed. 			

POME FRUITS (except Mayhaw)

Apple; azarole; crabapple; loquat; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote

PEST	DOSAGE		REMARKS
	lb ai/A	fl oz/A	
Aphids Codling Moth Cutworms Green Fruitworm Leafhoppers Leafminers Leafrollers Lygus spp. Plant Bugs Plum Curculio San Jose Scale (Crawlers) Stink Bugs Tarnished Plant Bugs	0.04 - 0.2	2.6 - 12.8	<p>Application by ground: Apply as a dilute (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre) spray in sufficient water to provide thorough coverage.</p> <p>Application by air: Apply the specified dosage in a minimum of 10 gallons of finished spray per acre by air.</p>
Two spotted Spider Mite Yellow Mite	0.06 - 0.2	3.8 - 12.8	
European Red Mite	0.08 - 0.2	5.12 - 12.8	
<p>RESTRICTIONS</p> <ul style="list-style-type: none"> Do not apply more than 0.2 lb ai/A (12.8 fl oz/A) per application. Do not apply more than 0.5 lb ai/A (32 fl oz/A) per year with no more than 0.45 lb ai/A (28.8 fl oz/A) applied after petal fall. Do not make more than 3 applications per year. Do not make applications less than 30 days apart. Do not apply within 14 days of harvest. Do not graze livestock in treated orchards or cut treated cover crops for feed. 			

POMEGRANATE

PEST	DOSAGE		REMARKS
	lb ai/A	fl oz/A	
Katydid Navel Orangeworm Omnivorous Leafrollers Leaf footed Plant Bugs Fuller Rose Beetles Aphids White scales Ground Beetles Brown Marmorated Stink Bugs	0.1 – 0.2	6.4 – 12.8	Apply foliar treatments in at least 50 gallons of finished spray per acre.
RESTRICTIONS <ul style="list-style-type: none"> • Do not apply more than 0.2 lb ai/A (12.8 fl oz/A) per application. • Do not apply more than 0.5 lb ai/A (32 fl oz/A) per year. • Do not make more than 3 applications per year. • Do not make applications less than 14 days apart. • Do not apply within 14 days of harvest. 			

FMC and BRIGADE are trademarks of FMC Corporation.
 © 2022 FMC Corporation. All rights reserved.

R-4649 101521 11-19-21