

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3313

**FOR DISTRIBUTION AND USE IN
THE STATE OF TEXAS ONLY**

**THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS
MADE AS PERMITTED UNDER FIFRA
SECTION 2(ee) AND HAS NOT BEEN
SUBMITTED TO OR APPROVED BY THE EPA.**

This label for BRIGADE 2EC Insecticide/Miticide is valid until 11/20/2024, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR CONTROL OF BROWN MARMORATED STINK BUGS

CROP	RATE OF APPLICATION	RESTRICTIONS
Artichokes	6.4 fl oz/A (0.1 lb ai/A)	<ul style="list-style-type: none">Do not make applications less than 15 days apart.Do not apply more than 32 fl oz/A of product (0.5 lb ai/A) per season.Do not apply within 5 days of harvest.
Bushberries	6.4 fl oz/A (0.1 lb ai/A)	<ul style="list-style-type: none">Do not make applications less than 7 days apart.Do not apply more than 32 fl oz/A of product (0.5 lb ai/A) per season.Do not apply within 1 day of harvest.
Caneberries	6.4 fl oz/A (0.1 lb ai/A)	<ul style="list-style-type: none">Do not apply more than 12.8 fl oz/A of product (0.2 lb ai/A) per season.Do not apply within 3 days of harvest.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

CROP	RATE OF APPLICATION	RESTRICTIONS
Cilantro, Coriander	6.4 fl oz/A (0.1 lb ai/A)	<ul style="list-style-type: none"> • Do not make applications less than 7 days apart. • Do not apply more than 32 fl oz/A of product (0.5 lb ai/A) per season. • Do not apply within 3 days of harvest.
Grapes	6.4 fl oz/A (0.1 lb ai/A)	<ul style="list-style-type: none"> • Do not apply more than 6.4 fl oz/A of product (0.1 lb ai/A) per season. • Do not apply within 30 days of harvest.
Hops	6.4 fl oz/A (0.1 lb ai/A)	<ul style="list-style-type: none"> • Do not make applications less than 21 days apart. • Do not apply more than 19.2 fl oz/A of product (0.3 lb ai/A) per season. • Do not apply within 14 days of harvest. • Use of ultra-low volume (ULV) application on hops is prohibited
Mayhaw	6.4 fl oz/A (0.1 lb ai/A)	<ul style="list-style-type: none"> • Do not make applications less than 7 days apart. • Do not apply more than 12.8 fl oz/A of product (0.2 lb ai/A) per season. • Do not apply within 30 days of harvest.
Root Crops (except sugarbeet)	6.4 fl oz/A (0.1 lb ai/A)	<ul style="list-style-type: none"> • Do not make applications less than 7 days apart. • Do not apply more than 32 fl oz/A of product (0.5 lb ai/A) per season. • Do not apply within 21 days of harvest.
Spinach	6.4 fl oz/A (0.1 lb ai/A)	<ul style="list-style-type: none"> • Do not make applications less than 7 days apart. • Do not apply more than 25.6 fl oz/A of product (0.4 lb ai/A) per season. • Do not apply within 40 days of harvest.
Tuberous and Corm Vegetables	6.4 fl oz/A (0.1 lb ai/A)	<ul style="list-style-type: none"> • Do not make more than 2 foliar applications per season. • Do not make applications less than 21 days apart. • Do not apply more than 32 fl oz/A of product (0.5 lb ai/A) per season, including soil applications. • Do not apply within 21 days of harvest.



RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

CROP	RATE OF APPLICATION	RESTRICTIONS
Conifer Seed Orchards	6.4 – 12.8 fl oz/A (0.1 – 0.2 lb ai/A)	<ul style="list-style-type: none">• Begin application 7 days following peak pollen flight and repeat at 30-day intervals to a maximum of 0.6 lb ai/A per season.• Do not make more than 6 applications per season or apply more than 38.4 fl oz/A of product (0.6 lb ai/A) per season.• Do not make applications less than 30 days apart

DIRECTIONS FOR USE:

Apply BRIGADE 2EC Insecticide/Miticicide at crop specific rate by ground or aerial equipment using sufficient spray volume to obtain thorough and uniform coverage of the target plant and foliage. Use a minimum of 2 gallons of finished spray per acre by air or 10 gallons of finished spray per acre by ground. The addition of 1 – 2 quarts of emulsified oil per acre to the spray solution may improve spray deposition and insect control. Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

Repeat applications of BRIGADE 2EC Insecticide/Miticicide as necessary to maintain control and prevent re-infestation from migrating adults into the crop. Brown marmorated stink bugs are very mobile and may re-infest crops quickly. Consult the label for maximum application rates, number and retreatment intervals. If another application is needed prior to the minimum retreatment interval, use another insecticide to maintain control.

