



EPA Reg. No. 71512-10-279

FIFRA 2(ee) Recommendation

**FOR DISTRIBUTION AND USE
ONLY IN THE FOLLOWING STATES:
CALIFORNIA and ARIZONA**

**THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS MADE AS
PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT
BEEN SUBMITTED TO OR APPROVED BY THE US EPA.**

**This recommendation for BELEAF[®] 50SG Insecticide is valid until January 30, 2028,
or until withdrawn, canceled or suspended.**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS
LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA
REGISTERED LABEL MUST BE FOLLOWED SUBJECT TO THE TERMS AND CONDITIONS OF THE
PRODUCT LABELING. THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE
TIME OF PESTICIDE APPLICATION.**

**FOR THE SUPPRESSION OF BEET LEAFHOPPER IN FRUITING VEGETABLES/TOMATO,
GROUND, AIR, and DRIP IRRIGATION**

Directions For Use

CROP	PEST	RATE	APPLICATION DIRECTIONS	REI
Tomato	Beet Leafhopper	4.28 oz/a (0.133 lbs ai/a)	GROUND and AERIAL: Apply as required by scouting. Begin applications before population begins to build and before damage is evident, according to local pest management guidelines.	12 hrs.
			DRIP: BELEAF 50SG Insecticide must be applied to ensure the product is in the root zone to provide effective control of target pests. Most effective application is when it is applied so that the roots are at or near the site of application. The earlier flonicamid is available to a developing plant, the earlier the protection begins. For transplanted crops, apply BELEAF 50SG Insecticide within 21 days of transplanting.	

RESTRICTIONS:

- Do not apply more than 4.28 oz/acre of BELEAF 50SG Insecticide (0.133 lbs. ai/acre) per application.
- Do not apply more than 8.4 oz/acre of BELEAF 50SG Insecticide (0.263 lbs. ai/acre) per year.
- Make no more than 2 drip applications per crop season.
- Do not apply more than one drip irrigation application at the 4.28 oz/acre (0.133 lbs. ai/acre) rate per season or in a single application.
- Allow a minimum of 7 days between applications.
- This product is moderately toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

R-4774 030823 02-01-2023

Beleaf[®] is a registered trademark of Ishihara Sangyo Kaisha, Ltd.
FMC and the FMC logo are trademarks of FMC Corporation
and/or an affiliate

© 2023 FMC Corporation. All rights reserved.