



EPA Reg. No. 71512-10-279

FIFRA SECTION 24(c) Special Local Need Label (SLN)
For drip irrigation use on leafy greens (crop subgroup 4-16A) for managing Thrips to reduce instances of Impatiens Necrotic Spot Virus (INSV) and Aphids
EPA SLN No. CA-230002

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA

Directions for Use: Drip Irrigation

- **It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**
- This state-specific Section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA registered label for Beleaf® 50SG Insecticide, EPA Reg. No. 71512-10-279 and this label.

Crop	Pest	Rate of Application
Leafy Greens crop subgroup 4-16A*	Aphids Thrips	2.8 to 4.28 oz per acre

*Amaranth, Chinese; amaranth, leafy; aster, Indian; blackjack; cat's whiskers; cham-chwi; cham-na-mul; chervil, fresh leaves; chipilin; chrysanthemum, garland; cilantro, fresh leaves; corn salad; cosmos; dandelion, leaves; dang-gwi, leaves; dillweed; dock; dol-nam-mul; ebolo; endive; escarole; fameflower; feather cockscomb; good king henry; huauzontle; jute, leaves; lettuce, bitter; lettuce, head; lettuce, leaf; orach; parsley, fresh leaves; plantain, buckhorn; primrose, English; purslane, garden; purslane, winter; radicchio; spinach; spinach, Malabar; spinach, New Zealand; spinach, tanier; Swiss chard; violet, Chinese, leaves; cultivars, varieties, and hybrids of these commodities

Specific Directions for Use:

- Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines.
- Use LOWER RATE for building populations and use HIGHER RATE for greater populations.
- Scout fields often and retreat as necessary to maintain populations below damaging levels. Rapidly growing plants may need retreatment.
- Beleaf® 50SG Insecticide must be applied to ensure the product is in the root zone to provide effective management of target pests.
- Most effective application is when it is applied so that the roots are at or near the site of application. The earlier flonicamid is available to a developing plant, the earlier the protection begins. For best results on transplanted crops, apply Beleaf® 50SG Insecticide within 28 days of transplanting.

Specific Use Restrictions:

- Make no more than 2 drip applications per crop season.
- Do not apply more than one drip irrigation application at the 4.28 oz/acre rate (0.133 lbs. ai/acre) or two drip applications at the 2.8 oz/acre rate (0.088 lbs. ai/acre) per crop season.
- Allow a minimum of 7 days between applications.
- When applying via drip irrigation, do not make foliar applications of flonicamid to the same crop during the same crop season.
- Do not apply within 14 days of harvest.

FMC
 FMC Corporation
 2929 Walnut Street
 Philadelphia, PA. 19104

Beleaf® is a trademark of Ishihara Sangyo Kaisha, Ltd
 FMC and the FMC logo are trademarks of FMC
 Corporation and/or an affiliate

R-4794 062623 03-30-23