

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3315

FOR DISTRIBUTION AND USE ONLY IN
DE, ME, MD, MA, MI, NJ, NC, OH, PA,
SC, VA, WV

THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS MADE
AS PERMITTED UNDER FIFRA SECTION 2(ee)
AND HAS NOT BEEN SUBMITTED TO OR
APPROVED BY THE EPA.

This recommendation for HERO Insecticide is valid until 03/25/25, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CROP	PEST	RATE OF APPLICATION
Field Corn (Grain and Silage) Popcorn Field Corn Grown for seed	Brown Marmorated Stinkbug	7.1 to 10.3 oz/A

DIRECTIONS FOR USE:

Apply 7.1 to 10.3 oz/A HERO insecticide by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2-5 gallons by air). When applying by air, 1-2 quarts of emulsifiable oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Apply at a minimum 3 to 5 days intervals or as needed for control under heavy insect pressure. Under heavy outbreak conditions tank mixing with another insecticide or use of HERO insecticide in a sequential insecticide program with an insecticide labeled for this insect pest is recommended. HERO insecticide may be used in a sequential insecticide program with STALLION insecticide where sequential applications may be warranted.

RESTRICTIONS:

- Do not apply more than 10.3 fl oz/A or 0.10 lb ai/A per application.
- Do not apply more than 41.2 fl oz/A or 0.40 lb ai/A per season.
- Do not apply within 30 days of harvest for grain and stover and 60 days for forage.
- Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application.
- Use of ultra-low volume (ULV) application on corn is prohibited.
- Do not make aerial or ground applications to corn if heavy rainfall is imminent.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000