

# LO-VOL 6

## 2,4-D LOW VOLATILE HERBICIDE

**TENKÖZ**<sup>®</sup>  
SERVING AGRICULTURE

**WEED AND BRUSH CONTROL IN SMALL GRAINS, FIELD CORN,  
RANGELAND, PASTURES, ROADSIDES AND FENCEROWS**

**ACTIVE INGREDIENT:**

2,4-Dichlorophenoxyacetic acid, isooctyl ester\* ..... 87.3%

**OTHER INGREDIENTS:** ..... 12.7%

**TOTAL:** ..... 100.0%

Isomer Specific AOAC Method, Equivalent to:

2,4-Dichlorophenoxyacetic Acid ..... 57.9%, 5.5 lbs./gal.

**EPA REG. NO. 71368-11-55467**

**EPA EST. NO. 228-IL-1**

## KEEP OUT OF REACH OF CHILDREN CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300. For Medical Emergencies Only, Call 877-325-1840.

### FIRST AID

<b>IF IN EYES</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

### NOTE TO PHYSICIAN

No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred. This product contains petroleum distillates. If large amounts, greater than 1 mg/kg body weight have been ingested, the stomach should be evacuated by gastric intubation with the aid of a cuffed endotracheal tube to prevent aspiration of petroleum distillates. After removal of stomach contents, wash stomach by instilling 30 to 50 grams of activated charcoal in 3 to 4 ounces of water through the stomach tube and again remove stomach contents. Avoid oily laxatives.

Manufactured For

Tenköz Inc., 100 N. Point Center E. • Suite 330, Alpharetta, GA 30022

Net Contents: 2.5 Gals. (3.78 L)

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Do not get in eyes, on skin or on clothing.

### PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

### All mixers, loaders, flaggers, and other applicators and handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes and socks, plus
- Chemical-resistant gloves (except pilots) and
- Chemical-resistant apron when mixing, loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Control Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE (personal protective equipment) may be reduced or modified as specified in the WPS. Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40CFR 170.240 (d)(6)].

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

### Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

### Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

### Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

### Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

### Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

## DIRECTIONS FOR USE

It Is A Violation Of Federal Law To Use This Product In A Manner Inconsistent With Its Labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any water-proof material, and shoes plus socks.

### GENERAL CAUTIONS AND RESTRICTIONS

Do not apply this product through any type of irrigation system.

Do not use in or near a greenhouse.

### MIXING INSTRUCTIONS

Add 1/2 the required amount of water to the spray tank, then add this product with agitation, and finally, the balance of the water with continued agitation. This material forms an emulsion in water, not a solution. This tends to separate on standing. Provide agitation to prevent such separation and insure uniform spray mixtures.

### COMPATIBILITY

If this herbicide is to be tank mixed with fertilizers or with other pesticides, compatibility should be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing.

Read and follow the label of each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

### COMBINATION WITH LIQUID NITROGEN FERTILIZER

Use 1/3 to 1/2 pint of this product per acre for weeding and feeding corn, small grains or grass pastures as directed on this label.

Use fertilizer at rates recommended by supplier or Extension Service Specialist.

Fill the spray tank about half full with the liquid fertilizer, then add this product with vigorous agitation, and complete filling the tank with fertilizer. Apply immediately and continue agitation in the spray tank during application. Application during very cold weather (near freezing) is not advisable. Do not allow mixture to stand overnight. Incompatibility may be encountered with some fertilizer brands or under some environmental conditions. If in doubt, test a small sample in the dilution ratio planned for application.

**NOTE:** Fertilizers can increase foliage contact burn of herbicides. Reducing the fertilizer rate and concentration will reduce the hazard of leaf burn.

### GENERAL INFORMATION

This product contains isooctyl ester of 2,4-D, the original and one of the best low volatile esters. In cropland, this herbicide is more effective than amines for controlling hard-to-kill weeds such as bindweed, thistle, smartweeds, wild garlic, curled dock, tansy ragwort and wild onions.

**INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT.**

Crop varieties vary in response to 2,4-D and some are easily injured. Apply this herbicide to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult your seed company, state Agricultural Extension Service or qualified crop consultant for advice.

## GENERAL WEED LIST

### ANNUAL AND BIENNIAL WEEDS

annual yellow sweetclover  
 \*beggarticks  
 bull thistle  
 coffeeweed  
 \*common broomweed  
 common burdock  
 common cocklebur  
 common evening primrose  
 common lambsquarters  
 croton (Texas or woolly)  
 hairy galinsoga  
 jimsonweed

\*knotweed  
 \*mallow (venice, dwarf, little)  
 marshelder  
 morning glory (common, ivy, woolly)  
 mustards (except blue mustard)  
 pepperweeds (except perennial)  
 \*\*pigweeds (Amaranthus spp.)  
 \*prickly lettuce  
 ragweed (common, giant)  
 rough fleabane  
 \*Russian thistle

salsify (western or common)  
 \*smartweeds (annual species)  
 sowthistle (annual or spiny)  
 sunflower  
 velvetleaf  
 \*vervains  
 vetches  
 \*wild carrot  
 wild lettuce  
 wild parsnips  
 wild radish

## GENERAL WEED LIST

### PERENNIAL WEEDS

\*alfalfa  
 \*bindweed (hedge, field & European)  
 blue lettuce  
 \*Canada thistle  
 catnip  
 chicory  
 dandelion  
 \*docks

\*dogbanes  
 \*goldenrod  
 \*ground ivy  
 \*hawkweed (orange)  
 healall  
 \*hoary cress  
 Jerusalem - artichoke  
 \*many-flowered aster

\*nettles (including stinging)  
 plantains  
 sowthistle (perennial)  
 \*tansy ragwort  
 \*vervains  
 \*western ironweed  
 \*wild garlic  
 \*wild onion

\*These species may require repeat applications and/or use of the higher rate recommended on this product label even under ideal conditions for applications.

\*\*Control of pigweeds in the High Plains area of Texas and Oklahoma may not be satisfactory with this product.

## SPECIFIC USE DIRECTIONS

### CEREAL GRAINS: WHEAT, BARLEY AND RYE (not underseeded with legumes)

APPLICATION TIMING	AMOUNT OF TENKOZ LO-VOL 6 PER ACRE	DIRECTIONS
<b>Postemergence</b> Annual and biennial weeds Perennial broadleaf weeds	1/3 to 1-1/3 pints* 2/3 to 1-1/3 pints*	Apply after grain is well tillered (usually about 4 to 8 inches high) but before boot. <b>Do not spray grain in boot to dough stage.</b>
Wild onion or garlic	1 to 1-1/3 pints*	Apply 1 pint when grain is well tillered and wild garlic or onion plants are small. Apply 1-1/3 pints after harvest in the crop stubble. For control of new fall growth of wild onion or garlic, refer to FALLOWLAND use directions.
<b>Emergency Weed Control in Wheat</b> Perennial broadleaf weeds	1.75 pints*	Apply when weeds are approaching bud stage, <b>but do not spray grain during the boot to dough stage.</b> The 1.75 pints per acre application can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury.
<b>Preharvest</b>	2/3 to 3/4 pint*	Apply when grains are in the hard dough stage to suppress large weeds that may interfere with harvest. Best results will be obtained when soil moisture is sufficient to induce succulent weed growth.

\*Use the lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control problem justifies the risk of grain damage. Do not apply to grain in the seedling stage.

**Note:** Maximum rate 1 pint per acre from jointing to before boot stage. Treatment at this growth stage (jointing to before boot) should be used where such increased risk of injury to crop is acceptable. Higher rates listed on this label are more likely to cause crop injury.

## RESTRICTIONS AND LIMITATIONS FOR USE ON CEREAL GRAINS

- For aerial application on grain: Apply this product in 2 or more gallons of water per acre.
- For ground application: A minimum of 10 to 15 gallons of water per acre is recommended for proper spray coverage.
- Do not mix with oil for crop uses.
- Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 2 weeks after treatment.
- Do not feed treated straw to livestock if a preharvest treatment or emergency treatment as described above is applied.
- Limited to 1.75 lbs ae (2.5 pints of product) per crop cycle.

### **Postemergence:**

Limited to one postemergence application per crop cycle. Maximum 1.75 pints per acre per application.

### **Preharvest:**

The preharvest interval (PHI) is 14 days.

Limited to one preharvest application per crop cycle.

Maximum of 3/4 pint per acre per application.

## FIELD CORN

APPLICATION TIMING	AMOUNT OF TENKOZ LO-VOL 6 EC PER ACRE	DIRECTIONS
<b>Preplant</b>	2/3 to 1-1/3 pints	To control emerged broadleaf weed seedlings or existing cover crops prior to planting corn, apply 7 to 14 days before planting. Do not use on light, sandy soil, or where soil moisture is inadequate for normal weed growth. Use high rate for control of less susceptible weeds or cover crops such as alfalfa.
<b>Preemergence</b>	1-1/3 to 1-1/2 pints	To control broadleaf weeds and suppress annual grasses, apply 3 to 5 days after planting, but before corn emerges. Use high rate on soil high in organic matter. Do not use on light, sandy soil, or where soil moisture is inadequate for normal weed growth.
<b>Postemergence</b> Annual broadleaf weeds Perennial broadleaf weeds	1/3 pint 1/2 pint	Avoid spraying just after corn leaves unfold as injury may occur. Apply when weeds are small and corn is less than 8 inches tall. If corn more than 8 inches tall must be sprayed, use drop pipes.  Spray when weeds are in the bud to bloom stage. If corn is more than 8 inches tall, use drop pipes to keep spray off corn leaves. Do not spray corn from tassel to dough stage. 2,4-D may make corn brittle. Winds or cultivation may cause stalk breakage while brittle.
<b>Preharvest</b>	2/3 to 1-1/3 pints	After the hard dough (denting) stage, to suppress weeds that interfere with harvest such as bindweed, cocklebur, dogbane, jimsonweed, ragweed, sunflower and velvetleaf, and to decrease production of weed seeds, spray with air or ground equipment. The high rate will be needed for tough weeds under stress.

## RESTRICTIONS AND LIMITATIONS FOR USE ON FIELD CORN

Do not use treated crop as fodder for 7 days following application.

Maximum of 4.25 pints per acre per crop cycle.

The preharvest interval (PHI) is 7 days.

### **Preplant or preemergence:**

Limited to one preplant or preemergence application per crop cycle.

Maximum of 1.5 pints per acre per application.

### **Postemergence:**

Limited to one postemergence application per crop cycle.

Maximum of 3/4 pint per acre per application.

### **Preharvest:**

Limited to one preharvest application per crop cycle.

Maximum of 2.25 pints per acre per application.

## GRAIN SORGHUM (milo)

APPLICATION TIMING	AMOUNT OF TENKOZ LO-VOL 6 PER ACRE	DIRECTIONS
Postemergence	1/2 to 3/4 pint	To control emerged annual and perennial broadleaf weeds when the sorghum is 6 to 15 inches tall (to top of canopy). If sorghum is taller than 8 inches to top of canopy, use drop nozzles to keep spray off crop foliage. <b>Do not treat during boot, flowering, or early dough stages.</b>

### RESTRICTIONS AND LIMITATIONS FOR USE ON SORGHUM

- A minimum application volume of 5 gallons per acre by air or 10 gallons per acre by ground is recommended.
- The preharvest interval (PHI) is 30 days.
- Do not permit meat or dairy animals to consume treated crop as fodder or forage for 30 days following application.

#### Postemergence (esters):

Limited to 1 application per crop cycle.

Maximum of 3/4 pint per acre per application.

## SOYBEANS (Preplant Only)\*

APPLICATION TIMING	AMOUNT OF TENKOZ LO-VOL 6 PER ACRE	DIRECTIONS
Preplant	1/2 to 2/3 pint	Apply not less than 7 days prior to planting soybeans, when weeds are small and actively growing. Use the higher rate on larger weeds and when perennials are present. Some weeds may require repeat treatment for adequate control (see GENERAL WEED LIST and below).
	> 2/3 to 1-1/3 pints	<p>Apply not less than 15 days prior to planting soybeans, when weeds are actively growing. Some weeds may require repeat treatment for adequate control (see GENERAL WEED LIST and below).</p> <p>In addition to those weeds found on the GENERAL WEED LIST, this product will suppress or control the following broadleaf weeds frequently encountered in reduced tillage soybean production systems: bullnettle, smallflowered bittercress, Carolina geranium, smallflowered buttercup, common and rough cinquefoil, red clover* horseweed or marestalk, mousetail, wild mustard, field pennycress, cutleaf evening primrose, common purslane, speedwell, and Virginia copperleaf.</p> <p>* These weeds are only partially controlled.</p> <p>Apply no more than 1-1/3 pints of this herbicide in one season prior to planting soybeans. After applying, plant soybean seed as deep as practical or at least 1-1/2 to 2 inches deep. Adjust the planter press wheel, if necessary, to ensure that planted seed is completely covered.</p> <p>If desired, this product may be applied pre-plant to soybeans in tank mixtures with other herbicides such as Poast®, Poast Plus®, Roundup®, Roundup D-Pak®, Honcho®, Gramoxone® Extra, Prowl®, Pursuit Plus®, Scepter®, Scepter 70 DG®, Squadron® and others that are registered for pre-plant soybean use.</p> <p>NOTE: Unacceptable injury to soybeans planted in fields previously treated with this herbicide may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather and agronomic factors such as the amount of weed vegetation and previous crop residue present that may be in effect between the time of herbicide application and the emergence of the soybean plant.</p>

\* Not currently registered in California.

## RESTRICTIONS AND LIMITATIONS FOR USE IN SOYBEANS (PRE-PLANT)

- Do not apply this product prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield.
- Do not apply this product when weather conditions such as temperature, air inversions, or wind favor drift from treated areas to susceptible plants.
- Do not mow or cultivate weeds prior to treating with this herbicide as poor control may result.
- Do not feed treated hay, forage, or fodder or graze treated soybeans to livestock.
- Only one application of this product may be made prior to planting soybeans per growing season.
- Do not feed or graze treated cover crops to livestock.

### Preplant Option 1:

Limited to 2 preplant applications per crop cycle.

Maximum of 0.75 pint per acre per preplant application.

### Preplant Option 2:

Limited to 1 application per crop cycle.

Maximum of 1.5 pints per acre per preplant application.

**Applicator may choose only one of the options for a single crop cycle (Preplant Option 1 or Preplant Option 2).**

## GRASSES GROWN FOR SEED PRODUCTION

WEEDS	AMOUNT OF TENKOZ LO-VOL 6 PER ACRE	DIRECTIONS
Annual broadleaf weeds	2/3 to 1 pint	Apply to established stands in the spring before the seed head comes into the boot stage. Do not spray in boot stage of growth. In seedling grasses apply in the spring after the grass has tillered or has at least 5 leaves but before the boot stage. Use only the low rate on seedling grasses. DO NOT USE IN CALIFORNIA
Biennial and Perennial broadleaf weeds	1-1/3 to 2 pints	

## RESTRICTIONS AND LIMITATIONS FOR USE ON GRASSES FOR SEED PRODUCTION

- Do not graze dairy animals or cut forage for hay within 7 days of application.

Limited to 2 applications per year.

Maximum of 2 pints per acre per application.

Minimum of 21 days between applications.

## ESTABLISHED GRASS PASTURES AND RANGELAND

WEEDS IN CROP	AMOUNT OF TENKOZ LO-VOL 6 PER ACRE	DIRECTIONS
Annual broadleaf weeds	1-1/3 pints	Apply when weeds are small and actively growing and prior to bud stage. Do not apply to newly seeded areas until grass is well established. Do not apply when grass is in the boot to milk stage if grass seed production is desired. Bentgrass and legumes may be injured by this treatment.
Perennial and biennial broadleaf weeds	2 to 2-2/3 pints	
Sand sage, sagebrush, shinnery oak and other 2,4-D susceptible woody species	2/3 to 2-3/4 pints	Apply aerially to brush in 2 or more gallons of water per acre. One gallon of fuel oil may be included in the mixture. Consult state or local brush control specialists for most effective rate, volume and timing of spray applications.

## FALLOW LAND AND CROP STUBBLE

WEEDS	AMOUNT OF TENKOZ LO-VOL 6 PER ACRE	DIRECTIONS
Annual broadleaf weeds	2/3 to 1-1/3 pints	Use the lower rate when weeds are small (2 to 3 inches tall) and growing actively. Use the higher rate on older and drought-stressed plants.
Biennial broadleaf weeds	2/3 to 1-2/3 pints	Spray while musk thistles or other biennial species are in the seedling to rosette stage, and before flower stalks are initiated. The lower rate can be used in spring during rosette stage. In fall or after flower stalks have developed, use the highest rate

(continued)

## FALLOW LAND AND CROP STUBBLE *(continued)*

WEEDS	AMOUNT OF TENKOZ LO-VOL 6 PER ACRE	DIRECTIONS
Perennial broadleaf weeds	1-1/3 to 3 pints	Spray weeds in bud to bloom stage, or in good vegetative growth. Do not disturb treated area for at least 2 weeks after treatment, or until weed tops are dead.
Wild garlic or onions	2-2/3 to 3 pints	Apply to new regrowth of wild onion or garlic which occurs in the fall following harvest of small grains, soybeans, corn or grain sorghum.

### RESTRICTIONS AND LIMITATIONS FOR USE ON PASTURES, RANGELAND, AND FALLOW LAND

- For ground application, a minimum of 10 to 15 gallons of water per acre is recommended for proper spray coverage.
- For aerial application, use a minimum of 2 gallons of water per acre.
- Do not graze animals on treated areas within 7 days after treatment.
- Do not permit dairy animals or meat animals being finished for slaughter to forage treated fields within 3 days of slaughter.

**Fallow land:**

Plant only labeled crops within 29 days following application.

Limited to 2 applications per year.

Maximum of 3 pints per acre per application.

Minimum of 30 days between applications.

**Pasture and Rangeland:**

Do not cut forage for hay within 7 days of application.

**Postemergence:**

For susceptible annual and biennial broadleaf weeds: Use 1.5 pints per acre per application.

For moderately susceptible biennial and perennial broadleaf weeds: Use 1.5 to 3.0 pints per acre per application.

For difficult to control weeds and woody plants: Use 3 pints/acre per application.

**Spot treatment:**

Use 3 pints per acre.

Maximum of two applications per year.

Maximum of 5.75 pints per acre per year.

Minimum of 30 days between applications.

If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.

### FENCEROWS, ROADSIDES, VACANT LOTS, AIRFIELDS, RAILROAD, HIGHWAY AND UTILITY RIGHTS-OF-WAY, AND OTHER NON-CROP AREAS

WEEDS	AMOUNT OF TENKOZ LO-VOL 6 PER ACRE	DIRECTIONS
Annual broadleaf weeds	1-1/3 to 2-2/3 pints	Apply when most weeds are still young and growing vigorously.
Perennial and biennial broadleaf weeds	2 to 3 pints	Apply when weeds are actively growing and near the bud stage, but before flowering. For best results on tansy ragwort and musk thistle, treat in rosette stage, before bolting. A second application is usually needed for best results on thistle, nettle and bindweed. Treat wild onion or garlic in early spring and in fall when they are young and growing actively.
Cattails and tules	1-1/3 quarts	Mix 1-1/3 quarts of this product per acre in 2 quarts kerosene or diesel oil then add this mixture to 100 gallons of water. This may require 300 to 500 gallons of water per acre, depending on the stand. The addition of a wetting agent such as SURFEL® spray adjuvant is suggested.
Woody plants	2/3 to 3/4 gallons	Apply preferably through low volume equipment such as DIRECTA-SPRA®, WOBBLER®, MINI-WOBBLER® or SPIROMETER®. Treat when 2,4-D susceptible species are in full leaf and growing actively. Oil or wetting agent and/or high volume spraying (300 to 500 gallons of water per acre) may be needed for acceptable control of some species or if brush is dense.



## RESTRICTIONS AND LIMITATIONS FOR USE IN NON-CROP AREAS

- For aerial application to solid stands of susceptible brush, use 1-1/3 to 2-2/3 quarts in 3 to 12 gallons volume per acre. 2 to 4 quarts of fuel oil may be included in this mixture.
- Applications will be less effective when conditions such as deficient soil moisture reduce brush growth.

### Postemergence (annual and perennial weeds):

Limited to 2 applications per year.

Maximum of 3 pints per acre per application.

Minimum of 30 days between applications.

### Postemergence (woody plants):

Limited to 1 application per year.

Maximum of 4.75 pints per acre per year.

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

## TANK MIXTURES FOR NON-CROP AREAS

### TENKOZ LO-VOL 6 AND TAHOE® 4E OR TAHOE® 3A Tank Mixtures for Non-Crop Areas

WEEDS	AMOUNT OF TENKOZ LO-VOL 6 PER ACRE	DIRECTIONS
<b>Broadleaf weed control</b>	1-1/3 to 2-2/3 pints TENKOZ LO-VOL 6 plus 2 to 6 pints TAHOE® 4E or 3 to 8 pints TAHOE® 3A	For wider spectrum control of broadleaf weeds and woody plants: Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing.
<b>Woody plant control</b> Broadcast foliar spray	2 to 3 pints TENKOZ LO-VOL 6 plus 1-1/2 to 3 quarts TAHOE® 4E or 2 to 4 quarts TAHOE® 3A	Apply as a broadcast spray in enough water to wet all parts of the brush foliage, stem and bark. This may require 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing but prior to flowering. Use the lower rates for biennials less than 3 inches rosette diameter. Use the higher rates for perennial weeds or for biennial weeds past the 3-inch rosette stage.
<b>Woody plant control</b> High volume leaf-stem treatment with plus ground equipment	5.75 pints TENKOZ LO-VOL 6 plus 1-1/2 to 12 pints TAHOE® 4E or 2 to 16 pints TAHOE® 3A	Apply as broadcast spray in enough water to thoroughly wet leaves, stems and root collars, or apply as a high volume stem foliage spray in enough volume to thoroughly wet leaves, stems and root collars (100 to 400 gallons of water per acre) or apply aerially in enough water (10 to 30 gallons per acre) using drift control equipment such as the MICRO-FOIL boom or an effective drift control agent such as LO-DRIFT spray additive. Use the higher rates and volumes when plants are dense or under drought conditions.
<b>Woody plant control</b> Aerial application (helicopter only)	5.75 pints TENKOZ LO-VOL 6 plus 3 to 4 quarts TAHOE® 4E or 4 to 6 quarts TAHOE® 3A	Apply in enough water to wet all parts of the brush foliage, stem and bark. This may require 10 to 30 gallons of water per acre using drift control equipment such as the MICRO-FOIL® boom or an effective drift control agent such as LO-DRIFT® spray additive. Use the higher rates and volumes when plants are dense or under drought conditions.

## TENKOZ LO-VOL 6 AND DIABLO® Herbicide Tank Mixtures For Non-Crop Areas

WEEDS	AMOUNT OF TENKOZ LO-VOL 6 PER ACRE	DIRECTIONS
Annual broadleaf weeds	1-1/3 to 2-2/3 pints TENKOZ LO-VOL 6 plus 1/2 to 1-1/2 pints DIABLO®	For wider spectrum control of broadleaf weeds and woody plants: Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing. Use the higher rates when treating dense or tall vegetative growth.
Perennial and biennial broadleaf weeds	2 to 3 pints TENKOZ LO-VOL 6 plus 1/2 to 6 pints DIABLO®	Apply as a broadcast spray in enough water to wet all parts of the brush foliage, stem and bark. This may require 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing but prior to flowering. Use the lower rates for biennials less than 3 inches rosette diameter. Use the higher rates for perennial weeds or for biennial weeds past the 3-inch rosette stage.
<b>Woody plant control</b> Broadcast, high volume, stem foliage or aerial application	5.75 pints TENKOZ LO-VOL 6 plus 2 to 8 quarts DIABLO®	Apply as broadcast spray in enough water to thoroughly wet leaves, stems and root collars, or apply as a high volume stem foliage spray in enough volume to thoroughly wet leaves, stems and root collars (100 to 400 gallons of water per acre) or apply aerially in enough water (10 to 30 gallons per acre) using drift control equipment such as the MICRO-FOIL boom or an effective drift control agent such as LO-DRIFT spray additive. Use the higher rates and volumes when plants are dense or under drought conditions.

### RESTRICTIONS AND LIMITATIONS FOR TANK MIXTURES FOR NON-CROP AREAS

- For postemergence annual and perennial weeds do not make more than 2 applications per year with a minimum of 30 days between applications.
- For postemergence woody plant control do not make more than 1 application per year

### SMALL AREA APPLICATIONS FOR NON-CROP USE

For control of broadleaf weeds in small non-crop areas with hand held or back-pack sprayers mix 2-2/3 fluid ounces of this herbicide per gallon of water. Thoroughly wet all weed foliage. Maintain agitation of mixture to prevent separation.

### FOREST MANAGEMENT

#### CONIFER RELEASE

To control alder, apply 2/3 to 2 quarts of this product in 9-1/2 to 15 gallons of water per acre as a foliage spray. Usually 1-1/3 quarts of this herbicide per acre is sufficient to provide good alder control with minimum conifer injury. Treat when 3/4 of the brush foliage has attained full size leaves and before new conifer growth reaches 2" in length. This is usually between early May and mid-June. Adjust treatment date depending on stage of growth of conifers and brush species. This may cause leader deformation on exposed firs, but they should overcome this during the second year after spraying.

To control tanoak, madrone, ceanothus, canyon live oak, and manzanita, and to release Douglas fir, hemlock, Sitka spruce or grand fir, apply 2 quarts of this product plus 3 quarts fuel oil in 8-1/2 to 15 gallons of water per acre before new growth on Douglas fir is 2" long. To control manzanita and ceanothus in ponderosa pine, apply a maximum of 5.75 pints of this product per acre before pine growth begins in spring.

To control hazel brush in the Lake states, apply 1-2/5 quarts of this product in 6 to 25 gallons of water per acre when new shoot growth of hazel is complete (usually mid-July).

In northern areas, if possible conifer injury can be tolerated, 1 to 2 quarts of this product applied aerially in 8 to 25 gallons of water per acre after conifers such as jack pine, red pine, black spruce, and white spruce have hardened off (usually mid-July) will provide control of competing hardwoods including alder, aspen, birch, hazel and willow.

#### SITE PREPARATION

**Budbreak Spray:** To control alder and other susceptible species before planting forest seedlings, apply 1-2/5 to 2-3/5 quarts of this product in 9 to 15 gallons of water per acre plus 2 quarts of fuel oil per acre after alder buds break, but before foliage is 1/4 full size.

**Foliage Spray:** To control alder before planting forest tree seedlings, apply 1-2/5 quarts of this product plus 2 quarts fuel oil in 9 to 15 gallons of water per acre after most alder leaves are full size.

Consult your regional or extension forester or state herbicide specialist for recommendations to fit local conditions.

This product will either kill, control or suppress the weeds listed in the label booklet for this product. Some of these species may require repeat spot applications even under ideal conditions.

## RESTRICTIONS AND LIMITATIONS FOR USE IN FOREST MANAGEMENT

### **Broadcast application:**

Limited to 1 broadcast application per year.  
Maximum of 5.75 pints per acre per broadcast application.

### **Basal spray, Cut Surface - Stumps, and Frill:**

Limit of one basal spray or cut surface application per year.  
Maximum of 11.5 pints per 100 gallons of spray solution.

### **Injection:**

Limit to one injection application per year.  
Maximum of 1.45 ml of 5.5 lb. ae formulation per injection site.

**Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA).**

**For further information, please refer to <http://www.epa.gov/espplitstatus/wtc/index.htm>**

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a dry, secured storage area. Keep container tightly closed when not in use.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### **CONTAINER DISPOSAL:**

**Nonrefillable Containers 5 Gallons or Less:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

## WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

## LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(RV032009)

All trademarks are the property of their respective owners.

**NOTES:**

SPECIMEN

**TENKOZ LO-VOL 6 2,4-D LOW VOLATILE HERBICIDE**  
**WEED AND BRUSH CONTROL IN SMALL GRAINS, FIELD CORN, RANGELAND, PASTURES, ROADSIDES**  
**AND FENCEROWS**

**ACTIVE INGREDIENT:**

2,4-Dichlorophenoxyacetic acid, isooctyl ester\* ..... 87.3%

**OTHER INGREDIENTS:** .....

.....12.7%

**TOTAL:** ..... 100.0%

Isomer Specific AOAC Method, Equivalent to:

\*2,4-Dichlorophenoxyacetic Acid ..... 57.9%, 5.5 lbs./gal.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

**SEE ATTACHED BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.**

**For Medical Emergencies Only, Call (877) 325-1840.**

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Do not get in eyes, on skin or on clothing.

**FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**NOTE TO PHYSICIAN**

No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

This product contains petroleum distillates. If large amounts, greater than 1 mg/kg body weight have been ingested, the stomach should be evacuated by gastric intubation with the aid of a cuffed endotracheal tube to prevent aspiration of petroleum distillates. After removal of stomach contents, wash stomach by instilling 30 to 50 grams of activated charcoal in 3 to 4 ounces of water through the stomach tube and again remove stomach contents. Avoid only laxatives.

**MANUFACTURED FOR**  
**TENKOZ INC.**

100 N. Point Center E. • Suite 330, Alpharetta, GA 30022

**DIRECTIONS FOR USE**

It Is A Violation Of Federal Law To Use This Product In A Manner Inconsistent With Its Labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a dry, secured storage area. Keep container tightly closed when not in use.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL: Nonrefillable Containers 5 Gallons or Less:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**EPA REG. NO. 71368-11-55467**  
**EPA EST. NO. 228-IL-1**

**Net Contents: 2.5 Gals. (3.78 L)**

**PULL HERE TO OPEN**  
**PRESS TO RESEAL**

**13047000**

**(RV032009)**