THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA Reg. No. 279-3338

Cadet® Herbicide applied post emergence alone or in combination with glyphosate plus Topsin® Fungicide (or alternative thiophanate-methyl fungicide) for suppression of White Mold (*Sclerotinia sclerotiorum*) in soybeans.

2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN THESE REGISTERED STATES: DELAWARE, MARYLAND, MICHIGAN, NEW JERSEY, OHIO, PENNSYLVANIA, VIRGINIA AND WEST VIRGINIA

THIS PRODUCT BULLETIN CONTAINS NEW
OR ADDITIONAL DIRECTIONS
FOR USE WHICH ARE RECOMMENDED BY
FMC CORPORATION UNDER
FIFRA SECTION 2(ee)

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

CROP	PEST	RATE OF APPLICATION
Soybeans	White Mold (Sclerotinia sclerotiorum) (Suppression only)	Cadet 0.5 – 0.9 fluid ounces/acre alone. or Cadet 0.4 – 0.6 fluid ounces/acre + Glyphosate 0.75 lb ai/A minimum or Cadet 0.4 – 0.6 fluid ounces/acre + Topsin at labeled rate.

DIRECTIONS FOR USE:

The effect of Cadet herbicide on white mold is not a fungicidal response, but one that involves Systemic Acquired Resistance (SAR). Higher rates will not increase the level of white mold suppression.

For optimum reduction in incidence of White Mold apply foliar broadcast to soybeans one week prior to first flower up to first flower (R1). Refer to the federal label for spray volumes and tank mix additives. Cadet may be applied alone in either conventional non-GMO soybeans or glyphosate resistant soybeans.

RESTRICTIONS:

Do not apply more than 1.25 fl. oz./acre or 0.0089 lb. ai/A per season. Do not apply within 60 days of soybean harvest.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED CADET HERBICIDE, EPA REG.NO. 279-3338, LABEL MUST BE FOLLOWED. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

