

# Specimen Label

TRICLOPYR

GROUP

4

HERBICIDE



# Turflon<sup>®</sup>

## Ester Ultra

## HERBICIDE

™Trademarks of Corteva Agriscience and its affiliated companies

**For the control of annual and perennial broadleaf weeds, bermudagrass, and kikuyugrass in ornamental turf.**

Active Ingredient:

triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, butoxyethyl ester.....

60.45%

Other Ingredients.....

39.55%

Total .....

100.00%

Acid Equivalent: triclopyr – 43.46% - 4 lb/gal

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

EPA Reg. No. 62719-566

**Keep Out of Reach of Children**

## CAUTION

**Causes Moderate Eye Irritation • Harmful If Swallowed • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid contact with skin, eyes, or clothing. Wear gloves and protective clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.**

#### Personal Protective Equipment (PPE)

**WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) – for this product, sod farm uses – must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (≥14 mils) made of barrier laminate, nitrile rubber, neoprene rubber, or Viton
- Shoes plus socks

**Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – for this product, non-sod farm uses – must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

**The requirements in this box apply to sod farm uses.**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves (≥14 mils) made of barrier laminate, nitrile rubber, neoprene rubber, or Viton
- Shoes plus socks

## Non-Agricultural Use Requirements

The requirements in this box apply to all use sites on this label except for sod farm uses.

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Entry Restrictions for Non-WPS Uses:** For applications to turf other than sod farms or turf grown for seed, do not allow entry into areas until sprays have dried.

## Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal. Open dumping is prohibited.

**Pesticide Storage:** Store above 28°F or agitate before use.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### Nonrefillable containers 5 gallons or less:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

### Refillable containers 5 gallons or larger:

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

### Nonrefillable containers 5 gallons or larger:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## Product Information

Use Turflon® Ester Ultra herbicide for the control of actively growing annual and perennial broadleaf weeds, bermudagrass, and kikuyugrass in perennial bluegrass, perennial ryegrass, or tall fescue ornamental turf including these turfgrasses in sod farms and golf courses.

## Use Precautions

Turflon Ester Ultra is formulated as a low volatile ester. However, the combination of spray contact with impervious surfaces such as roads and rocks, and increasing ambient air temperatures, may result in an increase in the volatility potential for this herbicide, increasing a risk for off-target injury to sensitive crops such as grapes and tomatoes.

## Use Restrictions

**Chemigation:** Do not apply this product through any type of irrigation system.

Do not apply Turflon Ester Ultra directly to, or otherwise permit it to come into direct contact with, cotton, grapes, tobacco, vegetable crops, flowers, fruit or orchard trees, shrubs, or other desirable broadleaf plants. Do not permit spray mists containing Turflon Ester Ultra to drift onto such plants.

To minimize turf injury, do not treat if turf is under heat or drought stress. Reapply Turflon Ester Ultra at 4-week intervals.

Mow newly seeded turf 2 or 3 times before treating. Do not water for 24 hours after application. Do not reseed for 3 weeks after application. (This restriction does not apply when bermudagrass turf is overseeded with perennial ryegrass at a minimum reseeding rate of 400 lb per acre.)

Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.

- Do not apply to ditches currently being used to transport irrigation water.
- Do not apply where runoff or irrigation water may flow onto agricultural land as injury to crops may result.
- Do not use on other turfgrass species such as bahiagrass, bentgrass, bermudagrass, centipedegrass, or St. Augustinegrass unless injury can be tolerated.
- Do not apply to exposed roots of shallow rooted trees and shrubs.
- Do not apply to golf course greens.
- For spot treatments, do not apply more than 2 quarts of Turflon Ester Ultra per acre or 1.5 fl oz per 1000 sq ft in a single application.

## Resistance Management

Triclopyr, the active ingredient in this product, is a Group 4 herbicide based upon the mode of action classification system of the Weed Science Society of America. Appropriate resistance management strategies should be followed.

- Development of plant populations resistant to the mode of action of aminopyralid is usually not a problem on non-cropland sites since these sites receive infrequent pesticide applications.
- Similar looking biotypes of a given weed species occurring in a treated area may vary in their susceptibility to an herbicide. Application of an herbicide below its labeled rate may allow more tolerant weeds to survive and a shift to more tolerant biotypes within the treated area.
- Where identified, spreading of resistant weeds to other fields may be prevented by cleaning harvesting and tillage equipment before moving to other areas and by planting weed-free seed.
- Scout before after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as mowing.
- Use tank mixtures with herbicides from a different group if such use is permitted. Where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- For further information or to report suspected resistance, contact your extension specialist or contact a customer service representative at 800-992-5994.

### Avoiding Injurious Spray Drift

Make applications only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants that are near enough to be injured. Spray drift can be reduced by keeping the spray boom as low as possible; by using higher volume sprays and lower pressures; using nozzles that deliver larger spray droplets; and by spraying when wind velocity is low.

### Broadleaf Weeds Controlled by Turflon Ester Ultra

black medic	curly dock	lespedeza	sweet clover
bull thistle	dandelion	matchweed	vetch
burdock	field bindweed	mustard	wild carrot (Queen Anne's lace)
Canada thistle	goldenrod	oxalis	wild lettuce
chicory	ground ivy	plantain	wild violet
clover	kikuyugrass (1)	ragweed	yarrow
creeping beggarweed	lambquarters	smartweed	

(1) See control of kikuyugrass under Application Instructions for Turf.

### Mixing Instructions

Add about one-half of the required amount of clean water to the spray tank. Add Turflon Ester Ultra and complete addition of water with agitation running. Mix thoroughly and continue moderate agitation while spraying. When Turflon Ester Ultra is mixed with water, it forms an emulsion, not a solution; separation may occur unless the spray mixture is agitated continuously.

**Tank Mixing:** To improve the spectrum of activity, Turflon Ester Ultra may be tank mixed at a rate of 1/2 to 1 pint per acre with labeled rates of low volatile amine or ester formulations of 2,4-D, MCPP, or other labeled postemergence broadleaf herbicides provided the tank mix product is registered for use on ornamental turf (perennial bluegrass, perennial ryegrass, or tall fescue ornamental turf including these turfgrasses in sod farms and golf courses). Refer to the product labels of tank mix products and follow all appropriate use directions, precautions, and limitations. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### Application Instructions for Turf

#### Broadcast Treatment of Ornamental Turf

Apply foliage sprays during warm weather, early spring through fall, when weeds are actively growing. Apply 1/2 to 1 quart of Turflon Ester Ultra per acre in enough water to control broadleaf weeds actively growing in perennial bluegrass, perennial ryegrass, or tall fescue. Only emerged weeds present at the time of application will be controlled. Apply using equipment that assures uniform coverage of the target area. When treating mature plants or hard to control species, and for applications made during drought conditions, use the higher rates of Turflon Ester Ultra. Applications made under drought conditions may provide less than desirable results.

#### Spot Treatment of Ornamental Turf

Mix 3/8 to 3/4 fl oz of Turflon Ester Ultra per 1000 sq ft in enough water to provide uniform coverage of the target area. Apply at any time broadleaf weeds are susceptible.

#### Control of Kikuyugrass

Apply Turflon Ester Ultra at a rate of 1/2 to 1 quart per acre. To improve activity, MSMA herbicide may be tank mixed with the 1/2 quart per acre rate of Turflon Ester Ultra. Three to 4 additional applications at 4- to 6-week intervals may be required to achieve control of kikuyugrass.

#### Suppression and Control of Bermudagrass

Apply Turflon Ester Ultra at the rate of 1 quart per acre. Three to 4 additional applications at 4-week intervals are required for adequate suppression of bermudagrass and to allow fescue or other desired turfgrass species to dominate. To improve suppression and control of bermudagrass, tank mix 1 quart of Turflon Ester Ultra per acre with a postemergence grass herbicide registered for this use pattern. Three to 4 additional applications of this tank mix at 4-week intervals should be made to achieve control. Reseeding following application will accelerate the transition to cool season turf.

### Suppression of Bermudagrass When Overseeding with Perennial Ryegrass in the States of Arizona, California, Nevada, New Mexico, and Utah

Regulation of bermudagrass growth by applying Turflon Ester Ultra is characterized by slowing and/or stopping of growth. Some browning of the bermudagrass may also occur. Hybrid varieties of bermudagrass are more susceptible to growth regulation than common bermudagrass and the effects may be more pronounced. The degree of growth regulation and browning is rate-related. Test the growth regulation effect of Turflon Ester Ultra on small areas until the user is comfortable with the results in his particular turf management system. A rate can then be chosen that will provide the desired level of response. The length of the growth regulation effect is also rate-related with the higher rates providing a longer period of suppression. The overseeded ryegrass will not be affected by this treatment. Do not apply to golf course putting greens.

Apply 4 to 8 fl oz of Turflon Ester Ultra per acre to hybrid bermudagrass and 6 to 10 fl oz of Turflon Ester Ultra per acre to common bermudagrass.

**Prior to overseeding:** Apply to fairways from 2 weeks period to overseeding to just before overseeding. For best results, apply Turflon Ester Ultra prior to any aggressive verticutting, scalping, or other seedbed preparation practices to ensure that adequate leaf surface of bermudagrass is available for update of Turflon Ester Ultra. A single application of Turflon Ester Ultra prior to overseeding usually provides about 30 days of bermudagrass growth suppression.

**After overseeding:** If temperatures are sufficient to support active growth of bermudagrass, additional applications of Turflon Ester Ultra can be made to extend the plant growth regulation effect. If additional applications are required, make them after the overseeded ryegrass is established, approximately 3 weeks after seeding. Allow a minimum of 4 weeks between applications.

### Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

### Warranty Disclaimer

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. To the extent permitted by law, all such risks shall be assumed by the buyer.

### Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

### Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. In no case shall Corteva Agriscience be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™®Trademarks of Corteva Agriscience and its affiliated companies

**Produced for**  
**Corteva Agriscience LLC**  
**9330 Zionsville Road**  
**Indianapolis, IN 46268**

Label Code: CD02-385-022

Replaced Label: D02-385-002 and D02-385-003

EPA accepted 03/21/19

## **Revisions:**

The following revisions were made via Non-Notification on April 11, 2024.

1. Removed pasture, rangeland, grazing, and non-crop applications and usages throughout the package label.
  - a. Removed 'rangeland and permanent grass pastures, and non-crop areas....grazed areas' usage from 'Marketing Information' section.
  - b. Removed 'rangeland and permanent grass pastures, and non-crop areas....grazed areas' usage from 'Product Information' section.
  - c. Removed non-crop restriction 'It is permissible to treat non-irrigation...' from 'Use Restrictions' section.
  - d. Removed 'Grazing,' 'Haying (harvesting of dried forage,' and 'Slaughter Restriction' from 'Use Restrictions' section.
  - e. Removed 'Application Instructions for Non-Crop Areas' section.
  - f. Removed 'Applications Methods' section for grazing applications.
  - g. Removed 'Foliage Treatment with Ground Equipment' section for grazing applications.
  - h. Removed 'Broadcast Applications with Ground Equipment' section for grazing applications.
  - i. Removed 'Aerial Application' section for rangeland and pasture applications.
  - j. Removed 'Basal Bark, Dormant Stem, and Cut Surface Treatments' section for grazing applications.
  - k. Removed 'Rangeland and Permanent Grass Pastures' application information section.
2. Removed 'or zoysiagrass ornamental turf' from 'Product Information' section.
3. Removed 'zoysiagrass' from 'Broadcast Treatment of Ornamental Turf' section.