

WINFIELD™

Herbivore™ Herbicide

FIFRA Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF MINNESOTA **FOR USE ON DRY BEANS POST – EMERGENCE**

This label is valid until otherwise amended, withdrawn, canceled or suspended.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Herbivore™ Herbicide and this Section 24(c) label before applying. Do not use in manner inconsistent with applicable use directions, restrictions, precautions and limitations provided on the label affixed to the container for Herbivore™ Herbicide and this Section 24(c) label.

ACTIVE INGREDIENT:
Halosulfuron-methyl.....75.0%
OTHER INGREDIENTS:..... 25.0%
TOTAL 100.0%

EPA Reg. No. 2749-528-1381

EPA SLN No. MN-110005

KEEP OUT OF REACH OF CHILDREN **CAUTION**

DIRECTIONS FOR USE

FOR USE ON DRY BEANS POST-EMERGENCE

The pre-harvest interval (PHI) is the required days between the last application of Herbivore™ Herbicide and the harvesting of the crop.

Crop	Rate Oz./Acre	PHI	LIMITATIONS
DRY BEANS	1/2 - 2/3	30	Do not apply more than 1.33 oz. of this product per acre per crop cycle, not to exceed 2 oz. (0.094 lb. active ingredient) per acre per 12-month period (includes applications to the crop and to Row Middles/Furrows). Do not make more than one post-emergence broadcast application per crop cycle.

For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 10 gallons of water per acre.

Use Rate 1/2 - 2/3 oz.:

Post-emergence- Use this product when plants have 1-3 trifoliate leaves, but before flowering. For best results, apply to weeds less than 6 inches high. Use a nonionic surfactant (NIS).

Tank Mixtures

If Herbivore™ Herbicide is utilized with a tank mixture partner(s) to improve annual broadleaf weed or post emergence grass control with graminicides, refer to the specific partner label(s) and observe all the precautionary statements and use directions including pre-harvest intervals, crop rotation restrictions, mixing and application instructions. Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures. For current information on the best tank mixture partner in your area, consult with the local dealer, distributor or State Agricultural Extension service.

Not all hybrids/varieties have been tested for sensitivity to this product. For untested varieties, a small amount of the field should be sprayed to determine potential sensitivity to its use. The user assumes responsibility for such use and any plant injury that may occur.

Herbivore™ Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Herbivore™ Herbicide and other Group 2 herbicides. Weed species with acquired resistance to Group 2 may eventually dominate the weed population if Group 2 herbicides are used repeatedly in the same field or in successive years as primary method of control for target species. This may result in partial or total loss of control of those species by Herbivore™ Herbicide or other Group 2 herbicides.

Herbivore and Winfield are trademarks of Winfield Solutions, LLC.

Manufactured for:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164

01/11/2012