

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3426

FOR DISTRIBUTION AND USE
ONLY IN TEXAS

THIS RECOMMENDATION, WHICH
CONTAINS ADDITIONAL DIRECTIONS
FOR USE, IS MADE AS PERMITTED
UNDER FIFRA SECTION 2(ee) AND HAS
NOT BEEN SUBMITTED TO OR
APPROVED BY THE EPA.

This recommendation for MUSTANG MAXX Insecticide is valid until 03/27/2025, or until withdrawn, cancelled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CROP	PEST	RATE OF APPLICATION
Section 3 Registered Crops*	Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)
DIRECTIONS FOR USE: For boll weevil control on volunteer cotton. Apply MUSTANG MAXX insecticide to registered crops (see section 3 label) where volunteer cotton is present. Multiple applications may be necessary to reduce pest populations to acceptable levels. Refer to each crop on the Mustang Maxx insecticide Section 3 label to determine the maximum rate allowed per season, application intervals and other restrictions.		
*RESTRICTIONS: *Refer to each crop section on the Mustang Maxx Insecticide Section 3 label for restrictions.		



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

Head and Stem Brassica Vegetables (1 day PHI) including: Broccoli; Chinese Broccoli (gailon, white flowering broccoli); Brussels Sprouts; Cauliflower; Cavalo broccolo; Kohlrabi; Cabbage; Chinese Cabbage (napa); Chinese Mustard Cabbage (gai choy). Leafy Brassica Greens (1-day PHI) including: Broccoli Raab (rapini); Chinese cabbage (bok choy); Collards; Kale; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens; Turnip Greens.

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air.
<ul style="list-style-type: none">Do not make applications less than 7 days apart.A maximum of 24 fl oz/A (0.15 lb ai/A) may be applied per season.		

Bulb Vegetables (*Allium* spp.) (7day PHI) including: Garlic; Garlic, Great-Headed (elephant); Green Eschalots; Japanese Bunching Onions; Leeks; Onion, Dry Bulb and Green; Onion, Welch; Shallots, Dry Bulb and Green; Spring Onion or Scallions

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in a minimum of 20 gal/A with ground equipment or in a minimum of 3 gal/A by aircraft. Begin applications when pests appear and repeat as necessary to maintain control.
<ul style="list-style-type: none">Do not make applications less than 7 days apart.Do not apply more than 20 fl oz/A (0.125 lb ai/A) per season.Do not graze livestock in treated areas or cut treated crops for feed.		

Corn (Field), Field Corn Grown for Seed, Popcorn

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
<ul style="list-style-type: none">Do not make applications less than 3 days apart.Do not apply more than 16 fl oz/A (0.1 lb ai/A) per season including At-planting plus foliar applications of MUSTANG MAXX Insecticide.Do not apply within 30 days of harvest for grain and stover and 60 days for forage.		



RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

Canola, Crambe, Rapeseed, Borage, Cuphea, Echium, Flax, Gold of Pleasure, Hare's-Ear Mustard, Lesquerella, Lunaria, Meadowfoam, Milkweed, Mustard, Oil Radish, Poppy Seed, Sesame, and Sweet Rocket (7 Day PHI).

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
<ul style="list-style-type: none">Do not make applications less than 7 days apart.Do not apply more than 24 fl oz/A (0.15 lb ai/A) per season.		

Cucurbit Vegetables Group (1 day PHI) including: Chayote (fruit); Chinese Waxgourd (Chinese Preserving Melon); Citron Melon; Cucumber; Gherkin; Gourd (edible) (including hyotan, cucuzza, hechima, Chinese orkra); *Mormordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon (hybrids and/or cultivars of *Cucumis melo*) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin; Summer Squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Winter Squash (includes butternut squash, calabaza, hubbard squash, acorn squash, and spaghetti squash); Watermelon (includes hybrids and varieties).

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
<ul style="list-style-type: none">Do not make applications less than 7 days apart.Do not apply more than 24 fl oz/A (0.15 lb ai/A) per season.		

Fruiting Vegetables (except Cucurbits) (1 day PHI) including: Eggplant; groundcherry (*Physalis* spp.); okra; pepino (Melon pear); pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); tomatillo; tomato.

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
<ul style="list-style-type: none">Do not make applications less than 7 days apart.Do not apply more than 24 fl oz/A (0.15 lb ai/A) per season.		



RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

Grass Forage, Fodder, and Hay Group and Grass Grown for Seed and Pasture and Rangeland (0 day PHI for forage and hay; 7 day PHI for straw and seed screenings) including: bahiagrass, barnyardgrass, bentgrass, Bermudagrass, Kentucky bluegrass, big bluestem, smooth brome grass, buffalograss, reed canarygrass, centipedegrass, crabgrass, cupgrass, dallisgrass, sand dropseed, Kentucky fescue, meadow foxtail, eastern gramagrass, side-oats grama, guinea grass, Indian grass, Johnsongrass, lovegrass, napiergrass, oatgrass, orchardgrass, pangolagrass, paspalum, redtop, Italian ryegrass, St. Augustine grass, sprangletop, squirreltailgrass, stargrass, switchgrass, timothy, crested wheatgrass, wildrye grass and zoysia grass. Also included are sudangrass and sorghum forages and their hybrids.

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment.
<ul style="list-style-type: none"> • Do not make applications less than 7 days apart for forage and hay; not less than 17 days for straw and seed screenings. • Do not spray livestock. Allow application to dry before letting livestock graze on treated area. A maximum of 4 fl oz/A (0.025 lb ai/A) may be applied per cutting. • For hay, a maximum of 16 fl oz/A (0.1 lb ai/A) per season may be applied. For forage, straw, and seed screenings, a maximum of 20 fl oz/A (0.125 lb ai/A) per season may be applied. • Applications may be made up to 0 days for forage and hay; 7 days for straw and seed screenings. 		

Leafy Vegetables (except Brassica) (1 day PHI): Amaranth (leafy amaranth, Chinese spinach, tampala); Arugula (Roquette); Cardoon; Celery; Celery, Chinese; Celtuce; Chervil; Chrysanthemum, edible-leaved and garland; Cilantro (not for use on cilantro grown for seed or coriander); Corn salad; Cress, garden; Cress, upland (yellow rocket, winter cress); Dandelion; Dock (sorrel); Endive (escarole); Fennel, Florence (finochio); Lettuce, head and leaf; Orach; Parsley; Purslane, garden; Purslane, winter; Radicchio (red chicory); Rhubarb; Spinach (including New Zealand and vine, Malabar spinach, Indian spinach); Swiss chard.

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air.
<ul style="list-style-type: none"> • Do not make applications less than 7 days apart. • A maximum of 24 fl oz/A (0.15 lb ai/A) may be applied per season. 		



RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

Legume Vegetables - Succulent and Dried (except Soybeans), 1 day PHI for succulent shelled or edible- podded peas or beans and 21 day PHI for dried shelled peas or beans

Succulent Edible-Podded Peas, Succulent Shelled Peas and Dried Shelled Peas (*Pisum* spp.) including:

Dwarf Pea; Edible-pod Pea; Snow Pea; Sugar Snap Pea; Pigeon pea; English Pea; Garden Pea; Green Pea; Lentil.

Succulent Edible-Podded Beans, Succulent Shelled Beans, and Dried Shelled Beans including: Runner Bean; Snap Bean; Wax Bean; Asparagus Bean; Chinese Longbean; Moth Bean; Yardlong Bean; Jackbean; Soybean (immature seed); Swordbean; Lima Bean; Broad Bean (Fava Bean); Blackeyed Pea; Southern Pea; Grain Lupin; Sweet Lupin; White Lupin; White Sweet Lupin; Field Bean; Kidney Bean; Navy Bean; Pinto Bean; Tepary Bean; Adzuki Bean; Catjang; Cowpea; Crowder Pea; Moth Bean; Mung Bean; Rice Bean; Urd Bean; Chickpea (Garbanzo Bean); Guar; Lablab bean.

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
<ul style="list-style-type: none">Do not make applications less than 5 days apart.Do not apply more than 24 fl oz/A (0.15 lb ai/A) per season including at-plant plus foliar applications.		

Peanut (7 day PHI)

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
<ul style="list-style-type: none">Do not make applications less than 14 days apart.Do not apply more than 24 fl oz/A (0.15 lb ai/A) per season.Do not graze livestock in treated area.Do not use treated vines or hay for animal feed.		



RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

Root and Tuber Vegetables Group 1 (except Sugar Beet) (1 day PHI) including: Arracacha; Arrowroot; Artichoke (Chinese and Jerusalem); Garden Beet; Edible Burdock; Edible Canna; Carrot; Cassava (Bitter and Sweet); Celeriac (Celery Root); Chayote (Root); Turnip-Rooted Chervil; Chicory; Chufa; Dasheen (Taro); Ginger; Ginseng; Horseradish; Leren; Turnip-Rooted Parsley; Parsnip; Potato; Oriental Radish (Daikon); Radish; Rutabaga; Salsify (Oyster Plant); Black Salsify; Spanish Salsify; Skirret; Sweet Potato; Tanier (Cocoyam); Turmeric; Turnip; Yam Bean; and Yam (True).

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
<ul style="list-style-type: none">Do not make applications less than 14 days apart.Do not apply more than 24 fl oz/A (0.15 lb ai/A) per season.Leaves of Root and Tuber Vegetables (except Sugar Beet tops) cannot be used for food or feed.		

Sorghum (Grain) and Millet (14 day PHI for grain and stover; 45 day PHI for forage)

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
<ul style="list-style-type: none">Do not make applications less than 10 days apart.Do not apply more than 20 fl oz/A (0.125 lb ai/A) per season.		

Soybeans (21 day PHI)

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
<ul style="list-style-type: none">Do not make applications less than 7 days apart.Do not apply more than 24 fl oz/A (0.15 lb ai/A) per season.Do not graze or harvest treated soybean forage, straw, or hay for livestock feed.		



RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

Sugarcane (21 day PHI)

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
<ul style="list-style-type: none">Do not make applications less than 21 days apart.Do not apply more than 16 fl oz/A (0.1 lb ai/A) per season.		

Sunflower, Castor Oil Plant, Chinese Tallowtree, Euphorbia, Evening Primrose, Jojoba, Niger Seed, Rose Hip, Stokes Aster, Tallowwood, Tea Oil Plant, and Vernonia (30 day PHI)

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
<ul style="list-style-type: none">Do not apply more than 20 fl oz/A (0.125 lb ai/A) per season including at-plant plus foliar applications.Do not make more than 5 applications at the maximum application rate per season.Do not graze livestock in treated areas or cut treated crops for feed.Avoid applications when honey bees are actively foraging by applying during the early morning or evening hours.		

Wheat and Triticale (14 day PHI for grain, forage, and hay)

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
<ul style="list-style-type: none">Do not make applications less than 14 days apart.Do not apply more than 20 fl oz/A (0.125 lb ai/A) per season.		

