

# RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's Certification.

ZETA-CYPERMETHRIN

GROUP

3A

INSECTICIDE



## FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3426

For Distribution and Use Only in  
Texas

THIS RECOMMENDATION, WHICH CONTAINS  
ADDITIONAL DIRECTIONS FOR USE, IS MADE  
AS PERMITTED UNDER FIFRA SECTION 2(ee)  
AND HAS NOT BEEN SUBMITTED TO OR  
APPROVED BY THE EPA.

This 2(ee) recommendation for MUSTANG<sup>®</sup> MAXX Insecticide is valid until December 31, 2025, or until withdrawn, cancelled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

**For the suppression of Cabbage maggot and Seedcorn maggot in  
Head and Stem Brassica Vegetables Crop Group 5-16 and  
Leafy Brassica Greens Crop Subgroup 4-16B**

PEST	RATE OF APPLICATION	USE DIRECTIONS
Cabbage maggot Seedcorn maggot	3.2 to 4.0 fl oz/A (0.02 to 0.025 lb ai/A)	<b>Application by ground:</b> Apply in a minimum of 15 gallons of finished spray per acre.

### RESTRICTIONS:

- Do not apply more than 4.0 fl oz/A of product (0.025 lb ai/A) per application.
- Do not apply more than 24 fl oz/A of product (0.15 lb ai/A) per year.
- Do not make more than 6 applications per year when applications are made at the maximum rate.
- Do not make applications less than 7 days apart.
- **PHI:** Do not apply within 1 day of harvest.

### REMARKS:

Performance and crop protection will be best when MUSTANG<sup>®</sup> MAXX insecticide is incorporated into an IPM program that includes measures that reduce risk for maggot infestation like the following: seed treatments (if needed for seedcorn maggot), avoid planting into cold, wet soils, heavily manured fields, or fields previously planted with brassica crops; crop rotation; and controlling alternate weed hosts.

**Coverage is critical.** Applications directed to the base of the plant or banded over the row are recommended for best performance. Consider using degree-day models or other means to predict periods of fly activity; multiple applications will likely be required for best protection but observe restrictions on re-application interval and days-to-harvest.



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000