

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



FIFRA 2(ee) Recommendation
EPA Reg. No. 279-3426
FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA, OREGON, AND WASHINGTON
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for MUSTANG MAXX Insecticide is valid until December 31, 2027, or until withdrawn, cancelled, or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CROP	PEST	RATE OF APPLICATION
Grapes: Table, Raisin, Wine	European Grapevine Moth Grape Leaf folder Light Brown Apple Moth Omnivorous Leafroller Orange Tortrix Western Grapeleaf Skeletonizer Spiders	4.0 fl oz/A (0.025 lb ai/A)

DIRECTIONS FOR USE:

- Thorough coverage is essential to achieve control.
- Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gal/A of finished spray by ground and 2 gal/A of finished spray by air).
- Multiple applications may be required.

Restrictions:

- Do not make applications less than 7 days apart.
- Do not apply more than 4.0 fl oz/A of product (0.025 fl oz/A) per application.
- Do not make more than 6 applications per year.
- Do not apply more than 24 fl oz/A of product (0.15 lb ai/A) per year.
- Do not apply within 1 day of harvest.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000