


**RESTRICTED USE PESTICIDE**

**Toxic to fish and aquatic organisms.**

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**

 <b>MUSTANG<sup>®</sup></b> <b>MAXX</b> INSECTICIDE  <b>EPA REG. NO. 279-3426</b>	<b><u>2(ee) Recommendation</u></b>
	<b>FOR DISTRIBUTION AND USE ONLY IN Delaware, Maryland, New Jersey, Pennsylvania, Virginia, West Virginia</b>
	<b>IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING</b>
	<b>THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)</b>

For the suppression or control of spotted lanternfly in listed crops.

CROP	RATE OF APPLICATION	RESTRICTIONS
Grape	4.0 fl oz/A (0.025 lb ai/A)	<ul style="list-style-type: none"><li>• Do not apply more than 4.0 fl oz of product (0.025 lb ai) per acre per application.</li><li>• Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season.</li><li>• Do not make applications less than 7 days apart.</li><li>• Do not apply within 1 day of harvest.</li></ul>

**Use Directions:**

Spray in sufficient water to provide thorough coverage for best control.

**Application by ground:** Apply in a minimum of 10 gallons of finished spray per acre.

**Application by air:** Apply in a minimum of 2 gallons of finished spray per acre.

CROP	RATE OF APPLICATION	RESTRICTIONS
Pome Fruit Group, including: Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental Pear, and Quince	4.0 fl oz/A (0.025 lb ai/A)	<ul style="list-style-type: none"><li>• Do not apply more than 4.0 fl oz of product (0.025 lb ai) per acre per application.</li><li>• Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season.</li><li>• Do not apply as a ULV spray.</li><li>• Do not feed or allow livestock to graze on cover crops from treated orchards.</li></ul>



		<ul style="list-style-type: none"> <li>• Do not make applications less than 7 days apart.</li> <li>• Do not apply within 14 days of harvest.</li> </ul>
<p><b>Use Directions:</b>          Avoid applications when honey bees are actively foraging by applying during the early morning or evening hours.          Spray in sufficient water to provide thorough coverage for best control.  <b>Application by ground:</b> Apply in a minimum of 20 gallons finished spray per acre for concentrate spray, or in a minimum of 100 gallons of finished spray for dilute spray.  <b>Application by air:</b> Apply in a minimum of 10 gallons finished spray per acre.</p>		

CROP	RATE OF APPLICATION	RESTRICTIONS
<b>Stone Fruit Group, including: Apricot, Cherry (Sweet and Tart), Nectarine, Peach, Plum (including Chickasaw Plum, Damson Plum, and Japanese Plum), Plumcot, and Prune (fresh)</b>	<b>4.0 fl oz/A (0.025 lb ai/A)</b>	<ul style="list-style-type: none"> <li>• Do not apply more than 4.0 fl oz of product (0.025 lb ai) per acre per application.</li> <li>• Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season.</li> <li>• Do not apply as a ULV spray.</li> <li>• Do not feed or allow livestock to graze on cover crops from treated orchards.</li> <li>• Do not make applications less than 7 days apart.</li> <li>• Do not apply within 14 days of harvest.</li> </ul>
<p><b>Use Directions:</b>          Spray in sufficient water to provide thorough coverage for best control.  <b>Application by ground:</b> Apply in a minimum of 20 gallons finished spray per acre for concentrate spray, or in a minimum of 100 gallons of finished spray for dilute spray.  <b>Application by air:</b> Apply in a minimum of 10 gallons finished spray per acre.</p>		

<p><b>REMARKS:</b>          Thorough coverage is essential to achieve control.</p> <p>Multiple applications may be required to maintain control and prevent reinfestation from migrating adults into the crop. Spotted lanternflies are mobile and may reinfest crops quickly. If another treatment is needed before the minimum retreatment interval, use another insecticide to maintain control.</p> <p><b>ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED MUSTANG® MAXX INSECTICIDE (EPA REG. NO. 279-3426) LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.</b></p>
---

Mustang® and FMC are trademarks of FMC Corporation or an affiliate. ©2018 FMC Corporation.

R-4087 081018 08-10-18

