

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA**



**EPA Reg. No. 279-3464**

**2(ee) Recommendation  
FOR DISTRIBUTION AND USE  
ONLY IN  
WASHINGTON, OREGON, IDAHO & MONTANA**

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL PESTS WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING**

Crop	Pest	Rate of Application
Spring Wheat Winter Wheat	Downy Brome <i>Bromus Tectorum</i>	See Table 1 for Rates and Timing

**Table 1. Use Rates of Anthem Flex in Spring and Winter Wheat**

Application Timing	Use Rate by Soil Texture <sup>1</sup>		Anthem Flex fl oz/A (lb a.i./A)
	Coarse	Medium	Fine
Preplant Surface	2.0 – 2.7 (0.078-0.085)	2.75 – 3.5 (0.086-0.109)	3.5 - 4.55 (0.109-0.141)
Preemergence	2.0 – 2.7 (0.078-0.085)	2.75 – 3.5 (0.086-0.109)	3.5 - 4.55 (0.109-0.141)

<sup>1</sup> Refer to soil texture section of the Anthem Flex Herbicide Section 3 label for definitions of soil texture groups.

**DIRECTIONS FOR USE: Spring and Winter Wheat**

- When applied as directed, Anthem Flex will provide control of downy brome.
- Apply as a preplant, preemergence or delayed preemergence.
- To reduce the risk of crop response apply preplant or preemergence to soils with a CEC greater than 15 and pH lower than 7.5 and an organic matter content greater than 2%.
- Anthem Flex Herbicide may be mixed with other approved herbicides to enhance the control of emerged weeds.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED ANTHEM FLEX HERBICIDE LABEL (EPA REG. NO.279-3464) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**