



FIFRA 2(ee) Recommendation
EPA Reg. No. 279-3588
FOR DISTRIBUTION AND USE IN TEXAS
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for Rhyme™ fungicide is valid until March 1, 2025, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR CHEMIGATION USE IN LISTED CROPS

Directions for Use					
When applied as directed through drip irrigation or micro sprinklers, Rhyme fungicide will provide control of listed foliar diseases or suppression of listed soilborne diseases.					
Crop	Pest	Rate (fl oz/A)	PHI	Retreatment Interval (Days)	Restrictions
Vegetables, Cucurbit Group 9 (See label for list of crops)	Anthracnose (<i>Colletotrichum</i> sp.) Gummy Stem Blight (<i>Didymella bryoniae</i>) Powdery Mildew (<i>Sphaerotheca fuliginea</i> , <i>Erysiphe cichoracearum</i>) Target Spot (<i>Corynespora cassiicola</i>) Soilborne diseases: Charcoal Rot (<i>Macrophomina phaseolina</i>) Fusarium sp. (<i>Fusarium oxysporum</i>)	7	0	14	<ul style="list-style-type: none"> Do not apply more than 7.0 fl. oz. (0.114 lb. ai) product/A/ application. Do not apply more than 28 fl. oz. of product/A/year. Do not make more than 4 applications/ year. Do not apply more than 0.455 lb. ai of flutriafol or flutriafol containing products/ A/year.
Use Directions:					
Apply preventively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development and to provide extended foliar protection. Best results for both foliar and soilborne pathogens have been obtained when Rhyme fungicide is applied preventively, as early as 14 days after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.					

Crop	Pest	Rate (fl oz/A)	PHI	Retreatment Interval (Days)	Restrictions
Vegetables, Fruiting Group 8-10 (See label for list of crops) Tomatoes (See label for list of crops)	Anthracnose (<i>Colletotrichum coccodes</i> , <i>C.acutatum</i>) Black Mold (<i>Alternaria alternata</i>) Early Blight (<i>Alternaria solani</i>) Cercospora Leaf Spot (<i>Cercospora capsici</i>) Powdery Mildew (<i>Leveillula taurica</i> , <i>Oidium neolycopersici</i>) Target Spot (<i>Corynespora cassicola</i>) Soilborne diseases: Charcoal Rot (<i>Macrophomina phaseolina</i>) Fusarium sp. (<i>Fusarium oxysporum</i>)	7	0	14	<ul style="list-style-type: none"> Do not apply more than 7.0 fl. oz. (0.114 lb ai) product/A/ application. Do not apply more than 28 fl. oz. of product/A/year. Do not make more than 4 applications/ year. Do not apply more than 0.455 lb. ai of flutriafol or flutriafol containing products/A/year.

Use Directions:

Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Best results for both foliar and soilborne pathogens have been obtained when Rhyme fungicide is applied preventively, shortly after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.

Crop	Pest	Rate (fl. oz./A)	PHI (days)	Retreatment Interval (Days)	Restrictions
Grapes (fresh, juice, table, wine, raisin)	Powdery Mildew (<i>Uncinula necator</i>)	5	14	14	<ul style="list-style-type: none"> Do not apply more than 5.0 fl. oz. (0.11 lb ai) product/A/ application. Do not apply more than 30 fl. oz. of product/A/year. Do not apply more than 0.488 lb. ai of flutriafol or flutriafol containing products/A/year.

Use Directions:

Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development and to provide extended foliar protection. Best results have been obtained when Rhyme fungicide is applied



preventively, as early as bud break, to ensure time for movement of flutriafol into the foliage. Once fruit are present, foliar applications are needed for fruit protection.

Crop	Pest	Rate (fl. oz./A)	PHI (days)	Retreatment Interval (Days)	Restrictions
Strawberry	Powdery Mildew <i>(Podosphaera aphanis)</i> Anthracnose <i>(Colletotrichum acutatum)</i> Soilborne diseases: Charcoal Rot <i>(Macrophomina phaseolina)</i>	7	7	14	<ul style="list-style-type: none"> Do not apply more than 7.0 fl. oz. (0.114 lb. ai) product/A/ application. Do not apply more than 28 fl. oz. of product/A/year. Do not make more apply more than 4 applications/year. Do not apply more than 0.455 lb. ai of flutriafol or flutriafol containing products/A/year.

Use Directions:

Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Best results for both foliar and soilborne pathogens have been obtained when Rhyme fungicide is applied preventively, shortly after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.

Table 1. Rhyme fungicide rate conversion chart for drip (trickle) chemigation soil application.

fl. oz./A	Fluid ounces Rhyme fungicide/1000 row feet based on planted row spacing												
	30" rows	34" rows	36" rows	38" rows	40" rows	44" rows	48" rows	60" rows	66" rows	72" rows	80" rows	84" rows	120" rows
5	0.29	0.33	0.34	0.36	0.38	0.42	0.46	0.57	0.63	0.69	0.77	0.80	1.15
7	0.40	0.46	0.48	0.51	0.54	0.59	0.64	0.80	0.88	0.96	1.07	1.12	1.61

R-4395 041320