



FIFRA 2(ee) Recommendation
EPA Reg. No. 279-3588
FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This label for Rhyme™ fungicide is valid until December 5, 2024, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Management of Botryosphaeria panicle and shoot blight and Phomopsis canker in Walnuts (including black and English (Persian) walnuts)

Directions For Use				
Apply <u>Rhyme</u> fungicide in sufficient water volume to ensure thorough coverage as a foliar application. Apply preventatively or when conditions are favorable for disease development.				
Crop	Disease	Rate (fl oz/A)	Application Directions	Retreatment Interval (Days)
Walnut	Botryosphaeria panicle and shoot blight Phomopsis canker	7	Apply preventatively and repeat as necessary on a 7-day interval when conditions are favorable for disease development.	7

RESTRICTIONS:

- No single application may exceed 7 fl oz of product/A.
- Do not apply more than 4 applications per year.
- Do not apply more than 28 fl oz of product/A per year.
- Do not apply more than 0.455 lb ai flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours
- Preharvest Interval (PHI) – Do not apply within 14 days of harvest

R-4357 120619 12-09-16