



FIFRA 2(ee) Recommendation
EPA Reg. No. 279-3588
FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This label for Rhyme™ fungicide is valid until December 5, 2024, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Management of Wood Disease Complex on Grapes and Tree Nuts

Directions for Use				
For management of wood diseases on grapes and tree nuts, apply Rhyme fungicide as directed through pressurized irrigation methods such as macro-sprinkler, micro-sprinkler, or drip systems. For best results, use Rhyme fungicide as part of a long-term management program.				
Crop	Disease	Rate (fl oz/A)	Application Directions	Retreatment Interval (Days)
<p>Grapes</p> <p>Established Vineyard (> 3 years old)</p>	<p>Vinewood diseases</p> <ul style="list-style-type: none"> <i>Botryosphaeria dothidea</i> <i>Eutypalata</i> Phomopsis canker <i>Phaeoacremonium parasiticum</i> <i>Neofusicoccum parvum</i> 	5	<p>Make initial application at the beginning of Spring Root Flush. Make second application 14 days later. Make third application at the start of Veraison. Make fourth application post-harvest during Fall Root Flush. All applications must be made through a pressurized irrigation system. To achieve maximum performance, use a minimum of 20 fl oz/A for the year</p>	14

Crop	Disease	Rate (fl oz/A)	Application Directions	Retreatment Interval (Days)
Grapes New Vineyards (1-3 years old)	Vinewood diseases <ul style="list-style-type: none"> <i>Botryosphaeria dothidea</i> <i>Eutypalata</i> Phomopsis canker <i>Phaeoacremonium parasiticum</i> <i>Neofusicoccum parvum</i> 	5	Make initial application at the beginning of Spring Root Flush. Make second application post-harvest during Fall Root Flush. All applications must be made through a pressurized irrigation system. To achieve maximum performance, use a minimum of 10 fl oz/A for the year	14
Tree Nuts (African Tree Nut, Almond, Brazil Nut, Burr Oak, Butternut, Cajou, Cashew, Castanha-Do-Maranhao, Coconut, Coquito Nut, Dika Nut, Guiana Chestnut, Hazelnut, Heartnut, Hickory Nut, Japanese Horse-Chestnut, Macadamia Nut, Monogongo Nut, Monkey-Pot, Pachira Nut, Pecan, Sapucaia Nut, and Walnut)	Trunk diseases <ul style="list-style-type: none"> <i>Botryosphaeria dothidea</i> <i>Cytospora sorbicola</i> <i>Neofusicoccum parvum</i> Phomopsis canker 	7	Make initial application of 7 fl oz/A at the beginning of Spring Root Flush, and reapply in 7 days. .Apply through a pressurized irrigation system. Reapply at the beginning of Fall Root Flush, and reapply in 7 days. To achieve maximum performance, make 4 applications at 7 oz/A. for the year	7

RESTRICTIONS:

Grapes

- No single application may exceed 5 fl oz of product/A.
- Do not apply more than 6 applications per year.
- Do not apply more than 30 fl oz of product/A per year.
- Do not apply more than 0.488 lb ai flutriafol/A per year.
- Restricted Entry Interval (REI) is 5 days for girdling or turning of grapes. For all other activities the REI is 12 hours.
- Preharvest Interval (PHI) – Do not apply within 14 days of harvest

Tree Nuts

- No single application may exceed 7 fl oz of product/A.
- Do not apply more than 4 applications per year.
- Do not apply more than 28 fl oz of product/A per year.
- Do not apply more than 0.455 lb ai flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours
- Preharvest Interval (PHI) – Do not apply within 14 days of harvest

R-4358 120619 12-09-19



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104