



## FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3588

**FOR DISTRIBUTION AND USE  
ONLY IN THE FOLLOWING STATES:  
AZ and NM**

**THIS RECOMMENDATION, WHICH CONTAINS  
ADDITIONAL DIRECTIONS FOR USE, IS MADE  
AS PERMITTED UNDER FIFRA SECTION 2(ee)  
AND HAS NOT BEEN SUBMITTED TO OR  
APPROVED BY THE US EPA.**

This recommendation for Rhyme™ Fungicide is valid until December 31, 2027, or until withdrawn, canceled or suspended.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.**

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

**FOR USE OF RHYME™ FUNGICIDE IN PECANS AS A SOIL DRENCH, DRIP, OR MICRO IRRIGATION SYSTEMS APPLICATION FOR CONTROL OF COTTON ROOT ROT (PHYMATOTRICHOPSIS OMNIVORA)**



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000

## Directions For Use

Read the label affixed to the container for Rhyme™ fungicide before applying

| Crop                                   | Disease   | Rate                                    | Application Directions  | Retreatment interval |
|--|---|---|---|----------------------|
| Pecans<br>( <i>Carya illinoensis</i> ) | Cotton Root Rot caused by ( <i>Phymatotrichopsis omnivora</i> ) | 7 fl oz/acre<br><br>(0.114 lbs ai/acre) | <b>Soil Drench:</b> To be made after the tree is planted. When Rhyme™ fungicide is applied as a soil drench, the application should be made from 2-3 feet around the base of the tree at concentrations not to exceed 7 fl oz/acre per application. The amount of fungicide to use per tree is based on tree and row spacing. | Minimum of 7 days    |
| Pecans<br>( <i>Carya illinoensis</i> ) | Cotton Root Rot caused by ( <i>Phymatotrichopsis omnivora</i> ) | 7 fl oz/acre<br><br>(0.114 lbs ai/acre) | <b>Drip or Micro-chemigation:</b> Rhyme™ fungicide may be applied through drip or micro chemigation systems where irrigation water configures around the trunk and root zone. The chemigation application can be made anytime after planting.   | Minimum of 7 days    |

### Restrictions

- No single application may exceed 7 fl. oz of product/acre
- Do not apply more than 4 applications per year
- Do not apply more than 28 fl. oz of product/acre per year
- Do not apply more than 0.455 lb. ai of flutriafol/acre per year
- Restricted Entry Interval (REI) is 12 hours

Do not apply within 7 days of harvest

R-4710 042622



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000